

CAVE MORPHOLOGY AND CONTROLS ON SPELEOGENESIS IN QUARTZITE: THE EXAMPLE OF THE ITAMBÉ DO MATO DENTRO AREA IN SOUTHEASTERN BRAZIL

MORFOLOGIJA IN SPELEOGENEZA JAM V KVARCITIH: PRIMER OBMOČJA ITAMBÉ DO MATO DENTRO V JOGOVZHODNI BRAZILIJU

Fabiana P. FABRI¹, Augusto S. AULER^{1*}, Allan S. CALUX², Roberto CASSIMIRO¹
& Cristina H. R. R. AUGUSTIN²

Abstract

UDC 551.435.84(81)

Fabiana P. Fabri, Augusto S. Auler, Allan S. Calux, Roberto Cassimiro & Cristina H. R. R. Augustin: Cave morphology and controls on speleogenesis in quartzite: The example of the Itambé do Mato Dentro area in southeastern Brazil

An area of 417 km² in southeastern Brazil was mapped in detail regarding karst features in quartzite. Landforms typically assigned to carbonate karst terrains are largely absent, except for discrete occurrences of towers and collapse dolines. Caves are the most conspicuous landform, although occurring at a lower density than in most Brazilian carbonate areas. A total of 11 caves were mapped and studied in detail. Caves tend to develop at shallow depths parallel to the slope surface, controlled by lithological facies and the hydraulic gradient. Petrographical analyses suggest that mica (sericite) lenses within the quartzite and both sericite and iron around quartz grains may favour grain disaggregation and later erosional removal in a shallow phreatic environment. Quartzite caves represent a common feature in southeastern Brazil but tend to be isolated features rather than an integrated and areally extensive hydrological system typical of many carbonate settings.

Key words: Quartzite caves, hydraulic gradient, sericite lenses.

Izveček

UDK 551.435.84(81)

Fabiana P. Fabri, Augusto S. Auler, Allan S. Calux, Roberto Cassimiro & Cristina H. R. R. Augustin: Morfologija in speleogeneza jam v kvarcilih: Primer območja Itambé do Mato Dentro v jugovzhodni Braziliji

V članku opisujemo rezultate podrobnega kartiranja krasa v kvarcitu na 417 km² velikem območju v jugovzhodni Braziliji. Z izjemo stolpov in udornic na območju ni površinskih kraških oblik, značilnih za karbonatni kras. Jame so najbolj značilen kraški pojav, čeprav je njihova gostota precej manjša od tiste na karbonatnem krasu. Na celotnem območju smo izmerili in detajlno raziskali 11 jam, ki so razvite plitvo in vzporedno s površjem. Glavna dejavnika razvoja jam sta litologija in hidravlični gradient. Petrografske analize kažejo, da je razvoj jam najbolj aktiven v območjih plitve freatične cone, kjer se v kvarcitu pojavljajo leče sljude (sericit) ter kjer sericit in železo obdajata kremenčeva zrna. V the območjih prihaja do razpadanja in mehanskega odnašanja zrn. V jugovzhodni Braziliji so kvarcitrne jame pogost pojav, vendar so med seboj pravilpoma nepovezane, kar kaže na to, da v primerjavi s karbonatnim krasom v kvarcitu ne prihaja do razvoja velikih povezanih kraških drenažnih sistemov.

Ključne besede: Jame v kvarcitu, hidravlični gradient, sericitne leče.

INTRODUCTION

Quartzite caves are now recognised as relatively common features in many tropical areas of the world (Wray 1997a; Auler 2012; Wray 2013). Although studies have been

performed since the early 1960s (White *et al.* 1966), systematic research on this topic is rather recent, with new information being produced every year. A significant

¹ Instituto do Carste, Rua Aquiles Lobo, 297, Belo Horizonte, MG, 30150-160, Brazil, aauler@gmail.com

² Departamento de Geografia, Instituto de Geociências, Universidade Federal de Minas Gerais, Avenida Antonio Carlos 6627, Belo Horizonte, MG, 31270-901, Brazil

*Corresponding author

Received/Prejeto: 01.12.2013

portion of new data comes from the tepui area of southern Venezuela, where the longest and largest quartzite caves in the world have recently been discovered (Aubrecht *et al.* 2013; Mecchia *et al.* 2014).

Genesis of quartzite caves has traditionally been interpreted in terms of an early dissolutional phase (termed “arenisation”) in which quartz grains boundaries are dissolved, increasing the porosity and resulting in a porous, friable rock that is subjected to later erosional removal of quartz grains (Martini 1979). The existence of dissolution processes is demonstrated by the frequent occurrence of silica speleothems, mostly coralloids (Wray 2007, 2009; Aubrecht *et al.* 2008), with opal being the dominant mineral. Hydrochemical studies (e.g., Piccini & Mecchia 2009; Mecchia *et al.* 2014) have supported a quantitatively restricted role of dissolution, with low SiO₂ content in most quartzite cave waters. However, given the long term geomorphic evolution of the area, even low levels of silica can be relevant to arenisation. The role of organic matter and microorganisms in trig-

gering quartz dissolution has also been suggested (Bennet 1991; Barton *et al.* 2009). The relative importance of an initial dissolutional phase is subject to debate (Sauro *et al.* 2013; Aubrecht *et al.* 2013), although there appears to be consensus that the morphology and evolution of quartzite cave systems are largely due to a later prolonged erosional phase. Indeed, some authors have recently argued that an early dissolutional phase may not be needed to explain the genesis of large quartzite cave systems (Aubrecht *et al.* 2011). Despite the recent progress in understanding the processes that control quartzite cave genesis, there is a lack of systematic research compared to the better studied carbonate areas.

Controls on quartzite cave development were examined through a study of a quartzite area in southeastern Brazil. Cave mapping, structural and petrographical studies were performed to investigate the cave inception processes and the relationship between the caves and their surrounding landscape.

STUDY AREA

The study area is located in the eastern state of Minas Gerais, southeastern Brazil (Fig. 1), mostly within the municipality of Itambé do Mato Dentro. This area lies in the southernmost expression of the vast Espinhaço Ridge, which extends over 1,500 km towards the north. Due to the need to perform detailed geomorphological work, a

more restricted rectangle of 417 km² was chosen, encompassing all significant caves.

Quartzite in the area belongs to three Units of the Sopa Brumadinho Formation of the Mid-Proterozoic Espinhaço Supergroup (Grossi Sad *et al.* 1997) (Fig. 2). The lower Serra do Lobo Unit is comprised of fine to coarse




Fig. 1: Location of the study area.




Fig. 3: Hypsometric map, fluvial network, location of caves and regional profiles.

METHODS

A previous speleological study (Cruz *et al.* 1999) indicated the presence of quartzite caves in the Cabeça de Boi Ridge. Further prospective work detected additional quartzite caves, identified mostly through information from local farmers. The study area was thus defined based on the location of caves and the presence of major ridges (Cabeça de Boi, Lobo, Linhares, Lapa and Espinhaço ridges).

The caves were mapped using conventional cave survey equipment (Suunto compasses and clinometers) and Leica Disto laser tapes. Survey grades reached BCRA grade 4C. The morphology of each cave was examined through cave maps and measurements at individual cave passages. Rock samples were collected at selected locations and analysed through thin sections, mineralogy being determined through a XPERT-PRO X-ray diffractometer. Structural data were measured at cave passages

using a Brunton Geo compass and processed through StereoNet 2.10 software.

Five regional topographic profiles were produced, together with local profiles along the main axis of cave development. These profiles were produced with the aid of Global Mapper 10.02 software, complemented with field measurements. Regional profiles were used to obtain the regional relief gradient. Map analyses of those slopes containing caves were used to infer the local relief gradient through the relation:

$$[(A_{max} - A_{min}) / D] \times 100 \text{ where:}$$

A_{max} – Maximum altitude of the profile
 A_{min} – Minimum altitude of the profile
 D – Distance between A_{max} and A_{min}

The cave drainage gradients were inferred through profiles of stream passages. The remaining quantitative morphometric cave data were obtained through measurements on cave maps using AutoCAD software.

KARST GEOMORPHOLOGY

Karst-like surface forms in the study area are largely absent, as observed in many quartzite terrains in Brazil. A few closed depressions were identified, some associated with cave collapse or elongated parallel to joints (Fig. 4a). Scarp dissection leads to individualisation of towers (Fig. 4b) that resemble limestone karst towers but do not necessarily imply dissolutional processes. Smaller features such as honeycomb weathering (Mustoe 1982) also occur.

UNDERGROUND LANDFORMS

A total of 11 caves were identified and studied. Their location is presented in Fig. 3. Tab. 1 presents general speleometric data. Most caves are rather small, with the notable exception of the Baixada das Crioulas I and II caves, which represented a single major (> 1 km long) cave separated by a short breakdown canyon. These fragmentary caves (approximately 40 % of the caves are less than 100 m long) pose difficulties to the interpretation




Fig. 4: A- Regional view of the study area showing a large closed depression (see arrow). B- Karst-like residual towers.

Tab. 1: Speleometric data for the studied caves. Cave density relates to the ratio between the cave area and the smallest rectangle that contains the entire cave. Sinuosity is represented by the ratio between the fluvial channel meandering length and the straight line distance between ends. * Measurement not possible due to cave pattern.

Caves	UTM Coordinates		Length (m)	Depth (m)	Area (m ²)	Volume (m ³)	Density	Maximum distance between extremes	Sinuosity	Number of entrances
	N	S								
Abrigo das Pinturas	7849576	683355	65	8	518	1677	0.156	65	*	1
Baixada das Crioulas I	7853631	672862	1074	75	6659.6	27572	0.05	517.13	1.12	10
Baixada das Crioulas II	7853583	673059	205	13.5	1776.7	18510	0.19	163.77	1.04	3
Braúna Seca	7852893	665102	37	7	193	611	0.386	22.5	1.03	2
Esteira	7848930	683629	96	28	335	1461	0.142	68.55	1.01	4
Funil I	7849054	679690	21	4	141	595	1,433	21	1.01	1
Funil II	7849054	679690	106	29	534	618	0.0945	105.22	1.01	1
Gentio	7848947	679991	14	2	53	166	0.571	13.68	1.03	1
Milagres I	7849656	683048	225	14	1100.3	3344	0.0586	125.72		3
Milagres II	7850686	675724	149	17	806	1136	0.113	143.62	1.01	2
Toquinha	7848930	683629	6	1.5	9.3	4.36	0.153	6	1	1

Tab. 2: Regional, local and stream passage gradients for the studied caves.

Caves	Lithology	Regional relief gradient (%)	Local relief gradient (%)	Drainage network gradient(%)	Cave length (m)
Abrigo das Pinturas	Itambé do Mato Dentro Unit. Medium to fine granulometry.	3 – 8	10.42	5.78	65
Baixada das Crioulas I	Serra do Lobo e Rio Preto Units. Medium to fine granulometry, with sericite films.	8 – 20	10.65	10.02	1074
Baixada das Crioulas II	Rio Preto Unit. Medium to fine granulometry, with sericite films.	8 – 20	10.65	10.02	205
Braúna Seca	Rio Peto Unit (phosphate lenses). Medium to fine granulometry..	8 – 20	42.96	30.02	37
Esteira	Itambé do Mato Dentro Unit. Medium to fine granulometry, with sericite films.	20 – 45	27.4	17.78	96
Funil I	Itambé do Mato Dentro Unit. Medium to fine granulometry, with sericite films.	8 – 20	14.32	22.9	21
Funil II	Itambé do Mato Dentro Unit. Medium to fine granulometry.	8 – 20	14.32	22.9	106
Gentio	Itambé do Mato Dentro Unit. Medium to fine granulometry.	8 – 20	14.32	22.9	14
Milagres I	Itambé do Mato Dentro Unit. Medium to fine granulometry, with sericite films.	3 – 8	8.28	5.78	225
Milagres II	Itambé do Mato Dentro Unit. Medium to fine granulometry.	8 – 20	17.78	9.14	149
Toquinha	Itambé do Mato Dentro Unit. Medium to fine granulometry.	8 – 20	31.76	23.56	6

of large scale morphology and genesis because they are not long enough to allow for a full display of the cave pattern.

Most caves occur as isolated landforms, resulting in a density of 0.026 caves/km². Caves develop mostly in the less resistant quartzite of the Rio Preto and Itambé do Mato Dentro Units. No correlation was found between slope gradient and cave development, and no caves were located in areas of very high gradient; the majority of caves occur in slopes of 8 – 20 % (Tab. 2 and Fig. 5).

The longer caves tend to contain a perennial stream, displaying a linear plan pattern with few or no tributaries (Fig. 6). Values for sinuosity are close to 1 (Tab. 1). The major caves tend to follow the slope gradient, being characterised by an inclined passage that frequently represents a link between upper (inlet) and lower (outlet) entrances. The majority of the caves possess less than 3 entrances, while most of the shorter ones displays a single entrance. An important exception is represented by Baixada das Crioulas I and II caves because they contain numerous entrances (Fig. 6) associated with inlets at the western/northwestern limit, possibly associated with the presence of a nearby scarp that favoured breakdown and surface water input to the caves.

Cave walls exhibit friable quartzite that precludes preservation of possible original hydrodynamic forms.

Irregular pillars occur in many caves. Cross sections are, on average, at most a few meters in terms of height and width. However, sections show considerable variation. Caves that contain streams (Baixada das Crioulas, Funil, Milagres, Esteira) show passages with a width/height ratio of approximately 2 (Fig. 7) and are square or lens-shaped. Keyhole passages have also been identified in the area. Joint-controlled rift passages occur in selected caves (Milagres).

The relatively homogeneous character of the geology surrounding the caves, comprised of mostly quartzite, results in an equally monotonous sedimentation inside caves. Sand residues are by far the commonest sediment, both of autogenic (crumbling breakdown of friable walls/ceiling) and allogenic sources (Fig. 8). Fine-grained clay and silt are observed at a few sites. Rare deposits of poorly sorted coarser and sometimes brecciated sediments have been recorded in some caves (Baixada das Crioulas, Funil II), usually related to paleo-fluvial terraces (Fig. 9).

Chemical sedimentation (speleothems) is limited mostly to coralloids and small stalactites. X-ray diffractometry has shown that speleothems are composed of quartz and opal-A, similar to the findings from sandstone and quartzite caves elsewhere (Wray 1997b; 1999; 2013; Wiegand *et al.* 2004; Aubrecht *et al.* 2008). Coralloids are not commonly associated to joints; this finding




Fig. 5: Selected regional profiles showing geology and cave locations. Profile locations in Fig. 3.




Fig. 6: Cave pattern in plan view of some of the studied caves. A - Baixada das Crioulas I and II caves. B - Esteira cave. C - Milagres II cave. D - Funil I cave. E - Milagres I cave. E - (Entrance)



Fig. 7: Lens-shaped passage with small streams at the wall/floor contact at Funil II cave. Fine-grained sedimentation at the centre of the passage.

indicates water input through rock porosity, as suggested by Wray (1999) in Australian caves and Mecchia *et al.* (2014) in the Venezuelan tepui caves. Thin section of stalactites allows the observation of an irregular display



Fig. 8: Sand residue and quartzite breakdown at Milagres I cave.

of opal layers parallel to a central axis. No evidence of an internal channel has been found, as noted by Wray (1997b). Another type of speleothem is represented by colloidal dark reddish brown flowstone, with crenulated surfaces that resemble millimetre wide rimstone dams. This type of deposit has been previously interpreted by Corrêa Neto *et al.* (1997) as expansive clays with allo-




Fig. 9: Alluvial coarse grained terrace at Baixada das Crioulas II cave. A- General view. B- Close-up view.

phane and amorphous bauxite, derived from the weathering of aluminium silicates. X-ray diffractometry analy-

ses indicate that quartz intermixed with clay material of undetermined mineralogy.

CONTROLS ON CAVE GENESIS

Approximately 800 measurements of joints, bedding and foliation planes were performed inside caves (Fig. 10). The most prominent joint direction is N0-20E, not closely correlated to cave passage orientation, which displays a more prominent direction at N80-90E. When passage type is taken into account, stream caves show limited

control by joints, joint direction correlation being found mostly in rift passages in dry caves, such as Milagres cave. Passage direction appears to be controlled by the dip direction of both bedding plane and foliation, showing these structural elements may be of importance in determining the position of passage initiation.

Baixada das Crioulas I and II caves exhibit a curious curvilinear pattern that is not found elsewhere in the area (Fig. 6). The cave passage closely follows a surface valley, suggesting a similar control for both surface and underground water flow.

The relationship between caves and the hydraulic gradient was analysed. Cave profiles, as previously mentioned, closely mirror the slope gradients (Fig. 11). Tab. 2 shows slope gradients for the caves, both in a regional and local (in the scale of the cave) context. There is no clear correlation between slope gradient and cave length, although all caves (with the exception of the horizontal Milagres I cave) develop within gradients $>10^\circ$. Furthermore, the caves appear to be aligned with the steepest slope gradient. Cave development appears to follow a favourable horizon whenever quartzite dip direction is aligned with the hydraulic gradient. Most caves develop in such conditions, being less conspicuous in horizontal rock/relief settings.




Fig. 10: Stereonet directions of cave passages, fractures, bedding plane and foliation for measurements in all caves.




Fig. 11: Relationship between cave and slope profiles, showing that caves tend to develop close to the surface. A- Funil I, II and Gento caves. B- Milagres II cave. C- Baixada das Crioulas I and II caves.

Quartzite samples taken at Funil II, Milagres I and II and Baixada das Crioulas I caves were subject to petrographical and geochemical analyses. They are comprised of fine to medium-grained quartzite with a high (>85%) content of quartz but with significant amounts

(5–10 %) of sericite. Iron oxides/hydroxides also occur associated with sericite. Sericite tends to occur as films associated with the foliation (Fig. 12) but also enveloping quartz grains. Sampling in both ceiling and pillars demonstrated that the residual pillars are characterised




Fig. 12: Sericite (see arrow) occur as continuous lenses that determines the rock foliation. Photomicrography from a sample from Baixada das Crioulas I cave.

by a lower (5–10 %) amount of sericite, while higher amounts (15–20 %) were recorded in the ceiling. Lower levels of sericite may thus result in more chemically resistant rock, suggesting that sericite/iron content may

play a major role in cave initiation in the area. One of the largest volumes among the studied caves is related to a mica-rich zone where the “big chamber” of Milagres I cave develops.

DISCUSSION

Caves in the Itambé do Mato Dentro area share many characteristics observed in quartzite caves elsewhere in Brazil (Corrêa Neto 2000; Wiegand *et al.* 2004; Auler 2012). Genesis of quartzite caves is traditionally attributed to dissolution along the boundaries of quartz grains, allowing for later erosional removal of constituents (Martini 1979; Wray 1997a, 2013). The role of dissolution in quartzite cave development is usually considered to be restricted to the initial phases of speleogenesis, being considerably small when compared to the later erosional phases. Some authors have even claimed that a dissolutional phase is not required to explain some of the largest and longest quartzite caves in the world (Aubrecht *et al.* 2011). In the study area, however, there is clear evidence of chemical alteration of an original, very hard and crystalline rock as observed in outcrops, resulting in a friable water-soaked rock that is easily removed under the existing hydraulic gradient.

Studies performed in Brazilian quartzite caves have stressed the importance of more readily weathered constituents in providing the initial route for ground water flow. These more favourable horizons (termed “inception horizons” by Lowe (1992)) can represent *loci* for the initial development of caves. Corrêa Neto (2000) first suggested that the alteration of micas, feldspars and clay

minerals may be an important mechanism in cave genesis at Ibitipoca ridge, southeastern Brazil. Melo & Gianini (2007) argue that kaolinite dissolution is the main trigger in the formation of karst landforms in the Furnas Formation of southern Brazil. In the Venezuelan tepuis, lenses of iron hydroxide or silt layers have been considered as possible inception horizons (Sauro 2014).

Bedding planes and foliation are the most common controlling structures on passage orientation in the study area. The existence of sericite layers precisely along the same horizons, as shown by petrographical analysis, may present a favourable condition for cave development, especially when the dip is in accordance with the hydraulic gradient. Furthermore, both sericite and iron oxides also occur alongside quartz grains. The weathering of mica releases both K and Mg, which may accelerate the dissolution of amorphous silica and SiO_2 (Icenhower & Dove 2000). Micas and clay minerals can also be important players in the dissolution of quartz in sandstone (Bjørkum *et al.* 1996; Meyer *et al.* 2006). Laterisation processes due to alteration of micas, feldspar and clay minerals may account for considerable generation of porosity (Corrêa Neto 2000; Aubrecht *et al.* 2011) and favour arenisation along or around specific beds (Mecchia *et al.* 2014). Iron can also be significantly mobilised through various pro-

cesses involving microbiological agents, contributing to porosity generation (Chalcraft & Pye 1984; Striebel & Schaferjohann 1997; Barton 2013; Sauro 2014).

The studied caves develop at a shallow depth (less than 50 m below the surface) and show a general concordance with the slope gradient. Although no direct

relationship between cave length and slope gradient was recorded in the area, optimal conditions for quartz removal appear to occur when both hydraulic gradient and bedding/foliation planes (containing sericite/iron) are aligned.

CONCLUSIONS

A study of 11 quartzite caves in southeastern Brazil has shown that caves are not a conspicuous geomorphological landform, but occur as isolated shallow features with a density of 0.026 caves/km², lower than in most carbonate karst areas in Brazil. Regional and local profiles show that caves develop parallel to the surface terrain and are not linked to deep groundwater flow paths. Bedding and foliation planes are the controlling structures in guiding the initial phase of speleogenesis. Thin sections demon-

strate that sericite layers occur aligned with the bedding planes and iron oxides/hydroxides are common around quartz grains. Weathering processes associated with both iron and sericite tend to free quartz grains (arenisation), allowing for the erosional removal of quartz, as demonstrated in other quartzite areas of the world. Although quantitatively small, chemical alteration of phyllosilicates is of paramount importance in defining the early circulation paths in an essentially shallow phreatic setting.

ACKNOWLEDGMENTS

Financial support was provided by FAPEMIG (grant CRA APQ 01652-09). We acknowledge the contribution of Instituto do Carste in helping organise cave surveys and data collection in the field. Régis de Paula Reis,

Tatiana Souza and the local guide Zequita helped with field work. Francesco Sauro and an anonymous reviewer provided insightful suggestions that helped improve the manuscript.

REFERENCES

- Aubrecht, R., Láncoz, T., Smida, B., Brewer-Carias, C., Mayoral, F., Schlogl, J., Audy, M., Vlcek, L., Kovacik, L. & M. Gregor, 2008: Venezuelan sandstone caves: a new view on their genesis, hydrogeology and speleothems.- *Geol. Croat*, 61, 345–362.
- Aubrecht, R., Láncoz, T., Gregor, M., Schlogl, J., Smida, B., Liscak, P., Brewer-Carias, C. & L. Vlcek, 2011: Sandstone caves on Venezuelan tepuis: Return to pseudokarst?- *Geomorphology*, 132, 351–365.
- Aubrecht, R., Láncoz, T., Gregor, M., Schlogl, J., Smida, B., Liscak, P., Brewer-Carias, C. & L. Vlcek, 2013: Reply to the comment on "Sandstone caves on Venezuelan tepuis: Return to pseudokarst?- *Geomorphology*, 197, 197–203.
- Auler, A.S., 2012: Quartzite caves of South America.- In: White, W.B. & D.C. Culver. (eds.) *Encyclopedia of Caves*. Academic Press, pp. 635–639, Chennai.
- Bennet, P.C., 1991: Quartz dissolution in organic rich aqueous systems.- *Geochimica et Cosmochimica Acta*, 55, 1781–1797.
- Barton, H.A., 2013. Biospeleogenesis.- In: Frumkin, A. (ed.) *Treatise on Geomorphology* Academic Press, 6, pp. 38–56, San Diego.
- Barton, H., Suarez, P., Muench, B., Giarrizzo, J., Broering, M., Banks, E. & K. Venkateswaran, 2009: The alkali speleogenesis of Roraima Sur Cave, Venezuela.- In: *Proceedings 15th International Congress of Speleology*, 19th–26th June 2009, Kerrville. National Speleological Society, 802–807.

- Björkum, P.A., 1996: How important is pressure in causing dissolution of quartz in sandstones?- *Journal of Sedimentary Research*, 66, 147–154.
- Chalcraft, D. & K. Pye, 1984: Humid tropical weathering of quartzite in southeastern Venezuela.- *Zeitschrift für Geomorphologie*, 28, 321–332.
- COMIG., 1996: Geological map of Conceição do Mato Dentro sheet.- Projeto Espinhaço 1:100.000.
- Corrêa Neto, A.V. & G. Dutra, 1997: A província espeleológica quartzítica Andrelândia, sudeste de Minas Gerais.- In: *Proceedings XXIV Brazilian Congress of Speleology*, 57–64.
- Corrêa Neto, A.V., 2000: Speleogenesis in quartzites in southeastern Minas Gerais, Brazil.- In: Klimchouk, A. et al. (ed.). *Speleogenesis. Evolution of Karst Aquifers*. National Speleological Society, pp. 452–457, Huntsville.
- Cruz, L.V., Pereira, M., Lopes, M.V.C.O. & R.C. Mourão, 1999: Avaliação espeleológica da Serra Cabeça de Boi, Fazenda Ponte – Itambé do Mato Dentro – MG.- *Proceedings XXV Brazilian Congress of Speleology*, 27–39.
- Grossi Sad, J.H., Mourão, M.A.A., Guimarães, M.L.V. & L.G. Knauer, 1997: Geologia da Folha Conceição do Mato Dentro, Minas Gerais. *Projeto Espinhaço*: CD-Rom.
- Icenhower, J. & P. Dove, 2000: The dissolution kinetics of amorphous silica into sodium chloride solutions: Effects of temperature and ionic strength.- *Geochimica et Cosmochimica Acta*, 64, 4193–4203.
- Lowe, D.J., 1992: The origin of limestone caverns: An inception horizon hypothesis.- PhD thesis, Manchester Metropolitan University, 512pp.
- Martini, J., 1979: Karst in the Black Reef Quartzite near Kaapsehoop, Transvaal.- *Annals Geological Survey of South Africa*, 13, 115–128.
- Mecchia, M. & L. Piccini, 1999: Hydrogeology and SiO₂ geochemistry of the Aonda Cave System, Ayuan-Tepui, Bolivar, Venezuela.- *Boletín de la Sociedad Venezolana de Espeleología*, 33, 1–11.
- Mecchia, M., Sauro, F., Piccini, L., De Waele, J., Sanna, L., Tisato, N., Lira, J. & F. Vergara, 2014: Geochemistry of surface and subsurface waters in quartz-sandstones: significance for the geomorphic evolution of tepui table mountains (Gran Sabana, Venezuela).- *Journal of Hydrology*, 511, 117–138.
- Melo, M.S. & P.C.F. Giannini, 2007: Sandstone dissolution landforms in the Furnas Formation, Southern Brazil.- *Earth Surface Processes and Landforms*, 32, 2149–2164.
- Meyer, E.E., Greene, G.W., Alcantar, N.A., Israelachvili, J.N. & J.R. Boles, 2006: Experimental investigation of the dissolution of quartz by a muscovite mica surface: Implications for pressure solution.- *Journal of Geophysical Research*, 111, 1–4.
- Mustoe, G.E., 1982: The origin of honeycomb weathering.- *Geological Society of America Bulletin*, 93, 108–115.
- Piccini, L. & M. Mecchia, 2009: Solution weathering rate and origin of karst landforms and caves in the quartzite of Ayuan-tepui (Gran Sabana, Venezuela).- *Geomorphology*, 106, 15–25.
- Sauro, F., Piccini, L., Mecchia, M. & J. De Waele, 2013: Comment on “Sandstone caves on Venezuelan tepuis: Return to pseudokarst?” by R. Aubrecht, T. Láncoz, M. Gregor, J. Schlögl, B. Smída, P. Liscák, Ch. Brewer-Carías, L. Vlcek, *Geomorphology* 132 (2011), 351–365.- *Geomorphology*, 197, 190–196.
- Sauro, F., 2014: Structural and lithological guidance on speleogenesis in quartz-sandstone: Evidence of the arenisation process.- *Geomorphology*, 226, 106–123.
- Striebel, T. & V. Schaferjohann, 1997: Karstification of sandstone in Central Europe: Attempts to validate chemical solution by analyses of water and precipitates. *Proceedings 12th International Congress of Speleology*, 10th–17th August 1997, La Chaux des Fonds. Swiss Speleological Society, 473–476.
- White, W.B., Jefferson, G. L. & J.F. Haman, 1966: Quartzite karst in southeastern Venezuela.- *International Journal of Speleology*, 2, 309–314.
- Wiegand, J., Fey, M., Haus, N. & I. Karmann, 2004: Geochemische und hydrochemische untersuchungen zur genese von sandstein-und quarzitkarst in der Chapada Diamantina und im eisernen viereck (Brasilien).- *Z. dt. Geol. Ges.*, 155, 61–90.
- Wray, R.A.L., 1997a: A global review of solutional weathering froms on quartz sandstones.- *Earth Sciences Reviews*, 42, 137–160.
- Wray, R.A.L., 1997b: The formation and significance of coralline silica speleothems in the Sydney Basin, southeastern Australia.- *Physical Geography*, 18, 1–17.
- Wray, R.A.L., 1999: Opal and chalcedony speleothems on quartz sandstone in the Sydney region, southeastern Australia.- *Australian Journal of Earth Sciences*, 46, 623–632.
- Wray, R.A.L., 2013: Solutional weathering and karstic landscapes on quartz sandstones and quartzite.- In: Frumkin, A. (ed) *Treatise on Geomorphology*. Vol. 6. Karst Geomorphology. Elsevier, pp. 463–483, San Diego.