

Identification of Best Growth Curve Model for Anatolian Black Cattle

Key words

Anatolian Black cattle;
live weight;
growth curve;
non-linear models

Çağrı Melikşah Sakar^{1*}, Seyrani Koncagül², İlker Ünal¹

¹International Center for Livestock Research and Training, Mamak, Ankara, ²Ankara University, Faculty of Agriculture, The Department of Animal Science, Ankara, Turkey

*Corresponding author: koncagul@ankara.edu.tr

Abstract: The aim of this study was to identify the model that best describes the growth trajectory from birth to 24 months of age in Anatolian Black Cattle (ABC) raised for conservation purposes. A total of 493 weight records of 113 animals at birth, 3, 6, 12, 18, and 24 months were collected. Six different non-linear models were used to describe the growth curve of animals: 2nd degree polynomial, 3rd degree polynomial, Logistic, Brody, Von Bertalanffy, and Gompertz models. In the study, R² values of the models were: 0.997, 0.999, 0.953, 0.979, 0.924, and 0.862; corel values (correlation between the observed and estimated curves) were 0.994, 0.998, 0.989, 0.993, 0.961, and 0.703; Residual Standard Deviations (RSD) were 3.216, 1.388, 11.533, 3.561, 14.736, and 27.141, respectively. Given these values, it was found that the 3rd degree polynomial model was the best to describe the growth curve of ABC. As a result of the analyses, it was noticed that the values predicted by this model deviated by 1-3 kg from the observed values in all periods and in all environmental factors examined (sex, dam age, parity, birth year and birth, season). It was found that these differences increased up to 4-5 kg only in the 18-month period. The results also showed that ABC continued to grow after 24 months of age. As a result, traits such as age at sexual maturity, breeding age, and slaughter age can be easily predicted by identifying the model that best describes growth and development in herds.

Received: 28 January 2023
Accepted: 12 September 2023

Introduction

Cattle breeding has an important place in Turkey's animal husbandry; while there are around 18 million cattle in Turkey, approximately 8% of this is local breeds (1). Domestic cattle breeding is important in terms of rural employment and development, rural sociology and the use of poor pasture areas. Anatolian Black Cattle (ABC) is one of the breeds that is most widely grown and spread among the domestic breeds in Turkey. This breed is grown especially in the Central Anatolian Region of Turkey and is mostly grown for meat and milk yield by breeders living in rural areas. It has adapted to these conditions since it has grown in unfavourable conditions in this region for many years. They have gained resistance to harsh winters, drought, hunger, thirst, and diseases (2, 3). Growth and development values in Turkey's domestic cattle breeds are generally slower than those of developed cattle breeds. In studies conducted with

the ABC breed in Turkey, live weights at birth, 3, 6 and 12 months of age were found to be 14.85 kg, 49.37 kg, 81.22 kg, and 97.29 kg, respectively (2). In other studies with the same breed, live weights from birth to 12 months of age were the following: 16.97- 21.35 kg, 63.21 - 68.18 kg, 101.04 - 110.33 kg, and 152.16 kg- 184.57 kg, respectively (3, 4, 5).

Growth in cattle is a function that continues throughout the life of the animal, from embryonic stages to adulthood, and can be explained mathematically by growth curve models (6). The change in any of the examined features over a certain period is defined as the growth curve (7). The growth curve shows the statistical relationship between the weight and time or age of animals, which is shaped under the influence of genetic potential and environmental factors (8). The growth of living organisms does not progress at a



Figure 1a: Anatolian Black cows and calves



Figure 1b: Animal weighing

constant rate throughout their lives (9). In the case of constant growth, linear models are used, and when the growth rate occurs at different times depending on age, nonlinear models (such as Negative Exponential, Brody, Logistics, Gompertz, Bertalanffy, Richards, and Weibull) are used (6, 10). The fact that living things have different growth rates in some periods necessitated the use of nonlinear models, which were more comprehensive models (8). There is still a need to investigate whether the most commonly preferred non-linear models are sensitive to the length of the growth period prior to truncation of the data (11).

The aim of this study was to evaluate non-linear models of the growth curve in ABC cattle taken at individual weights from birth to 24 months and to determine the model that best explains growth. For this purpose, 2nd degree polynomial, 3rd degree polynomial, Logistic, Brody, Von Bertalanffy, and Gompertz models were analysed.

Materials and methods

Animals

The animal material of this study consisted of Anatolian Black Cattle (ABC) grown in the “International Center for Livestock Research and Training” (39°97' N, 33°10' E; elevation 826 m) located in Ankara. This breed has been conserved within the scope of the project “Conservation of Domestic Genetic Resources and Sustainable Use” conducted by the General Directorate of Agriculture Research and Policies. The study was carried out on a total of 113 heads of ABC born between 2015 and 2020.

ABC calves were raised with their dams from birth, and they were allowed to suckle their dams freely. The cows were not milked on the farm. Feeding of ABC bred cows was two meals a day, morning and evening, ad libitum in the form of total mixed feed. ABC cows were given 80% barley bales and 20% dry meadow grass as roughage.

Table 1: Descriptive statistics of body weight at different ages in Anatolian Black Cattle

Statistics	BW	3MW	6MW	12MW	18MW	24MW
N	113	93	96	98	35	58
Minimum (kg)	13.00	37.00	52.50	89.00	144.00	178.00
Maximum (kg)	30.00	99.00	153.00	283.00	332.00	444.00
Female (kg)	17.25	61.70	90.42	144.30	188.65	225.34
Male (kg)	19.55	67.66	103.08	162.75	245.50	303.90
Mean (kg)	18.57	65.10	98.33	155.60	217.89	264.64
Standard Error	0.312	1.340	2.120	3.830	7.970	7.910
Coefficient of Variation (CV%)	17.86	19.88	21.17	24.40	21.64	22.78

Notes: BW=birth weight, 3MW=3 month weight, 6MW=6 month weight, 12MW=12 month weight, 18MW=18 month weight, 24MW=24 month weight.

Table 2: Non-linear models used to describe the growth of Anatolian Black Cattle

Model	Equation	Reference
2 nd Degree Pol.	$y_t = \beta_0 + \beta_1 t + \beta_2 t^2$	12
3 rd Degree Pol.	$y_t = \beta_0 + \beta_1 t + \beta_2 t^2 + \beta_3 t^3$	12
Logistic	$y_t = A (1 + be^{-kt})^{-1}$	13
Brody	$y_t = A (1 - be^{-kt})$	14
Von Bertalanffy	$y_t = A (1 - be^{-kt})^3$	15
Gompertz	$y_t = A \exp(-be^{-kt})$	16

y_t = observed BW at age t (kg); $\beta_0, \beta_1, \beta_2, \beta_3$: regression coefficients of 2nd and 3rd degree polynomial models; A: the asymptotic limit of the BW when age t approaches infinity (kg); b: the integration constant, related to the initial weights of the animal and without a well-defined biological interpretation; k: ratio of the relative intensity of growth (maturation rate); t: time (month).

Figure 1a shows Anatolian Black cows and calves, while Figure 1b shows the weighing of an animal.

Data set

In this study, birth weight, 3, 6, 12, 18, and 24 month live weights of 113 calves born between 2015 and 2020 were used. These values were determined by weighing them with precision scales up to 200 g. The characteristics of the data are presented in Table 1. In addition, information on sex, dam age, parity, birth year, and month was also recorded.

Table 3: Model comparison for growth of Anatolian Black Cattle

Model	β_0	β_1	β_2	β_3	R ²	RSD	corel
2 nd degree polynomial	21.80±0.619	0.44±0.0132	0.01±0.001	-	0.997	3.216	0.994
3 rd degree polynomial	18.72±0.361	0.33±0.099	0.01±0.002	0.01±0.001	0.999	1.388	0.998
	A	b	k	t ₁			
Logistic	206.09±8.417	-	-1.79±1.788	8.84±8.839	0.953	11.533	0.989
Brody	225.56±15.625	0.83±0.027	0.01±0.001	-	0.979	3.561	0.993
Von Bertalanffy	169.96±15.650	-0.19±0.124	0.01±0.005	-	0.924	14.736	0.961
Gompertz	129.05±4.530	2.751±0.079	0.40±0.000	-	0.862	27.141	0.703

$\beta_0, \beta_1, \beta_2, \beta_3$: regression coefficients of 2nd and 3rd degree polynomial models; A: the asymptotic limit of the BW when age t approaches infinity (kg); b: the Integration constant, related to the initial weights of the animal and without a well-defined biological interpretation; k: ratio of the relative intensity of growth (maturation rate); t: time (month).

R²: coefficient of determination; RSD: Residual Standard Deviation; corel: correlation between observed and estimated growth curves

Predicting the growth curve

In the study, six different non-linear models were used in the estimation of growth curves, and these models are presented in Table 2.

In the study, R² (coefficient of determination), RSD (Residual Standard Deviation), and corel (correlation) between the observed and estimated growth curves were used to compare the models.

Statistical analysis

Statistical analyses were carried out using the Proc Nlin in SAS (17). Growth curve models were fitted for each animal separately, and then the best-fitted model parameters and the other phenotypic data were analyzed using the Proc Glm in SAS (17). For analysis, sex (female, male), age of dam (2-3, 4-7, 8-10, 11+), parity (1, ...7), birth year (2015, ... 2020), and birth season (winter, spring, summer, autumn) were included in the model, and these were taken as fixed effects in the GLM analyses. To determine the differences between groups, the Tukey test was used.

Results and discussion

The growth curve parameters as derived from 6 different models using 493 weight records for ABC are presented in Table 3. The 3rd degree polynomial model showed the highest R² and corel and lowest RSD, indicating the best goodness of fit. On the other hand, the Gompertz model was the least fitted to estimate the ABC weight based on its lowest value of R². That is, the 3rd degree polynomial model with four parameters ($\beta_0, \beta_1, \beta_2, \beta_3$), which has the highest R² and smallest RSD value, best explains the change in live weight




Figure 2: Observed and estimated growth curve of females and males by the models and sex of animals




Figure 3: Growth curve with 3rd degree polynomial model by sex

according to age in ABC. In Figure 2, the growth curve of the observed weights by gender and estimated according to the models is presented. As can be seen in Figure 2, the model most compatible with the values observed in ABC from birth to 24 months is the 3rd degree polynomial model.

In similar studies, the 3rd degree polynomial model was determined to be the most suitable model in the Holstein breed by Heinrichs and Hargrove (12); in the Ayrshire, Brown Swiss and Shorthorn breeds by Heinrichs and Hargrove (18); and in the Holstein and Brown Swiss breeds by Akbulut

(19). On the other hand, the most suitable models were the Gompertz and Von Bertalanffy model ($R^2=0.70$) in Madura breed by Hartati and Putra (6); Richards model ($R^2=0.999$) in Holstein by Tutkun (10); the Logistic model in the pre-weaning period; the Gompertz and Richards models in the post-weaning period in the Holstein breed by Koşkan and Özkaya (20); Richards model ($R^2=0.968, 0.960$) in Brown-Swiss and Holsteins by Bayram and Akbulut (21); the Richards model ($R^2=0.976$) in Anatolian Buffaloes by Şahin et al. (22). The most suitable models differ in studies conducted with different breeds and environmental conditions. In practice,

Table 4: Least square means and standard errors (SE) of the 3rd degree polynomial model parameters by different environmental factors

Factor	Group	n	$\beta_0 \pm SE$	$\beta_1 \pm SE$	$\beta_2 \pm SE$	$\beta_3 \pm SE$
Sex	Female	48	17.61±0.608	0.01±0.182	0.011±0.0028	0.0009±0.0004
	Male	65	20.25±0.573	0.35±0.171	0.005±0.0027	0.0005±0.0003
Dam Age	2-3	31	21.18±1.773	-0.52±0.530	0.006±0.0083	-0.0006±0.0011
	4-7	46	17.45±0.797	0.80±0.238	0.002±0.0037	0.0006±0.0005
	8-10	17	18.12±1.037	0.17±0.310	0.006±0.0048	0.0007±0.0006
	11+	19	18.99±0.979	0.26±0.293	0.019±0.0046	0.0020±0.0006
Parity	1	29	14.40±1.704	0.92±0.510	0.010±0.0080	0.0023±0.0010
	2	24	18.00±0.985	-0.45±0.295	0.016±0.0046	0.0012±0.0006
	3	19	19.95±0.995	-0.34±0.297	0.018±0.0046	0.0008±0.0006
	4	15	19.48±1.084	-0.39±0.324	0.012±0.0050	-0.0001±0.0007
	5	12	20.90±1.189	0.13±0.355	0.003±0.0055	-0.0002±0.0007
	6	8	20.33±1.511	0.63±0.452	-0.002±0.0071	0.0001±0.0009
	7	6	19.47±1.683	0.73±0.503	-0.002±0.0079	0.0005±0.0010
Year	2015	18	17.81±1.026	0.49±0.307	0.003±0.0047	0.0005±0.0006
	2016	23	19.03±0.876	0.15±0.262	0.007±0.0041	0.0001±0.0005
	2017	17	18.39±0.967	0.17±0.289	0.003±0.0045	-0.0001±0.0006
	2018	24	19.70±0.784	0.21±0.234	0.008±0.0036	0.0011±0.0005
	2019	10	19.27±1.147	-0.10±0.343	0.012±0.0054	0.0016±0.0007
	2020	21	19.39±1.041	0.14±0.311	0.013±0.0047	0.0009±0.0006
Season	Winter	17	17.49±0.960 ^b	-0.15±0.287	0.179±0.0044 ^a	0.0019±0.001
	Spring	55	19.78±0.696 ^{ab}	0.37±0.208	0.002±0.0032 ^b	0.0002±0.000
	Summer	26	20.58±0.777 ^a	0.41±0.232	0.004±0.0036 ^b	0.0001±0.000
	Autumn	15	17.88±1.038 ^b	0.08±0.310	0.008±0.0048 ^b	0.0006±0.001

^{a,b} The means with the different superscripts within the factor in the same column are different (P<0.05).

$\beta_0, \beta_1, \beta_2, \beta_3$: regression coefficients of 3rd degree polynomial model

determining the weight-age relationship of cattle requires a lot of expense and time (21). In order to make reliable estimations in different regions and different breeds and to use the obtained parameters for selection purposes, first the identification of the appropriate model is necessary.

In the rest of this paper, the results of the 3rd degree polynomial model were presented since this model was

determined to be the best fitted to real measurements obtained from animals. Table 4 reflects the least square means and their corresponding standard errors of $\beta_0, \beta_1, \beta_2, \beta_3$ parameters by environmental factors. As a result of the analysis, β_0 values vary between 17.45-21.18 (except for parity 1) in all environmental factors. The differences between these values were found to be statistically significant only in the seasonal group (P<0.05). In addition, β_2 values




Figure 4: Growth curve with 3rd degree polynomial model by age of dam




Figure 5: Growth curve with 3rd degree polynomial model by parity

were found to be statistically significant between seasons ($P < 0.05$).

In Figures 3, 4, 5, 6, and 7, the growth curves of animals according to sex, dam age, parity, birth year, and birth season are presented using the 3rd degree polynomial model. As Figure 3 is examined, it has been determined that males have a higher weight than females in both observed and predicted values in all periods. Sahin et al. (22) found that adult live weight was higher in males in all models (Logistic, Gompertz, Richards, Brody) examined in Anatolian buffaloes. Hartati and Putra (6) reported that the animals had similar growth characteristics in all models (Logistik, Gompertz, Von Bertalanffy) which were examined in both sexes in Madura cattle. Growth in both males and females in the study continued until the age of 24 months, which

can be clearly seen in the linearly plotted growth curve in Figure 3. Akbulut (19), using the 3rd degree polynomial model, determined that the growth in Holstein and Brown Swiss breeds continued linearly up to 18 months.

When Figures 4 and 5 were examined, the differences according to dam age and parity became more pronounced after 18 months of age. According to the chosen model, it was estimated that calves from the 11+ dam age group had higher live weights in periods BW, 3M, and 24M. It was also estimated that the 8-10 dam age group had higher live weights in periods 6M and 12M, while the 2-3 dam age group had higher live weights in the 18M period. When live weights measured in different periods were examined according to parity, it was determined that the animals born from the 5th parity cows had a higher live weight in the birth




Figure 6: Growth curve with 3rd degree polynomial model by birth year




Figure 7: Growth curve with 3rd degree polynomial model by birth season

period. In addition, while the animals born from the 7th parity cows had a higher live weight in the 3M, 6M, and 12M periods, the animals born from the 2nd parity cows had a higher live weight in the 18M and 24M periods.

When Figures 6 and 7 are examined, the differences according to the year of birth and season of birth begin to appear mostly in 6M. While studying the differences in live weights by years, it was observed that the calves born in 2020 had higher live weights at birth, 6M, and 12M periods. Additionally, those born in 2018 had higher live weights in the 3M period, and those born in 2019 had higher live weights in the 18th and 24M periods. When the differences in live weights according to the seasons were examined, it was found that the animals born in the summer season had higher weights at birth and 3M periods. It was also

observed that while the animals born in the spring season had higher live weights at the 6M and 12M periods, the animals born in the autumn season had higher live weights at the 18M and 24M periods.

According to this model, the estimated values showed a deviation of around 1-2 kg in female and male animals at all periods compared to the observed values, while in males they showed a deviation of 3-5 kg only in the 12M and 18M periods (Figure 3). In other graphs (Figure 4-7), the differences between the generally estimated values and the observed values are between 1-3 kg, and the differences were found to be around 4-5 kg only in 18M periods. This indicates that the 3rd degree polynomial model is the most appropriate model for the growth values of ABC.

Monitoring the growth and development of animals during some periods in the growth process will be of great benefit to the farms in terms of herd management, care, and feeding regulation (22). In order to obtain reliable estimates of the growth curve parameters, it may be necessary to collect growth data until the point when the growth curve starts to flatten or the growth rate slows down (11). Changes in body weight in animals reflect the influence of environmental factors and management systems, particularly nutrition (23). In addition, by monitoring the growth of the animals, early intervention can be made for animals that have a problem in their development.

Conclusions

According to the results of the study, the 3rd degree polynomial model was determined to be the most suitable model in ABC according to the R², RSD, and corel values. By using the 3rd degree polynomial model on the farm, the general growth and development of the animals can be followed, and conditions such as sexual maturity age, breeding age, and appropriate slaughter age can be easily predicted. Examining the growth curve is important for breeders to decide on the optimum body weight of the animals, the appropriate age, and the ideal weight. Growth curve parameters can be successfully applied to animals and may benefit the development and design of selection strategies.

Acknowledgements

The population of the study consisted of Anatolian Black Cattle breeds practiced in connection with the "Conservation of Domestic Genetic Resources and Sustainable Use". Therefore, the authors kindly acknowledge the contribution of the General Directorate of Agricultural Research and Policies (Ministry of Agriculture and Forestry) of the Republic of Turkey, which has given the necessary permission to provide the animal material used in the study.

The author of the study kindly declares no competing interest.

References

1. Tuik. Data portal for statistics [online]. Ankara: Turkish statistical institute, 2023. <https://data.tuik.gov.tr/Kategori/GetKategori?p=Tarim-111> (17. 8. 2023)
2. Sakar ÇM, Zülkadir U. Determination of the relationship between Anatolian black cattle growth properties and myostatin, GHR and Pit-1 gene. *Anim Biotechnol* 2022; 33: 536–45.
3. Ünal İ, Tuncer HI, Sakar ÇM, Ünay E. The effect of maternal age on some body measurements in Anatolian Black calves. *Black Sea J Agric* 2019; 2: 47–50.
4. Sakar ÇM, Zülkadir U. Determination of some growth and development characteristics between birth and twelve months age in Yerli Kara cattle. *J Agric Sci* 2022; 28: 33–9.
5. Kılıçel R. Examination of different breeding types in Anatolian Black and Brown Swiss cattles reared in Şile region. Konya; Selçuk University, 2014. Doctoral Thesis
6. Hartati H, Pintaka BPW. Predicting the growth curve of body weight in Madura cattle. *Kafkas Univ Vet Fac Derg* 2021; 27: 431–7.
7. Aytekin İ, Karabacak A, Zülkadir U, Keskin İ, Boztepe S. Usage of some models for describing the growth curves of Akkaraman and Anatolian Merino lambs raised in open and closed sheepfolds at the fattening period. *Selc J Agric Food Sci* 2009; 23: 30–5.
8. Bilgin ÖC, Esenbuğa N. Parameter estimation in nonlinear growth models. *J Anim Prod* 2003; 44: 81–90.
9. Efe E. Growth curves. Adana: Institute of Science Department of Animal Science, 1990. Doctoral Thesis
10. Tutkun M. Growth curve prediction of Holstein-Fresian bulls using different non-linear model functions. *Appl Ecol Environ Res* 2019; 17: 4409–16.
11. Koncagül S, Çadırcı S. Comparison of three non-linear models when data truncated at different lengths of growth period in Japanese quails. *Arch Geflügelk* 2009; 73: 7–12.
12. Heinrichs AJ, Hargrove GL. Standards of weight and height for Holstein heifers. *J Dairy Sci* 1987; 70: 653–60.
13. Nelder JA. The fitting of a generalization of the Logistic curve. *Biometrics* 1961; 17: 89–110.
14. Brody S. Bioenergetics and growth, with special reference to the efficiency complex in domestic animals. New York: Reinhold Publishing Corporation, 1945.
15. Von Bertalanffy L. Quantitative laws in metabolism and growth. *Quarterly Rev Biol* 1957; 32: 217–31.
16. Gompertz B. On the nature of the function expressive of the law of human mortality, and on a new mode of determining the value of life contingencies. *Philosoph Trans* 1825; 115: 513–85.
17. SAS. SAS Enterprise Guide 8.3 [online]. Cary: SAS Institute Inc, 2023 <https://support.sas.com/en/software/enterprise-guide-support.html#documentation> (17. 8. 2023)
18. Heinrichs AJ, Hargrove GL. Standards of weight and height for Ayrshire, Brown Swiss and milking Shorthorn heifers, *J Dairy Sci* 1994; 77: 1676–81.
19. Akbulut Ö. Growth analysis of Brown Swiss and Holstein heifers under harsh climatic conditions. *Turkish J Vet Anim Sci* 1999; 23: 131–7.
20. Koskan O, Ozkaya S. Determination of growth curves of female Holstein calves using five non-linear models. *Pak J Agric Sci* 2014; 51: 225–8.
21. Bayram B, Akbulut Ö. Analysis of growth curve in Brown Swiss and Holstein Friesian cattle using linear and non-linear models. *J Anim Prod* 2009; 50: 33–40.
22. Şahin A, Ulutaş Z, Karadavut U, Yıldırım A, Arslan S. Comparison of growth curve using some nonlinear models in Anatolian Buffalo calves. *Kafkas Univ Vet Fac Derg* 2014; 20: 357–62.
23. Entwistle K, Talib C, Siregar A, Budiarti S, Turner W, Lindsay D. Bali cattle performance: Current population dynamics and performance and some strategies for improvement (A preliminary report). *Indonesian J Anim Vet Sci* 2012; 17: 31–9.

Določitev najboljšega modela krivulje rasti za anatolsko črno govedo

Ç. M. Sakar, S. Koncagül, I. Ünal

Izvleček: Namen te študije je bil določiti model, ki najbolje opisuje potek rasti od rojstva do 24 mesecev starosti anatolskega črnega goveda (ABC), vzrejenega za namene ohranjanja. Zbranih je bilo 493 podatkov o telesni teži 113 živali ob rojstvu ter pri 3, 6, 12, 18 in 24 mesecih starosti. Za opis krivulje rasti živali je bilo uporabljenih šest različnih nelinearnih modelov, in sicer polinom 2. stopnje, polinom 3. stopnje, logistični model ter modeli Brody, Von Bertalanffy in Gompertz. V študiji so bile vrednosti R^2 modelov naslednje: 0,997, 0,999, 0,953, 0,979, 0,924 in 0,862; vrednosti correl (korelacija med opazovanimi in ocenjenimi krivuljami) so bile 0,994, 0,998, 0,989, 0,993, 0,961 in 0,703; ostanki standardnih odklonov (RSD) so bili 3,216, 1,388, 11,533, 3,561, 14,736 in 27,141. Glede na te vrednosti je bilo ugotovljeno, da je polinomski model 3. stopnje najbolje opisal krivuljo rasti ABC. Na podlagi analiz je bilo ugotovljeno, da so vrednosti, ki jih je napovedal ta model, za 1-3 kg odstopale od ugotovljenih vrednosti v vseh obdobjih in pri vseh preučevanih okoljskih dejavnikih (spol, starost matere, pariteta, leto rojstva in sezona rojstva). Ugotovljeno je bilo, da so se te razlike povečale na 4-5 kg le v 18-mesečnem obdobju. Rezultati so tudi pokazali, da se je ABC še naprej povečevala po 24 mesecih starosti. Posledično je mogoče lastnosti, kot so starost ob spolni zrelosti, plemenska starost in klavna starost, enostavno napovedati z določitvijo modela, ki najbolje opisuje rast in razvoj v čredah.

Ključne besede: anatolsko črno govedo; živa teža; krivulja rasti; nelinearni modeli