

# On Hermitian varieties in $\text{PG}(6, q^2)$

Angela Aguglia \* 

*Dipartimento di Meccanica, Matematica e Management, Politecnico di Bari,  
Via Orabona 4, I-70125 Bari, Italy*

Luca Giuzzi † 

*DICATAM, University of Brescia, Via Branze 53, I-25123 Brescia, Italy*

Masaaki Homma 

*Department of Mathematics and Physics, Kanagawa University,  
Hiratsuka 259-1293, Japan*

Received 8 June 2020, accepted 15 February 2021, published online 10 August 2021

---

## Abstract

In this paper we characterize the non-singular Hermitian variety  $\mathcal{H}(6, q^2)$  of  $\text{PG}(6, q^2)$ ,  $q \neq 2$  among the irreducible hypersurfaces of degree  $q + 1$  in  $\text{PG}(6, q^2)$  not containing solids by the number of its points and the existence of a solid  $S$  meeting it in  $q^4 + q^2 + 1$  points.

*Keywords:* Unital, Hermitian variety, algebraic hypersurface.

*Math. Subj. Class. (2020):* 51E21, 51E15, 51E20

---

## 1 Introduction

The set of all absolute points of a non-degenerate unitary polarity in  $\text{PG}(r, q^2)$  determines the Hermitian variety  $\mathcal{H}(r, q^2)$ . This is a non-singular algebraic hypersurface of degree  $q + 1$  in  $\text{PG}(r, q^2)$  with a number of remarkable properties, both from the geometrical and the combinatorial point of view; see [6, 16]. In particular,  $\mathcal{H}(r, q^2)$  is a 2-character set with respect to the hyperplanes of  $\text{PG}(r, q^2)$  and 3-character blocking set with respect to the

---

\*Corresponding author. The author was supported by the Italian National Group for Algebraic and Geometric Structures and their Applications (GNSAGA - INdAM).

†The author was supported by the Italian National Group for Algebraic and Geometric Structures and their Applications (GNSAGA - INdAM).

*E-mail addresses:* [angela.aguglia@poliba.it](mailto:angela.aguglia@poliba.it) (Angela Aguglia), [luca.giuzzi@unibs.it](mailto:luca.giuzzi@unibs.it) (Luca Giuzzi), [homma@kanagawa-u.ac.jp](mailto:homma@kanagawa-u.ac.jp) (Masaaki Homma)

lines of  $\text{PG}(r, q^2)$  for  $r > 2$ . An interesting and widely investigated problem is to provide combinatorial descriptions of  $\mathcal{H}(r, q^2)$ .

First, we observe that a condition on the number of points and the intersection numbers with hyperplanes is not in general sufficient to characterize Hermitian varieties; see [1, 2]. On the other hand, it is enough to consider in addition the intersection numbers with codimension 2 subspaces in order to get a complete description; see [7].

In general, a hypersurface  $\mathcal{H}$  of  $\text{PG}(r, q)$  is viewed as a hypersurface over the algebraic closure of  $\text{GF}(q)$  and a point of  $\text{PG}(r, q^i)$  in  $\mathcal{H}$  is called a  $\text{GF}(q^i)$ -point. A  $\text{GF}(q)$ -point of  $\mathcal{H}$  is also said to be a rational point of  $\mathcal{H}$ . Throughout this paper, the number of  $\text{GF}(q^i)$ -points of  $\mathcal{H}$  will be denoted by  $N_{q^i}(\mathcal{H})$ . For simplicity, we shall also use the convention  $|\mathcal{H}| = N_q(\mathcal{H})$ .

In the present paper, we shall investigate a combinatorial characterization of the Hermitian hypersurface  $\mathcal{H}(6, q^2)$  in  $\text{PG}(6, q^2)$  among all hypersurfaces of the same degree having also the same number of  $\text{GF}(q^2)$ -rational points.

More in detail, in [12, 13] it has been proved that if  $\mathcal{X}$  is a hypersurface of degree  $q + 1$  in  $\text{PG}(r, q^2)$ ,  $r \geq 3$  odd, with  $|\mathcal{X}| = |\mathcal{H}(r, q^2)| = (q^{r+1} + (-1)^r)(q^r - (-1)^r)/(q^2 - 1)$   $\text{GF}(q^2)$ -rational points, not containing linear subspaces of dimension greater than  $\frac{r-1}{2}$ , then  $\mathcal{X}$  is a non-singular Hermitian variety of  $\text{PG}(r, q^2)$ . This result generalizes the characterization of [8] for the Hermitian curve of  $\text{PG}(2, q^2)$ ,  $q \neq 2$ .

The case where  $r > 4$  is even is, in general, currently open. A starting point for a characterization in arbitrary even dimension can be found in [3] where the case of a hypersurface  $\mathcal{X}$  of degree  $q + 1$  in  $\text{PG}(4, q^2)$ ,  $q > 3$  is considered. There, it is shown that when  $\mathcal{X}$  has the same number of rational points as  $\mathcal{H}(4, q^2)$ , does not contain any subspaces of dimension greater than 1 and meets at least one plane  $\pi$  in  $q^2 + 1$   $\text{GF}(q^2)$ -rational points, then  $\mathcal{X}$  is a Hermitian variety.

In this article we deal with hypersurfaces of degree  $q + 1$  in  $\text{PG}(6, q^2)$  and we prove that a characterization similar to that of [3] holds also in dimension 6. We conjecture that this can be extended to arbitrary even dimension.

**Theorem 1.1.** *Let  $S$  be a hypersurface of  $\text{PG}(6, q^2)$ ,  $q > 2$ , defined over  $\text{GF}(q^2)$ , not containing solids. If the degree of  $S$  is  $q + 1$  and the number of its rational points is  $q^{11} + q^9 + q^7 + q^4 + q^2 + 1$ , then every solid of  $\text{PG}(6, q^2)$  meets  $S$  in at least  $q^4 + q^2 + 1$  rational points. If there is at least a solid  $\Sigma_3$  such that  $|\Sigma_3 \cap S| = q^4 + q^2 + 1$ , then  $S$  is a non-singular Hermitian variety of  $\text{PG}(6, q^2)$ .*

Furthermore, we also extend the result of [3] to the case  $q = 3$ .

## 2 Preliminaries and notation

In this section we collect some useful information and results that will be crucial to our proof.

A Hermitian variety in  $\text{PG}(r, q^2)$  is the algebraic variety of  $\text{PG}(r, q^2)$  whose points  $\langle v \rangle$  satisfy the equation  $\eta(v, v) = 0$  where  $\eta$  is a sesquilinear form  $\text{GF}(q^2)^{r+1} \times \text{GF}(q^2)^{r+1} \rightarrow \text{GF}(q^2)$ . The radical of the form  $\eta$  is the vector subspace of  $\text{GF}(q^2)^{r+1}$  given by

$$\text{Rad}(\eta) := \{w \in \text{GF}(q^2)^{r+1} : \forall v \in \text{GF}(q^2)^{r+1}, \eta(v, w) = 0\}.$$

The form  $\eta$  is non-degenerate if  $\text{Rad}(\eta) = \{0\}$ . If the form  $\eta$  is non-degenerate, then the corresponding Hermitian variety is denoted by  $\mathcal{H}(r, q^2)$  and it is a non-singular algebraic

variety, of degree  $q + 1$  containing

$$(q^{r+1} + (-1)^r)(q^r - (-1)^r)/(q^2 - 1)$$

$\text{GF}(q^2)$ -rational points. When  $\eta$  is degenerate we shall call *vertex*  $R_t$  of the degenerate Hermitian variety associated to  $\eta$  the projective subspace  $R_t := \text{PG}(\text{Rad}(\eta)) := \{\langle w \rangle : w \in \text{Rad}(\eta)\}$  of  $\text{PG}(r, q^2)$ . A degenerate Hermitian variety can always be described as a cone of vertex  $R_t$  and basis a non-degenerate Hermitian variety  $\mathcal{H}(r - t, q^2)$  disjoint from  $R_t$  where  $t = \dim(\text{Rad}(\eta))$  is the vector dimension of the radical of  $\eta$ . In this case we shall write the corresponding variety as  $R_t\mathcal{H}(r - t, q^2)$ . Indeed,

$$R_t\mathcal{H}(r - t, q^2) := \{X \in \langle P, Q \rangle : P \in R_t, Q \in \mathcal{H}(r - t, q^2)\}.$$

Any line of  $\text{PG}(r, q^2)$  meets a Hermitian variety (either degenerate or not) in either  $1, q + 1$  or  $q^2 + 1$  points (the latter value only for  $r > 2$ ). The maximal dimension of projective subspaces contained in the non-degenerate Hermitian variety  $\mathcal{H}(r, q^2)$  is  $(r - 2)/2$ , if  $r$  is even, or  $(r - 1)/2$ , if  $r$  is odd. These subspaces of maximal dimension are called *generators* of  $\mathcal{H}(r, q^2)$  and the generators of  $\mathcal{H}(r, q^2)$  through a point  $P$  of  $\mathcal{H}(r, q^2)$  span a hyperplane  $P^\perp$  of  $\text{PG}(r, q^2)$ , the *tangent hyperplane* at  $P$ .

It is well known that this hyperplane meets  $\mathcal{H}(r, q^2)$  in a degenerate Hermitian variety  $P\mathcal{H}(r - 2, q^2)$ , that is in a Hermitian cone having as vertex the point  $P$  and as base a non-singular Hermitian variety of  $\Theta \cong \text{PG}(r - 2, q^2)$  contained in  $P^\perp$  with  $P \notin \Theta$ .

Every hyperplane of  $\text{PG}(r, q^2)$  that is not tangent meets  $\mathcal{H}(r, q^2)$  in a non-singular Hermitian variety  $\mathcal{H}(r - 1, q^2)$ , and is called a *secant hyperplane* of  $\mathcal{H}(r, q^2)$ . In particular, a tangent hyperplane contains

$$1 + q^2(q^{r-1} + (-1)^r)(q^{r-2} - (-1)^r)/(q^2 - 1)$$

$\text{GF}(q^2)$ -rational points of  $\mathcal{H}(r, q^2)$ , whereas a secant hyperplane contains

$$(q^r + (-1)^{r-1})(q^{r-1} - (-1)^{r-1})/(q^2 - 1)$$

$\text{GF}(q^2)$ -rational points of  $\mathcal{H}(r, q^2)$ .

We now recall several results which shall be used in the course of this paper.

**Lemma 2.1** ([15]). *Let  $d$  be an integer with  $1 \leq d \leq q + 1$  and let  $\mathcal{C}$  be a curve of degree  $d$  in  $\text{PG}(2, q)$  defined over  $\text{GF}(q)$ , which may have  $\text{GF}(q)$ -linear components. Then the number of its rational points is at most  $dq + 1$  and  $N_q(\mathcal{C}) = dq + 1$  if and only if  $\mathcal{C}$  is a pencil of  $d$  lines of  $\text{PG}(2, q)$ .*

**Lemma 2.2** ([10]). *Let  $d$  be an integer with  $2 \leq d \leq q + 2$ , and  $\mathcal{C}$  a curve of degree  $d$  in  $\text{PG}(2, q)$  defined over  $\text{GF}(q)$  without any  $\text{GF}(q)$ -linear components. Then  $N_q(\mathcal{C}) \leq (d - 1)q + 1$ , except for a class of plane curves of degree 4 over  $\text{GF}(4)$  having 14 rational points.*

**Lemma 2.3** ([11]). *Let  $\mathcal{S}$  be a surface of degree  $d$  in  $\text{PG}(3, q)$  over  $\text{GF}(q)$ . Then*

$$N_q(\mathcal{S}) \leq dq^2 + q + 1$$

**Lemma 2.4** ([8]). *Suppose  $q \neq 2$ . Let  $\mathcal{C}$  be a plane curve over  $\text{GF}(q^2)$  of degree  $q + 1$  without  $\text{GF}(q^2)$ -linear components. If  $\mathcal{C}$  has  $q^3 + 1$  rational points, then  $\mathcal{C}$  is a Hermitian curve.*

**Lemma 2.5** ([7]). *A subset of points of  $\text{PG}(r, q^2)$  having the same intersection numbers with respect to hyperplanes and spaces of codimension 2 as non-singular Hermitian varieties is a non-singular Hermitian variety of  $\text{PG}(r, q^2)$ .*

From [9, Theorem 23.5.1, Theorem 23.5.3] we have the following.

**Lemma 2.6.** *If  $\mathcal{W}$  is a set of  $q^7 + q^4 + q^2 + 1$  points of  $\text{PG}(4, q^2)$ ,  $q > 2$  such that every line of  $\text{PG}(4, q^2)$  meets  $\mathcal{W}$  in  $1, q + 1$  or  $q^2 + 1$  points, then  $\mathcal{W}$  is a Hermitian cone with vertex a line and base a unital.*

Finally, we recall that a *blocking set with respect to lines* of  $\text{PG}(r, q)$  is a point set which blocks all the lines, i.e., intersects each line of  $\text{PG}(r, q)$  in at least one point.

### 3 Proof of Theorem 1.1

We first provide an estimate on the number of points of a curve of degree  $q + 1$  in  $\text{PG}(2, q^2)$ , where  $q$  is any prime power.

**Lemma 3.1.** *Let  $\mathcal{C}$  be a plane curve over  $\text{GF}(q^2)$ , without  $\overline{\text{GF}(q^2)}$ -lines as components and of degree  $q + 1$ . If the number of  $\text{GF}(q^2)$ -rational points of  $\mathcal{C}$  is  $N < q^3 + 1$ , then*

$$N \leq \begin{cases} q^3 - (q^2 - 2) & \text{if } q > 3 \\ 24 & \text{if } q = 3 \\ 8 & \text{if } q = 2. \end{cases} \tag{3.1}$$

*Proof.* We distinguish the following three cases:

- (a)  $\mathcal{C}$  has two or more  $\text{GF}(q^2)$ -components;
- (b)  $\mathcal{C}$  is irreducible over  $\text{GF}(q^2)$ , but not absolutely irreducible;
- (c)  $\mathcal{C}$  is absolutely irreducible.

Suppose first  $q \neq 2$ .

**Case (a)** Suppose  $\mathcal{C} = \mathcal{C}_1 \cup \mathcal{C}_2$ . Let  $d_i$  be the degree of  $\mathcal{C}_i$ , for each  $i = 1, 2$ . Hence  $d_1 + d_2 = q + 1$ . By Lemma 2.2,

$$N \leq N_{q^2}(\mathcal{C}_1) + N_{q^2}(\mathcal{C}_2) \leq [(q + 1) - 2]q^2 + 2 = q^3 - (q^2 - 2)$$

**Case (b)** Let  $\mathcal{C}'$  be an irreducible component of  $\mathcal{C}$  over the algebraic closure of  $\text{GF}(q^2)$ . Let  $\text{GF}(q^{2t})$  be the minimum defining field of  $\mathcal{C}'$  and  $\sigma$  be the Frobenius morphism of  $\text{GF}(q^{2t})$  over  $\text{GF}(q^2)$ . Then

$$\mathcal{C} = \mathcal{C}' \cup \mathcal{C}'^\sigma \cup \mathcal{C}'^{\sigma^2} \cup \dots \cup \mathcal{C}'^{\sigma^{t-1}},$$

and the degree of  $\mathcal{C}'$ , say  $e$ , satisfies  $q + 1 = te$  with  $e > 1$ . Hence any  $\text{GF}(q^2)$ -rational point of  $\mathcal{C}$  is contained in  $\bigcap_{i=0}^{t-1} \mathcal{C}'^{\sigma^i}$ . In particular,  $N \leq e^2 \leq (\frac{q+1}{2})^2$  by Bezout's Theorem and  $(\frac{q+1}{2})^2 < q^3 - (q^2 - 2)$ .

**Case (c)** Let  $\mathcal{C}$  be an absolutely irreducible curve over  $\text{GF}(q^2)$  of degree  $q + 1$ . Either  $\mathcal{C}$  has a singular point or not.

In general, an absolutely irreducible plane curve  $\mathcal{M}$  over  $\text{GF}(q^2)$  is  $q^2$ -Frobenius non-classical if for a general point  $P(x_0, x_1, x_2)$  of  $\mathcal{M}$  the point  $P^{q^2} = P^{q^2}(x_0^{q^2}, x_1^{q^2}, x_2^{q^2})$  is

on the tangent line to  $\mathcal{M}$  at the point  $P$ . Otherwise, the curve  $\mathcal{M}$  is said to be Frobenius classical. A lower bound of the number of  $\text{GF}(q^2)$ -points for  $q^2$ -Frobenius non-classical curves is given by [4, Corollary 1.4]: for a  $q^2$ -Frobenius non-classical curve  $\mathcal{C}'$  of degree  $d$ , we have  $N_{q^2}(\mathcal{C}') \geq d(q^2 - d + 2)$ . In particular, if  $d = q + 1$ , the lower bound is just  $q^3 + 1$ .

Going back to our original curve  $\mathcal{C}$ , we know that  $\mathcal{C}$  is Frobenius classical because  $N < q^3 + 1$ . Let  $F(x, y, z) = 0$  be an equation of  $\mathcal{C}$  over  $\text{GF}(q^2)$ . We consider the curve  $\mathcal{D}$  defined by  $\frac{\partial F}{\partial x}x^{q^2} + \frac{\partial F}{\partial y}y^{q^2} + \frac{\partial F}{\partial z}z^{q^2} = 0$ . Then  $\mathcal{C}$  is not a component of  $\mathcal{D}$  because  $\mathcal{C}$  is Frobenius classical. Furthermore, any  $\text{GF}(q^2)$ -point  $P$  lies on  $\mathcal{C} \cap \mathcal{D}$  and the intersection multiplicity of  $\mathcal{C}$  and  $\mathcal{D}$  at  $P$  is at least 2 by Euler's theorem for homogeneous polynomials. Hence by Bézout's theorem,  $2N \leq (q + 1)(q^2 + q)$ . Hence

$$N \leq \frac{1}{2}q(q + 1)^2.$$

This argument is due to Stöhr and Voloch [18, Theorem 1.1]. This Stöhr and Voloch's bound is lower than the estimate for  $N$  in case (a) for  $q > 4$  and it is the same for  $q = 4$ . When  $q = 3$  the bound in case (a) is smaller than the Stöhr and Voloch's bound. Finally, we consider the case  $q = 2$ . Under this assumption,  $\mathcal{C}$  is a cubic curve and neither case (a) nor case (b) might occur. For a degree 3 curve over  $\text{GF}(q^2)$  the Stöhr and Voloch's bound is loose, thus we need to change our argument. If  $\mathcal{C}$  has a singular point, then  $\mathcal{C}$  is a rational curve with a unique singular point. Since the degree of  $\mathcal{C}$  is 3, singular points are either cusps or ordinary double points. Hence  $N \in \{4, 5, 6\}$ . If  $\mathcal{C}$  is nonsingular, then it is an elliptic curve and, by the Hasse-Weil bound, see [19],  $N \in I$  where  $I = \{1, 2, \dots, 9\}$  and for each number  $N$  belonging to  $I$  there is an elliptic curve over  $\text{GF}(4)$  with  $N$  points, from [14, Theorem 4.2]. This completes the proof.  $\square$

Henceforth, we shall always suppose  $q > 2$  and we denote by  $\mathcal{S}$  an algebraic hypersurface of  $\text{PG}(6, q^2)$  satisfying the following hypotheses of Theorem 1.1:

- (S1)  $\mathcal{S}$  is an algebraic hypersurface of degree  $q + 1$  defined over  $\text{GF}(q^2)$ ;
- (S2)  $|\mathcal{S}| = q^{11} + q^9 + q^7 + q^4 + q^2 + 1$ ;
- (S3)  $\mathcal{S}$  does not contain projective 3-spaces (solids);
- (S4) there exists a solid  $\Sigma_3$  such that  $|\mathcal{S} \cap \Sigma_3| = q^4 + q^2 + 1$ .

We first consider the behavior of  $\mathcal{S}$  with respect to the lines.

**Lemma 3.2.** *An algebraic hypersurface  $\mathcal{T}$  of degree  $q + 1$  in  $\text{PG}(r, q^2)$ ,  $q \neq 2$ , with  $|\mathcal{T}| = |\mathcal{H}(r, q^2)|$  is a blocking set with respect to lines of  $\text{PG}(r, q^2)$*

*Proof.* Suppose on the contrary that there is a line  $\ell$  of  $\text{PG}(r, q^2)$  which is disjoint from  $\mathcal{T}$ . Let  $\alpha$  be a plane containing  $\ell$ . The algebraic plane curve  $\mathcal{C} = \alpha \cap \mathcal{T}$  of degree  $q + 1$  cannot have  $\text{GF}(q^2)$ -linear components and hence it has at most  $q^3 + 1$  points because of Lemma 2.2. If  $\mathcal{C}$  had  $q^3 + 1$  rational points, then from Lemma 2.4,  $\mathcal{C}$  would be a Hermitian curve with an external line, a contradiction since Hermitian curves are blocking sets. Thus  $N_{q^2}(\mathcal{C}) \leq q^3$ . Since  $q > 2$ , by Lemma 3.1,  $N_{q^2}(\mathcal{C}) < q^3 - 1$  and hence every plane through  $r$  meets  $\mathcal{T}$  in at most  $q^3 - 1$  rational points. Consequently, by considering all planes through  $r$ , we can bound the number of rational points of  $\mathcal{T}$  by  $N_{q^2}(\mathcal{T}) \leq (q^3 - 1) \frac{q^{2r-4} - 1}{q^2 - 1} =$

$q^{2r-3} + \dots < |\mathcal{H}(r, q^2)|$ , which is a contradiction. Therefore there are no external lines to  $\mathcal{T}$  and so  $\mathcal{T}$  is a blocking set w.r.t. lines of  $\text{PG}(r, q^2)$ .  $\square$

**Remark 3.3.** The proof of [3, Lemma 3.1] would work perfectly well here under the assumption  $q > 3$ . The alternative argument of Lemma 3.2 is simpler and also holds for  $q = 3$ .

By the previous Lemma and assumptions (S1) and (S2),  $\mathcal{S}$  is a blocking set for the lines of  $\text{PG}(6, q^2)$ . In particular, the intersection of  $\mathcal{S}$  with any 3-dimensional subspace  $\Sigma$  of  $\text{PG}(6, q^2)$  is also a blocking set with respect to lines of  $\Sigma$  and hence it contains at least  $q^4 + q^2 + 1$   $\text{GF}(q^2)$ -rational points; see [5].

**Lemma 3.4.** *Let  $\Sigma_3$  be a solid of  $\text{PG}(6, q^2)$  satisfying condition (S4), that is  $\Sigma_3$  meets  $\mathcal{S}$  in exactly  $q^4 + q^2 + 1$  points. Then,  $\Pi := \mathcal{S} \cap \Sigma_3$  is a plane.*

*Proof.*  $\mathcal{S} \cap \Sigma_3$  must be a blocking set for the lines of  $\text{PG}(3, q^2)$ ; also it has size  $q^4 + q^2 + 1$ . It follows from [5] that  $\Pi := \mathcal{S} \cap \Sigma_3$  is a plane.  $\square$

**Lemma 3.5.** *Let  $\Sigma_3$  be a solid of satisfying condition (S4). Then, any 4-dimensional projective space  $\Sigma_4$  through  $\Sigma_3$  meets  $\mathcal{S}$  in a Hermitian cone with vertex a line and basis a Hermitian curve.*

*Proof.* Consider all of the  $q^6 + q^4 + q^2 + 1$  subspaces  $\bar{\Sigma}_3$  of dimension 3 in  $\text{PG}(6, q^2)$  containing  $\Pi = \mathcal{S} \cap \Sigma_3$ .

From Lemma 2.3 and condition (S3) we have  $|\bar{\Sigma}_3 \cap \mathcal{S}| \leq q^5 + q^4 + q^2 + 1$ . Hence,

$$|\mathcal{S}| = (q^7 + 1)(q^4 + q^2 + 1) \leq (q^6 + q^4 + q^2)q^5 + q^4 + q^2 + 1 = |\mathcal{S}|.$$

Consequently,  $|\bar{\Sigma}_3 \cap \mathcal{S}| = q^5 + q^4 + q^2 + 1$  for all  $\bar{\Sigma}_3 \neq \Sigma_3$  such that  $\Pi \subset \bar{\Sigma}_3$ .

Let  $C := \Sigma_4 \cap \mathcal{S}$ . Counting the number of rational points of  $C$  by considering the intersections with the  $q^2 + 1$  subspaces  $\Sigma'_3$  of dimension 3 in  $\Sigma_4$  containing the plane  $\Pi$  we get

$$|C| = q^2 \cdot q^5 + q^4 + q^2 + 1 = q^7 + q^4 + q^2 + 1.$$

In particular,  $C \cap \Sigma'_3$  is a maximal surface of degree  $q + 1$ ; so it must split in  $q + 1$  distinct planes through a line of  $\Pi$ ; see [17]. So  $C$  consists of  $q^3 + 1$  distinct planes belonging to distinct  $q^2$  pencils, all containing  $\Pi$ ; denote by  $\mathcal{L}$  the family of these planes. Also for each  $\Sigma'_3 \neq \Sigma_3$ , there is a line  $\ell'$  such that all the planes of  $\mathcal{L}$  in  $\Sigma'_3$  pass through  $\ell'$ . It is now straightforward to see that any line contained in  $C$  must necessarily belong to one of the planes of  $\mathcal{L}$  and no plane not in  $\mathcal{L}$  is contained in  $C$ .

In order to get the result it is now enough to show that a line of  $\Sigma_4$  meets  $C$  in either 1,  $q + 1$  or  $q^2 + 1$  points. To this purpose, let  $\ell$  be a line of  $\Sigma_4$  and suppose  $\ell \not\subset C$ . Then, by Bezout's theorem,

$$1 \leq |\ell \cap C| \leq q + 1.$$

Assume  $|\ell \cap C| > 1$ . Then we can distinguish two cases:

1.  $\ell \cap \Pi \neq \emptyset$ . If  $\ell$  and  $\Pi$  are incident, then we can consider the 3-dimensional subspace  $\Sigma'_3 := \langle \ell, \Pi \rangle$ . Then  $\ell$  must meet each plane of  $\mathcal{L}$  in  $\Sigma'_3$  in different points (otherwise  $\ell$  passes through the intersection of these planes and then  $|\ell \cap C| = 1$ ). As there are  $q + 1$  planes of  $\mathcal{L}$  in  $\Sigma'_3$ , we have  $|\ell \cap C| = q + 1$ .

2.  $\ell \cap \Pi = \emptyset$ . Consider the plane  $\Lambda$  generated by a point  $P \in \Pi$  and  $\ell$ . Clearly  $\Lambda \notin \mathcal{L}$ . The curve  $\Lambda \cap \mathcal{S}$  has degree  $q+1$  by construction, does not contain lines (for otherwise  $\Lambda \in \mathcal{L}$ ) and has  $q^3+1$   $\text{GF}(q^2)$ -rational points (by a counting argument). So from Lemma 2.4 it is a Hermitian curve. It follows that  $\ell$  is a  $q+1$  secant.

We can now apply Lemma 2.6 to see that  $C$  is a Hermitian cone with vertex a line.  $\square$

**Lemma 3.6.** *Let  $\Sigma_3$  be a space satisfying condition (S4) and take  $\Sigma_5$  to be a 5-dimensional projective space with  $\Sigma_3 \subseteq \Sigma_5$ . Then  $\mathcal{S} \cap \Sigma_5$  is a Hermitian cone with vertex a point and basis a Hermitian hypersurface  $\mathcal{H}(4, q^2)$ .*

*Proof.* Let

$$\Sigma_4 := \Sigma_4^1, \Sigma_4^2, \dots, \Sigma_4^{q^2+1}$$

be the 4-spaces through  $\Sigma_3$  contained in  $\Sigma_5$ . Put  $C_i := \Sigma_4^i \cap \mathcal{S}$ , for all  $i \in \{1, \dots, q^2+1\}$  and  $\Pi = \Sigma_3 \cap C_1$ . From Lemma 3.5  $C_i$  is a Hermitian cone with vertex a line, say  $\ell_i$ . Furthermore  $\Pi \subseteq \Sigma_3 \subseteq \Sigma_4^i$  where  $\Pi$  is a plane. Choose a plane  $\Pi' \subseteq \Sigma_4^1$  such that  $m := \Pi' \cap C_1$  is a line  $m$  incident with  $\Pi$  but not contained in it. Let  $P_1 := m \cap \Pi$ . It is straightforward to see that in  $\Sigma_4^i$  there are exactly 1 plane through  $m$  which is a  $(q^4+q^2+1)$ -secant,  $q^4$  planes which are  $(q^3+q^2+1)$ -secant and  $q^2$  planes which are  $(q^2+1)$ -secant. Also  $P_1$  belongs to the line  $\ell_1$ . There are now two cases to consider:

- (a) There is a plane  $\Pi'' \neq \Pi'$  not contained in  $\Sigma_4^i$  for all  $i = 1, \dots, q^2+1$  with  $m \subseteq \Pi'' \subseteq \mathcal{S} \cap \Sigma_5$ .

We first show that the vertices of the cones  $C_i$  are all concurrent. Consider  $m_i := \Pi'' \cap \Sigma_4^i$ . Then  $\{m_i : i = 1, \dots, q^2+1\}$  consists of  $q^2+1$  lines (including  $m$ ) all through  $P_1$ . Observe that for all  $i$ , the line  $m_i$  meets the vertex  $\ell_i$  of the cone  $C_i$  in  $P_i \in \Pi$ . This forces  $P_1 = P_2 = \dots = P_{q^2+1}$ . So  $P_1 \in \ell_1, \dots, \ell_{q^2+1}$ .

Now let  $\bar{\Sigma}_4$  be a 4-dimensional space in  $\Sigma_5$  with  $P_1 \notin \bar{\Sigma}_4$ ; in particular  $\Pi \not\subseteq \bar{\Sigma}_4$ . Put also  $\bar{\Sigma}_3 := \Sigma_4^1 \cap \bar{\Sigma}_4$ . Clearly,  $r := \bar{\Sigma}_3 \cap \Pi$  is a line and  $P_1 \notin r$ . So  $\bar{\Sigma}_3 \cap \mathcal{S}$  cannot be the union of  $q+1$  planes, since if this were to be the case, these planes would have to pass through the vertex  $\ell_1$ . It follows that  $\bar{\Sigma}_3 \cap \mathcal{S}$  must be a Hermitian cone with vertex a point and basis a Hermitian curve. Let  $\mathcal{W} := \bar{\Sigma}_4 \cap \mathcal{S}$ . The intersection  $\mathcal{W} \cap \Sigma_4^i$ , as  $i$  varies, is a Hermitian cone with basis a Hermitian curve, so, the points of  $\mathcal{W}$  are

$$|\mathcal{W}| = (q^2+1)q^5 + q^2 + 1 = (q^2+1)(q^5+1);$$

in particular,  $\mathcal{W}$  is a hypersurface of  $\bar{\Sigma}_4$  of degree  $q+1$  such that there exists a plane of  $\bar{\Sigma}_4$  meeting  $\mathcal{W}$  in just one line (such planes exist in  $\bar{\Sigma}_3$ ). Also suppose  $\mathcal{W}$  to contain planes and let  $\Pi''' \subseteq \mathcal{W}$  be such a plane. Since  $\Sigma_4^i \cap \mathcal{W}$  does not contain planes, all  $\Sigma_4^i$  meet  $\Pi'''$  in a line  $t_i$ . Also  $\Pi'''$  must be contained in  $\bigcup_{i=1}^{q^2+1} t_i$ . This implies that the set  $\{t_i\}_{i=1, \dots, q^2+1}$  consists of  $q^2+1$  lines through a point  $P \in \Pi \setminus \{P_1\}$ .

Furthermore each line  $t_i$  passing through  $P$  must meet the radical line  $\ell_i$  of the Hermitian cone  $\mathcal{S} \cap \Sigma_4^i$  and this forces  $P$  to coincide with  $P_1$ , a contradiction. It follows that  $\mathcal{W}$  does not contain planes.

So by the characterization of  $\mathcal{H}(4, q^2)$  of [3] we have that  $\mathcal{W}$  is a Hermitian variety  $\mathcal{H}(4, q^2)$ .

We also have that  $|\mathcal{S} \cap \Sigma_5| = |P_1 \mathcal{H}(4, q^2)|$ . Let now  $r$  be any line of  $\mathcal{H}(4, q^2) = \mathcal{S} \cap \bar{\Sigma}_4$  and let  $\Theta$  be the plane  $\langle r, P_1 \rangle$ . The plane  $\Theta$  meets  $\Sigma_4^i$  in a line  $q_i \subseteq \mathcal{S}$  for each  $i = 1, \dots, q^2 + 1$  and these lines are concurrent in  $P_1$ . It follows that all the points of  $\Theta$  are in  $\mathcal{S}$ . This completes the proof for the current case and shows that  $\mathcal{S} \cap \Sigma_5$  is a Hermitian cone  $P_1 \mathcal{H}(4, q^2)$ .

- (b) All planes  $\Pi''$  with  $m \subseteq \Pi'' \subseteq \mathcal{S} \cap \Sigma_5$  are contained in  $\Sigma_4^i$  for some  $i = 1, \dots, q^2 + 1$ . We claim that this case cannot happen. We can suppose without loss of generality  $m \cap \ell_1 = P_1$  and  $P_1 \notin \ell_i$  for all  $i = 2, \dots, q^2 + 1$ . Since the intersection of the subspaces  $\Sigma_4^i$  is  $\Sigma_3$ , there is exactly one plane through  $m$  in  $\Sigma_5$  which is  $(q^4 + q^2 + 1)$ -secant, namely the plane  $\langle \ell_1, m \rangle$ . Furthermore, in  $\Sigma_4^1$  there are  $q^4$  planes through  $m$  which are  $(q^3 + q^2 + 1)$ -secant and  $q^2$  planes which are  $(q^2 + 1)$ -secant. We can provide an upper bound to the points of  $\mathcal{S} \cap \Sigma_5$  by counting the number of points of  $\mathcal{S} \cap \Sigma_5$  on planes in  $\Sigma_5$  through  $m$  and observing that a plane through  $m$  not in  $\Sigma_5$  and not contained in  $\mathcal{S}$  has at most  $q^3 + q^2 + 1$  points in common with  $\mathcal{S} \cap \Sigma_5$ . So

$$|\mathcal{S} \cap \Sigma_5| \leq q^6 \cdot q^3 + q^7 + q^4 + q^2 + 1.$$

As  $|\mathcal{S} \cap \Sigma_5| = q^9 + q^7 + q^4 + q^2 + 1$ , all planes through  $m$  which are neither  $(q^4 + q^2 + 1)$ -secant nor  $(q^2 + 1)$ -secant are  $(q^3 + q^2 + 1)$ -secant. That is to say that all of these planes meet  $\mathcal{S}$  in a curve of degree  $q + 1$  which must split into  $q + 1$  lines through a point because of Lemma 2.1.

Take now  $P_2 \in \Sigma_4^2 \cap \mathcal{S}$  and consider the plane  $\Xi := \langle m, P_2 \rangle$ . The line  $\langle P_1, P_2 \rangle$  is contained in  $\Sigma_4^2$ ; so it must be a  $(q + 1)$ -secant, as it does not meet the vertex line  $\ell_2$  of  $C_2$  in  $\Sigma_4^2$ . Now,  $\Xi$  meets every of  $\Sigma_4^i$  for  $i = 2, \dots, q^2 + 1$  in a line through  $P_1$  which is either a 1-secant or a  $q + 1$ -secant; so

$$|\mathcal{S} \cap \Xi| \leq q^2(q) + q^2 + 1 = q^3 + q^2 + 1.$$

It follows that  $|\mathcal{S} \cap \Xi| = q^3 + q^2 + 1$  and  $\mathcal{S} \cap \Xi$  is a set of  $q + 1$  lines all through the point  $P_1$ . This contradicts our previous construction.

□

**Lemma 3.7.** *Every hyperplane of  $\text{PG}(6, q^2)$  meets  $\mathcal{S}$  either in a non-singular Hermitian variety  $\mathcal{H}(5, q^2)$  or in a cone with vertex a point over a Hermitian hypersurface  $\mathcal{H}(4, q^2)$ .*

*Proof.* Let  $\Sigma_3$  be a solid satisfying condition (S4). Denote by  $\Lambda$  a hyperplane of  $\text{PG}(6, q^2)$ . If  $\Lambda$  contains  $\Sigma_3$  then, from Lemma 3.6 it follows that  $\Lambda \cap \mathcal{S}$  is a Hermitian cone  $P\mathcal{H}(4, q^2)$ .

Now assume that  $\Lambda$  does not contain  $\Sigma_3$ . Denote by  $S_5^j$ , with  $j = 1, \dots, q^2 + 1$  the  $q^2 + 1$  hyperplanes through  $\Sigma_4^1$ , where as before,  $\Sigma_4^1$  is a 4-space containing  $\Sigma_3$ . By Lemma 3.6 again we get that  $S_5^j \cap \mathcal{S} = P^j \mathcal{H}(4, q^2)$ . We count the number of rational points of  $\Lambda \cap \mathcal{S}$  by studying the intersections of  $S_5^j \cap \mathcal{S}$  with  $\Lambda$  for all  $j \in \{1, \dots, q^2 + 1\}$ . Setting  $\mathcal{W}_j := S_5^j \cap \mathcal{S} \cap \Lambda$ ,  $\Omega := \Sigma_4^1 \cap \mathcal{S} \cap \Lambda$  then

$$|\mathcal{S} \cap \Lambda| = \sum_j |\mathcal{W}_j \setminus \Omega| + |\Omega|.$$

If  $\Pi$  is a plane of  $\Lambda$  then  $\Omega$  consists of  $q + 1$  planes of a pencil. Otherwise let  $m$  be the line in which  $\Lambda$  meets the plane  $\Pi$ . Then  $\Omega$  is either a Hermitian cone  $P_0 \mathcal{H}(2, q^2)$ , or  $q + 1$



planes of a pencil, according as the vertex  $P^j \in \Pi$  is an external point with respect to  $m$  or not.

In the former case  $\mathcal{W}_j$  is a non singular Hermitian variety  $\mathcal{H}(4, q^2)$  and thus  $|\mathcal{S} \cap \Lambda| = (q^2 + 1)(q^7) + q^5 + q^2 + 1 = q^9 + q^7 + q^5 + q^2 + 1$ .

In the case in which  $\Omega$  consists of  $q+1$  planes of a pencil then  $\mathcal{W}_j$  is either a  $P_0\mathcal{H}(3, q^2)$  or a Hermitian cone with vertex a line  $\ell$  and basis a Hermitian curve  $\mathcal{H}(2, q^2)$ .

If there is at least one index  $j$  such that  $\mathcal{W}_j = \ell\mathcal{H}(2, q^2)$ , then there must be a 3-dimensional space  $\Sigma'_3$  of  $S'_5 \cap \Lambda$  meeting  $\mathcal{S}$  in a generator. Hence, from Lemma 3.6 we get that  $\mathcal{S} \cap \Lambda$  is a Hermitian cone  $P'\mathcal{H}(4, q^2)$ .

Assume that for all  $j \in \{1, \dots, q^2 + 1\}$ ,  $\mathcal{W}_j$  is a  $P_0\mathcal{H}(3, q^2)$ . In this case

$$|\mathcal{S} \cap \Lambda| = (q^2 + 1)q^7 + (q + 1)q^4 + q^2 + 1 = q^9 + q^7 + q^5 + q^4 + q^2 + 1 = |\mathcal{H}(5, q^2)|.$$

We are going to prove that the intersection numbers of  $\mathcal{S}$  with hyperplanes are only two that is  $q^9 + q^7 + q^5 + q^4 + q^2 + 1$  or  $q^9 + q^7 + q^4 + q^2 + 1$ .

Denote by  $x_i$  the number of hyperplanes meeting  $\mathcal{S}$  in  $i$  rational points with  $i \in \{q^9 + q^7 + q^4 + q^2 + 1, q^9 + q^7 + q^5 + q^2 + 1, q^9 + q^7 + q^5 + q^4 + q^2 + 1\}$ . Double counting arguments give the following equations for the integers  $x_i$ :

$$\begin{cases} \sum_i x_i = q^{12} + q^{10} + q^8 + q^6 + q^4 + q^2 + 1 \\ \sum_i ix_i = |\mathcal{S}|(q^{10} + q^8 + q^6 + q^4 + q^2 + 1) \\ \sum_{i=1} i(i-1)x_i = |\mathcal{S}|(|\mathcal{S}| - 1)(q^8 + q^6 + q^4 + q^2 + 1). \end{cases} \quad (3.2)$$

Solving (3.2) we obtain  $x_{q^9+q^7+q^5+q^2+1} = 0$ . In the case in which  $|\mathcal{S} \cap \Lambda| = |\mathcal{H}(5, q^2)|$ , since  $\mathcal{S} \cap \Lambda$  is an algebraic hypersurface of degree  $q+1$  not containing 3-spaces, from [19, Theorem 4.1] we get that  $\mathcal{S} \cap \Lambda$  is a Hermitian variety  $\mathcal{H}(5, q^2)$  and this completes the proof.  $\square$

*Proof of Theorem 1.1.* The first part of Theorem 1.1 follows from Lemma 3.4. From Lemma 3.7,  $\mathcal{S}$  has the same intersection numbers with respect to hyperplanes and 4-spaces as a non-singular Hermitian variety of  $\text{PG}(6, q^2)$ , hence Lemma 2.5 applies and  $\mathcal{S}$  turns out to be a  $\mathcal{H}(6, q^2)$ .  $\square$

**Remark 3.8.** The characterization of the non-singular Hermitian variety  $\mathcal{H}(4, q^2)$  given in [3] is based on the property that a given hypersurface is a blocking set with respect to lines of  $\text{PG}(4, q^2)$ , see [3, Lemma 3.1]. This lemma holds when  $q > 3$ . Since Lemma 3.2 extends the same property to the case  $q = 3$  it follows that the result stated in [3] is also valid in  $\text{PG}(4, 3^2)$ .

## 4 Conjecture


We propose a conjecture for the general  $2n$ -dimensional case.


*Let  $\mathcal{S}$  be a hypersurface of  $\text{PG}(2d, q^2)$ ,  $q > 2$ , defined over  $\text{GF}(q^2)$ , not containing  $d$ -dimensional projective subspaces. If the degree of  $\mathcal{S}$  is  $q+1$  and the number of its rational points is  $|\mathcal{H}(2d, q^2)|$ , then every  $d$ -dimensional subspace of  $\text{PG}(2d, q^2)$  meets  $\mathcal{S}$  in at least  $\theta_{q^2}(d-1) := (q^{2d-2} - 1)/(q^2 - 1)$  rational points. If there is at least a  $d$ -dimensional*


subspace  $\Sigma_d$  such that  $|\Sigma_d \cap \mathcal{S}| = |\text{PG}(d-1, q^2)|$ , then  $\mathcal{S}$  is a non-singular Hermitian variety of  $\text{PG}(2d, q^2)$ .

Lemma 3.1 and Lemma 3.2 can be a starting point for the proof of this conjecture since from them we get that  $\mathcal{S}$  is a blocking set with respect to lines of  $\text{PG}(2d, q^2)$ .

## ORCID iDs

Angela Aguglia  <https://orcid.org/0000-0001-6854-8679>

Luca Giuzzi  <https://orcid.org/0000-0003-3975-7281>

Masaaki Homma  <https://orcid.org/0000-0003-4568-6408>

## References

- [1] A. Aguglia, Quasi-Hermitian varieties in  $\text{PG}(r, q^2)$ ,  $q$  even, *Contrib. Discrete Math.* **8** (2013), 31–37.
- [2] A. Aguglia, A. Cossidente and G. Korchmáros, On quasi-Hermitian varieties, *J. Combin. Des.* **20** (2012), 433–447, doi:10.1002/jcd.21317.
- [3] A. Aguglia and F. Pavese, On non-singular Hermitian varieties of  $\text{PG}(4, q^2)$ , *Discrete Math.* **343** (2020), 111634, 5, doi:10.1016/j.disc.2019.111634.
- [4] H. Borges and M. Homma, Points on singular Frobenius nonclassical curves, *Bull. Braz. Math. Soc. (N.S.)* **48** (2017), 93–101, doi:10.1007/s00574-016-0008-6.
- [5] R. C. Bose and R. C. Burton, A characterization of flat spaces in a finite geometry and the uniqueness of the Hamming and the MacDonald codes, *J. Combinatorial Theory* **1** (1966), 96–104.
- [6] R. C. Bose and I. M. Chakravarti, Hermitian varieties in a finite projective space  $\text{PG}(N, q^2)$ , *Canadian J. Math.* **18** (1966), 1161–1182, doi:10.4153/CJM-1966-116-0.
- [7] S. De Winter and J. Schillewaert, Characterizations of finite classical polar spaces by intersection numbers with hyperplanes and spaces of codimension 2, *Combinatorica* **30** (2010), 25–45, doi:10.1007/s00493-010-2441-2.
- [8] J. W. P. Hirschfeld, L. Storme, J. A. Thas and J. F. Voloch, A characterization of Hermitian curves, *J. Geom.* **41** (1991), 72–78, doi:10.1007/BF01258509.
- [9] J. W. P. Hirschfeld and J. A. Thas, *General Galois geometries*, Springer Monographs in Mathematics, Springer, London, 2016, doi:10.1007/978-1-4471-6790-7.
- [10] M. Homma and S. J. Kim, Around Sziklai’s conjecture on the number of points of a plane curve over a finite field, *Finite Fields Appl.* **15** (2009), 468–474, doi:10.1016/j.ffa.2009.02.008.
- [11] M. Homma and S. J. Kim, An elementary bound for the number of points of a hypersurface over a finite field, *Finite Fields Appl.* **20** (2013), 76–83, doi:10.1016/j.ffa.2012.11.002.
- [12] M. Homma and S. J. Kim, The characterization of Hermitian surfaces by the number of points, *J. Geom.* **107** (2016), 509–521, doi:10.1007/s00022-015-0283-1.
- [13] M. Homma and S. J. Kim, Number of points of a nonsingular hypersurface in an odd-dimensional projective space, *Finite Fields Appl.* **48** (2017), 395–419, doi:10.1016/j.ffa.2017.08.011.
- [14] R. Schoof, Nonsingular plane cubic curves over finite fields, *J. Combin. Theory Ser. A* **46** (1987), 183–211, doi:10.1016/0097-3165(87)90003-3.
- [15] B. Segre, Le geometrie di Galois, *Ann. Mat. Pura Appl. (4)* **48** (1959), 1–96, doi:10.1007/BF02410658.

- [16] B. Segre, *Forme e geometrie hermitiane, con particolare riguardo al caso finito*, *Ann. Mat. Pura Appl. (4)* **70** (1965), 1–201, doi:10.1007/BF02410088.
- [17] J.-P. Serre, *Lettre à M. Tsfasman*, *Astérisque* (1991), 11, 351–353 (1992), journées Arithmétiques, 1989 (Luminy, 1989).
- [18] K.-O. Stöhr and J. F. Voloch, *Weierstrass points and curves over finite fields*, *Proc. London Math. Soc. (3)* **52** (1986), 1–19, doi:10.1112/plms/s3-52.1.1.
- [19] A. Weil, *Sur les courbes algébriques et les variétés qui s'en déduisent*, volume 7 of *Publ. Inst. Math. Univ. Strasbourg*, Hermann et Cie., Paris, 1948.