

Paired domination stability in graphs

Aleksandra Gorzkowska 

*AGH University, Department of Discrete Mathematics,
al. Mickiewicza 30, 30-059 Krakow, Poland*

Michael A. Henning 

*Department of Mathematics and Applied Mathematics, University of Johannesburg,
Auckland Park, 2006 South Africa*

Monika Piłśniak 

*AGH University, Department of Discrete Mathematics,
al. Mickiewicza 30, 30-059 Krakow, Poland*

Elżbieta Tumidajewicz * 

*AGH University, Department of Discrete Mathematics,
al. Mickiewicza 30, 30-059 Krakow, Poland, and
Department of Mathematics and Applied Mathematics, University of Johannesburg,
Auckland Park, 2006 South Africa*

Received 29 December 2020, accepted 16 July 2021, published online 27 May 2022

Abstract

A set S of vertices in a graph G is a paired dominating set if every vertex of G is adjacent to a vertex in S and the subgraph induced by S contains a perfect matching (not necessarily as an induced subgraph). The paired domination number, $\gamma_{\text{pr}}(G)$, of G is the minimum cardinality of a paired dominating set of G . A set of vertices whose removal from G produces a graph without isolated vertices is called a non-isolating set. The minimum cardinality of a non-isolating set of vertices whose removal decreases the paired domination number is the γ_{pr}^- -stability of G , denoted $\text{st}_{\gamma_{\text{pr}}}^-(G)$. The paired domination stability of G is the minimum cardinality of a non-isolating set of vertices in G whose removal changes the paired domination number. We establish properties of paired domination stability in graphs. We prove that if G is a connected graph with $\gamma_{\text{pr}}(G) \geq 4$, then $\text{st}_{\gamma_{\text{pr}}}^-(G) \leq 2\Delta(G)$ where $\Delta(G)$ is the maximum degree in G , and we characterize the infinite family of trees that achieve equality in this upper bound.

Keywords: Paired domination, paired domination stability.

*Corresponding author.

1 Introduction

In 1983 Bauer, Harary, Nieminen and Suffel [3] introduced and studied the concept of domination stability in graphs. Stability for other domination type parameters has been studied in the literature. For example, total domination stability, 2-rainbow domination stability, exponential domination stability, Roman domination stability are studied in [1, 2, 12, 15, 16], among other papers. In this paper we study the paired version of domination stability.

Let $G = (V, E)$ be a graph with vertex set $V = V(G)$ and edge set $E = E(G)$. Two vertices u and v are *neighbors* if they are adjacent, that is, if $uv \in E$. A *dominating set* of G is a set D of vertices such that every vertex in $V(G) \setminus D$ has a neighbor in D . The minimum cardinality of a dominating set is the *domination number*, $\gamma(G)$, of G . Domination is well studied in the literature. A recent book on domination in graphs can be found in [10]. A small sample of papers on domination critical graphs can be found in [3, 4, 5, 6, 9, 17, 18]. Adopting the notation coined by Bauer et al. [3], the γ^- -stability (γ^+ -stability, resp.) of G , denoted by $\gamma^-(G)$ ($\gamma^+(G)$, resp.), is the minimum number of vertices whose removal decreases (increases, resp.) the domination number. The minimum number of vertices whose removal decreases or increases the domination number is the *domination stability*, $\text{st}_\gamma(G)$, of G , and so $\text{st}_\gamma(G) = \min\{\gamma^-(G), \gamma^+(G)\}$.

We refer to a graph without isolated vertices as an *isolate-free graph*. Unless otherwise stated, let G be an isolate-free graph. A *total dominating set*, abbreviated TD-set, of G is a set D of vertices of G such that every vertex, including vertices in the set D , has a neighbor in D . The minimum cardinality of a TD-set of G is the *total domination number*, $\gamma_t(G)$, of G . We call a TD-set of G of cardinality $\gamma_t(G)$ a γ_t -set of G . A vertex v is *totally dominated* by a set D in G if the vertex v has a neighbor in D . We refer the reader to the book [14] for fundamental concepts on total domination in graphs. Total domination critical graphs are studied, for example, in [7, 13]. The total version of domination stability was first studied by Henning and Krzywkowski [12].

A *paired dominating set*, abbreviated PD-set, of an isolate-free graph G is a dominating set S of G with the additional property that the subgraph $G[S]$ induced by S contains a perfect matching M (not necessarily induced). With respect to the matching M , two vertices joined by an edge of M are *paired* and are called *partners* in S . The *paired domination number*, $\gamma_{\text{pr}}(G)$, of G is the minimum cardinality of a PD-set of G . We call a PD-set of G of cardinality $\gamma_{\text{pr}}(G)$ a γ_{pr} -set of G . We note that the paired domination number $\gamma_{\text{pr}}(G)$ is an even integer. For a recent survey on paired domination in graphs, we refer the reader to the book chapter [8].

Every PD-set is a TD-set, implying that $\gamma_t(G) \leq \gamma_{\text{pr}}(G)$. A *non-isolating set* of vertices in G is a set $S \subseteq V$ such that the graph $G - S$ is isolate-free, where $G - S$ is the graph obtained from G by removing S and all edges incident with vertices in S . Let $\text{NI}(G)$ denote the set of all non-isolating sets of vertices of G .

Adopting the standard notation for domination stability given in [3, 12], the γ_{pr}^- -stability

(resp., γ_{pr}^+ -stability) of G , denoted by $st_{\gamma_{pr}}^-(G)$ (resp., $st_{\gamma_{pr}}^+(G)$) is the minimum cardinality of a set in $NI(G)$ whose removal decreases (increases, resp.) the paired domination number. Thus,

$$st_{\gamma_{pr}}^-(G) = \min_{S \in NI(G)} \{ |S| : \gamma_{pr}(G - S) < \gamma_{pr}(G) \}$$

and

$$st_{\gamma_{pr}}^+(G) = \min_{S \in NI(G)} \{ |S| : \gamma_{pr}(G - S) > \gamma_{pr}(G) \}.$$

If there is no set in $NI(G)$ whose removal increases the paired domination number, then we define $st_{\gamma_{pr}}^+(G) = \infty$. For example, $st_{\gamma_{pr}}^-(P_5) = 1$ while $st_{\gamma_{pr}}^+(P_5) = \infty$. The *paired domination stability*, $st_{\gamma_{pr}}(G)$, of G is the minimum cardinality of a set in $NI(G)$ whose removal increases or decreases the paired domination number. Thus,

$$st_{\gamma_{pr}}(G) = \min_{S \in NI(G)} \{ |S| : \gamma_{pr}(G - S) \neq \gamma_{pr}(G) \} = \min \{ st_{\gamma_{pr}}^-(G), st_{\gamma_{pr}}^+(G) \}.$$

Let G be a graph and let $S \in NI(G)$. If $\gamma_{pr}(G - S) < \gamma_{pr}(G)$ and $|S| = st_{\gamma_{pr}}^-(G)$, then we call S a $st_{\gamma_{pr}}^-$ -set of G . If $\gamma_{pr}(G - S) > \gamma_{pr}(G)$ and $|S| = st_{\gamma_{pr}}^+(G)$, then we call S a $st_{\gamma_{pr}}^+$ -set of G . If $\gamma_{pr}(G - S) \neq \gamma_{pr}(G)$ and $|S| = st_{\gamma_{pr}}(G)$, then we call S a $st_{\gamma_{pr}}$ -set of G .

Defining the *null graph* K_0 , which has no vertices, as a graph, we have the following results due to Bauer et al. [3] and Rad et al. [15] for the γ^- -stability of a graph.

Theorem 1.1 ([3, 15]). *If G is an isolate-free graph of order n , then the following holds.*

- (a) $st_{\gamma}(G) \leq \delta(G) + 1$.
- (b) *If $G \not\cong K_n$, then $st_{\gamma}(G) \leq n - 1$.*

Considering the null graph, the paired domination stability of a non-trivial graph is always defined. If G is a graph of order n and $\gamma_{pr}(G) = 2$, then $st_{\gamma_{pr}}^-(G) = n$ since removing all vertices from the graph G produces the null graph with paired domination number zero.

For notation and graph theory terminology we generally follow [14]. In particular, for $r, s \geq 1$, a *double star* $S(r, s)$ is the tree with exactly two vertices that are not leaves, one of which has r leaf-neighbors and the other s leaf-neighbors. A *rooted tree* is a tree T in which we specify one vertex r called the *root*. For each vertex v of T different from r , its *parent* is the neighbor of v on the unique (r, v) -path, while every other neighbor of v is a *child* of v in T . If w is a vertex of T different from v and the (unique) (r, w) -path contains v , then w is a *descendant* of v in T . We note that every child of v is a descendant of v . The *diameter* $diam(G)$ of G is the maximum distance among all pairs of vertices of G . A *diametral path* in G is a shortest path between two vertices in G of length equal to $diam(G)$. For an integer $k \geq 1$, $[k] = \{1, \dots, k\}$.

2 Main results

Our first aim is to show that the paired domination stability of a graph can be very different from its total domination stability studied in [12].

Theorem 2.1. *For $k \geq 1$ an arbitrary integer, the following holds.*

- (a) *There exist connected graphs G such that $st_{\gamma_{pr}^-}(G) - st_{\gamma_t^-}(G) = k$.*
- (b) *There exist connected graphs H such that $st_{\gamma_t^-}(H) - st_{\gamma_{pr}^-}(H) = k$.*

Our second aim is to establish properties of paired domination stability in graphs. Thereafter, we establish upper bounds on the paired domination stability and the γ_{pr}^- -stability of a graph. For this purpose, we shall need the following family of trees defined by Henning and Krzywkowski [12]. For integers $k \geq 2$ and $\Delta \geq 2$, the authors in [12] define $T_{k,\Delta}$ as the “graph obtained from the disjoint union of k double stars $S(\Delta - 1, \Delta - 1)$ by adding $k - 1$ edges between the leaves of these double stars so that the resulting graph is a tree with maximum degree Δ .” Let $\mathcal{F}_{k,\Delta}$ be the family of all such trees $T_{k,\Delta}$, and let

$$\mathcal{F}_\Delta = \bigcup_{k \geq 2} \mathcal{F}_{k,\Delta}.$$

The following result establishes an upper bound on the γ_{pr}^- -stability of a tree, and characterizes the trees with maximum possible γ_{pr}^- -stability.

Theorem 2.2. *If T is a tree with maximum degree Δ satisfying $\gamma_{pr}(T) \geq 4$, then the following hold.*

- (a) $st_{\gamma_{pr}^-}(T) \leq 2\Delta$, with equality if and only if $T \in \mathcal{F}_\Delta$.
- (b) $st_{\gamma_{pr}^-}(T) \leq 2\Delta - 1$, and this bound is sharp for all $\Delta \geq 2$.

For general graphs, we establish the following upper bound on the γ_{pr}^- -stability in terms of the maximum degree of the graph.

Theorem 2.3. *If G is a connected graph with $\gamma_{pr}(G) \geq 4$, then $st_{\gamma_{pr}^-}(G) \leq 2\Delta(G)$, and this bound is sharp.*

As an immediate consequence of Theorem 2.3, we have the following upper bound on the paired domination stability of a graph.

Corollary 2.4. *If G is a connected graph with $\gamma_{pr}(G) \geq 4$, then $st_{\gamma_{pr}^-}(G) \leq 2\Delta(G)$.*

3 Paired stability versus domination and total stability

In this section, we show that paired domination stability and the domination stability of a graph can be very different. By Theorem 1.1, for every nontrivial graph G , we have $st_\gamma(G) \leq \delta(G) + 1$. In particular, $st_\gamma(T) \leq 2$ for every nontrivial tree T . This is in contrast to the paired domination stability, where for any given $\Delta \geq 2$, we show that there exist a family of trees T with maximum degree Δ satisfying $st_{\gamma_{pr}^-}(T) = 2\Delta - 1$.

For $\Delta = 2$, the authors in [12] define \mathcal{H}_Δ as the family of all paths of order at least 7 and congruent to 3 modulo 4, that is, $\mathcal{H}_\Delta = \{P_n \mid n \equiv 3 \pmod{4} \text{ and } n \geq 7\}$. For integers $\Delta \geq 3$ and $\Delta \geq k \geq 2$, they define $H_{k,\Delta}$ as the graph “obtained from the disjoint union of k double stars $S(\Delta - 1, \Delta - 1)$ by selecting one leaf from each double star and identifying these k leaves into one new vertex” and they define the family

$$\mathcal{H}_\Delta = \bigcup_{k \geq 2} H_{k,\Delta}.$$

We determine next the paired domination stability of a tree in the family \mathcal{H}_Δ .

Proposition 3.1. *For $\Delta \geq 3$, if $T \in \mathcal{H}_\Delta$, then $\text{st}_{\gamma_{\text{pr}}}(T) = 2\Delta - 1$.*

Proof. For integers $\Delta \geq k \geq 2$ where $\Delta \geq 3$, consider a tree $T \in \mathcal{H}_{k,\Delta}$. By definition of the family $\mathcal{H}_{k,\Delta}$, the tree T is constructed from the disjoint union of k double stars S_1, \dots, S_k , each isomorphic to $S(\Delta - 1, \Delta - 1)$, by selecting one leaf from each double star and identifying these k chosen leaves into one new vertex, which we call v_c . Let x_i and y_i be the two central vertices of the double star S_i for $i \in [k]$, where x_i is adjacent to v_c in T . Let $D = \cup_{i=1}^k \{x_i, y_i\}$. Since $\Delta \geq 3$, every vertex in D is a support vertex of T , implying that every PD-set in T contains the set D and therefore $\gamma_{\text{pr}}(T) \geq |D| = 2k$. Since the set D is a PD-set of T (with the vertices x_i and y_i paired for all $i \in [k]$), we have $\gamma_{\text{pr}}(T) \leq |D| = 2k$. Consequently, $\gamma_{\text{pr}}(T) = 2k$ and D is the unique γ_{pr} -set of T .

Let S be a $\text{st}_{\gamma_{\text{pr}}}$ -set of T . Thus, S is a set in $\text{NI}(T)$ with $|S| = \text{st}_{\gamma_{\text{pr}}}(T)$ satisfying $\gamma_{\text{pr}}(T - S) \neq \gamma_{\text{pr}}(T) = 2k$. We show that $|S| \geq 2\Delta - 1$. Suppose, to the contrary, that $|S| \leq 2\Delta - 2$. If the set S contains both x_i and y_i for some $i \in [k]$, then since S is a non-isolating set of T every leaf neighbor of x_i and y_i is also in S , implying that $|S| \geq 2\Delta - 1$, a contradiction. Hence, the set S contains at most one of x_i and y_i for every $i \in [k]$. Let D^* be a γ_{pr} -set of $T - S$, and so $|D^*| \neq 2k$.

Suppose that $v_c \in S$. In this case, if $|S| = 1$, then the paired domination numbers of T and $T - S$ are the same, a contradiction. Hence, $|S| \geq 2$. If neither x_i nor y_i belong to S for some $i \in [k]$, then by the minimality of the non-isolating set S , no vertex of T_i different from v_c belongs to S , and so $|D^* \cap V(T_i)| = 2$. If S contains y_i but not x_i for some $i \in [k]$, then every leaf neighbor of y_i is in S and by the minimality of the set S , no leaf neighbor of x_i belongs to S , and so $|D^* \cap V(T_i)| = 2$. Analogously, if S contains x_i but not y_i for some $i \in [k]$, then $|D^* \cap V(T_i)| = 2$. This is true for all $i \in [k]$, implying that $|D^*| = \sum_{i=1}^k |D^* \cap V(T_i)| = 2k$, a contradiction. Hence, $v_c \notin S$.

As observed earlier, the set S contains at most one of x_i and y_i for every $i \in [k]$. If $y_i \in S$ and $y_j \in S$ for some $i, j \in [k]$ where $i \neq j$, then $|S| \geq 2\Delta$, a contradiction. If $y_i \in S$ and $x_j \in S$ for some $i, j \in [k]$ where $i \neq j$, then $|S| \geq 2\Delta - 1$, a contradiction. If $x_i \in S$ and $x_j \in S$ for some $i, j \in [k]$ where $i \neq j$, then $|S| \geq 2\Delta - 2$. In this case, by the minimality of S we have $S = (N[x_i] \cup N[x_j]) \setminus \{v_c, y_i, y_j\}$ and $|S| = 2\Delta - 2$. But then $T - S$ consists of three components, namely two stars isomorphic to $K_{1,\Delta-1}$ and one component belonging to the family $T \in \mathcal{H}_{k-2,\Delta}$ with paired domination number $2(k-2)$. Thus, $\gamma_{\text{pr}}(T - S) = 2 + 2 + 2(k-2) = 2k$, a contradiction. Therefore, $\text{st}_{\gamma_{\text{pr}}}(T) = |S| \geq 2\Delta - 1$, as claimed.

Conversely, if we take $S = N(x_1) \cup N(y_1) \setminus \{v_c\}$, then $S \in \text{NI}(T)$ and $T - S \in \mathcal{H}_{k-1,\Delta}$. Thus, $\gamma_{\text{pr}}(T - S) = 2(k-1) < \gamma_{\text{pr}}(T)$, and so $\text{st}_{\gamma_{\text{pr}}}(T) \leq \text{st}_{\gamma_{\text{pr}}}^-(T) \leq |S| = 2\Delta - 1$. Consequently, $\text{st}_{\gamma_{\text{pr}}}(T) = \text{st}_{\gamma_{\text{pr}}}^-(T) = 2\Delta - 1$. \square

As observed earlier, $\text{st}_\gamma(T) \leq 2$ for every nontrivial tree T . By Proposition 3.1, paired domination stability therefore differs significantly from domination stability. We show next that the paired domination stability and the total domination stability of a graph can also be very different.

Proposition 3.2. *For $k \geq 1$ an integer, there exist trees T such that $\text{st}_{\gamma_{\text{pr}}}^-(T) - \text{st}_{\gamma_t}^-(T) = k$.*

Proof. Let $k \geq 1$ be a given integer, and let $T = T_k$ be obtained from a path P_5 given by $v_1v_2v_3v_4v_5$ by attaching k leaf neighbors to each of v_1, v_2 and v_3 (see Figure 1). We

note that $\{v_1, v_2, v_3, v_4\}$ is the unique γ_t -set of T and the unique γ_{pr} -set of T . In particular, $\gamma_t(T) = \gamma_{pr}(T) = 4$. If $S = \{v_5\}$, then the set S is a non-isolating set of T and $\gamma_t(T - S) = |\{v_1, v_2, v_3\}| = 3 < \gamma_t(T)$, implying that $st_{\gamma_t}^-(T) = 1$.

We show next that $st_{\gamma_{pr}}^-(T) = k + 1$. Let S be a non-isolating set of T such that $\gamma_{pr}(T - S) < \gamma_{pr}(T)$. We show that $|S| \geq k + 1$. Suppose, to the contrary, that $|S| \leq k$. Let D be a γ_{pr} -set of $T - S$, and so $|D| = \gamma_{pr}(T - S) = 2$. Let L_i denote the set of leaf neighbors of v_i for $i \in [4]$. If $v_i \in S$ for some $i \in [3]$, then S contains all k leaf neighbors of v_i , and so $|S| \geq k + 1$, a contradiction. Hence, $S \cap \{v_1, v_2, v_3\} = \emptyset$. If $\{v_1, v_3\} \subset D$, then $|D| \geq 4$, a contradiction. If $v_1 \notin D$, then $L_1 \subseteq S$, implying that $S = L_1$ and $|S| = k$. However in this case, $\{v_2, v_3, v_4\} \subset D$. If $v_3 \notin D$, then $L_3 \subseteq S$, implying that $S = L_3$ and $|S| = k$. However in this case, $\{v_1, v_2, v_4\} \subset D$. In both cases, $|D| \geq 4$, a contradiction. Therefore, $|S| \geq k + 1$, implying that $st_{\gamma_{pr}}^-(T) \geq k + 1$. Conversely, if $S = L_1 \cup L_4$, then S is a non-isolating set of T such that $\gamma_{pr}(T - S) = |\{v_2, v_3\}| < \gamma_{pr}(T)$, implying that $st_{\gamma_{pr}}^-(T) \leq |S| = k + 1$. Consequently, $st_{\gamma_{pr}}^-(T) = k + 1$. Thus, $st_{\gamma_{pr}}^-(T) - st_{\gamma_t}^-(T) = k$. \square




Figure 1: A tree from the family T_k in the proof of Proposition 3.2.

Proposition 3.3. For $k \geq 1$ an integer, there exist trees T such that $st_{\gamma_t}^-(T) - st_{\gamma_{pr}}^-(T) = k$.

Proof. Let $k \geq 1$ be a given integer, and let $\ell \geq 2k + 1$ be an integer. For $i \in [k]$, let Q_i be obtained from a path $v_{i_1}v_{i_2}v_{i_3}v_{i_4}v_{i_5}$ by attaching ℓ leaf neighbors to each of v_{i_3}, v_{i_4} and v_{i_5} , and let L_{i_3}, L_{i_4} and L_{i_5} be the resulting sets of leaf neighbors of v_{i_3}, v_{i_4} and v_{i_5} , respectively. Let Q be obtained from a path $v_1v_2v_3$ by attaching ℓ leaf neighbors to each of v_1 and v_2 , and attaching k leaf neighbors to v_3 . Let L_i be the resulting set of leaf neighbors of v_i for $i \in [3]$. Let T be obtained from the disjoint union of the paths Q, Q_1, \dots, Q_k by adding the k edges $v_3v_{i_1}$ for $i \in [k]$. Let A be the set of support vertices of T , and so $|A| = 3(k + 1)$.

Every TD-set of T contains all its support vertices, implying that $\gamma_t(T) \geq |A|$. Since the set A is a TD-set of T , we have $\gamma_t(T) \leq |A|$. Consequently, $\gamma_t(T) = |A| = 3(k + 1)$. Every PD-set of T contains the set A and at least one additional vertex from each path Q_i that is a neighbor of v_{i_3} or v_{i_5} for $i \in [k]$, and at least one additional vertex that is a neighbor of v_1 or v_3 since the vertices of every PD-set are paired, implying that $\gamma_{pr}(T) = |A| + k + 1 = 4(k + 1)$.

Let S be a non-isolating set of T such that $\gamma_{pr}(T - S) < \gamma_{pr}(T)$. If $|S| < k$, then every support vertex of T remains a support vertex of $T - S$, implying that $\gamma_{pr}(T - S) \geq \gamma_{pr}(T)$, a contradiction. Hence, $|S| \geq k$. Conversely, if $S^* = L_3$, then the set $A \setminus \{v_3\}$ of all support vertices of $T - S^*$, together with the vertices v_{i_2} for $i \in [k]$, form a PD-set of $T - S^*$, implying that $\gamma_{pr}(T - S^*) \leq 4k + 2 < 4k + 4 = \gamma_{pr}(T)$. Hence, $st_{\gamma_{pr}}^-(T) \leq |S^*| = k$. Consequently, $st_{\gamma_{pr}}^-(T) = k$.

We show next that $st_{\gamma_t}^-(T) = 2k$. Let $A' = A \setminus \{v_3\}$, and so $|A'| = |A| - 1 = 3k + 2$. Let S be a non-isolating set of T such that $\gamma_t(T - S) < \gamma_t(T)$. We show that $|S| \geq 2k$. Suppose, to the contrary, that $|S| \leq 2k - 1$. Let D be a γ_t -set of $T - S$, and so $|D| = \gamma_t(T - S) \leq 3k + 2$. Since $|S| < 2k < \ell$ and each vertex in A' has ℓ leaf neighbors in T , we note that every vertex of A' is a support vertex of $T - S$, implying that $A' \subseteq D$, and so $3k + 2 \geq |D| \geq |A'| = 3k + 2$, implying that $D = A'$. In particular, $v_3 \notin D$, implying that all k leaf neighbors of v_3 belong to S ; that is, $L_3 \subseteq S$. If $v_{i_1} \notin S$ for some $i \in [k]$, then in order to totally dominate the vertex v_{i_1} , the vertex $v_{i_2} \in D$, contradicting our earlier observation that $D = A'$. Hence, $v_{i_1} \in S$ for all $i \in [k]$, and so $|S| \geq |L_3| + k = 2k$, a contradiction. Therefore, our original supposition that $|S| \leq 2k - 1$ is incorrect, implying that $|S| \geq 2k$ and $st_{\gamma_{pr}}^-(T) \geq 2k$. Conversely, if S^* consists of all $2k$ neighbors of v_3 different from v_2 in T , then S^* is a non-isolating set of T such that $\gamma_t(T - S^*) = |A'| < \gamma_t(T)$, implying that $st_{\gamma_t}^-(T) \leq |S^*| = 2k$. Consequently, $st_{\gamma_t}^-(T) = 2k$. Thus, $st_{\gamma_t}^-(T) - st_{\gamma_{pr}}^-(T) = k$. \square




Figure 2: A tree from the family T in the proof of Proposition 3.3.

Theorem 2.1 follows from Propositions 3.2 and 3.3. As further examples, we remark that if P is the Petersen graph, then $\gamma_t(P) = 4$ and $\gamma_{pr}(P) = 6$. Further, if v is an arbitrary vertex of P , then $\gamma_t(P - v) = 4$, and so $st_{\gamma_t}^-(P) \geq 2$. Moreover, if S consists of two non-adjacent vertices of P , then $\gamma_t(P - S) = 3$, and so $st_{\gamma_t}^-(P) \leq 2$. Consequently, $st_{\gamma_t}^-(P) = 2$. However if v is an arbitrary vertex of P , then $\gamma_{pr}(P - v) = 4$, implying that $st_{\gamma_{pr}}^-(P) = 1$. Moreover, let G_k be a graph obtained from the Petersen graph by replacing every vertex by a copy of a complete graph K_k for some $k \geq 1$, and adding all edges between two resulting complete graphs that correspond to two vertices of G_k (see Fig. 3). The resulting graph G_k is a $(4k - 1)$ -regular, $3k$ -connected graph that satisfies $\gamma_t(G_k) = 4$ and $st_{\gamma_t}^-(G_k) = 2k$, and $\gamma_{pr}(G_k) = 6$ and $st_{\gamma_{pr}}^-(G_k) = k$. This yields the following result.

Proposition 3.4. *For $k \geq 1$ an integer, there exists $(4k - 1)$ -regular, $3k$ -connected graphs G such that $st_{\gamma_t}^-(G) - st_{\gamma_{pr}}^-(G) = k$.*

4 Properties of paired domination stability

In this section, we present properties of paired domination stability in graphs. We begin with the following property of paired domination in graphs.




Figure 3: A graph G_k obtained from the Petersen graph by replacing every vertex by K_k .

Proposition 4.1. *Every connected isolate-free graph G contains a spanning tree T such that $\gamma_{pr}(T) = \gamma_{pr}(G)$.*

Proof. Since adding edges to a graph cannot increase its paired domination number, if T is an isolate-free spanning subgraph of a graph G , then $\gamma_{pr}(G) \leq \gamma_{pr}(T)$. Let D be a γ_{pr} -set of G , and so D is a PD-set of G and $|D| = \gamma_{pr}(G)$. Let M be a perfect matching in the subgraph $G[D]$ induced by D . Let T' be a spanning subgraph of G that consists of the edges in M and for each vertex v outside D , an edge of G that joins v to exactly one vertex of the dominating set D . If the resulting spanning subgraph T' is a tree, then we let $T = T'$. Otherwise, if the resulting spanning subgraph T' is a forest with $\ell \geq 2$ components, then we add $\ell - 1$ edges from the edge set of the graph G between these components, avoiding cycles, to construct a tree, which we call T . Since D is a PD-set in the resulting tree T , we note that $\gamma_{pr}(T) \leq |D| = \gamma_{pr}(G)$. Since T is an isolate-free spanning subgraph of G , we have $\gamma_{pr}(T) \geq \gamma_{pr}(G)$. Consequently, T is a spanning tree of G satisfying $\gamma_{pr}(T) = \gamma_{pr}(G)$. \square

By our earlier convention, if G is a graph of order n and $\gamma_{pr}(G) = 2$, then $st_{\gamma_{pr}}^-(G) = n$ since removing all vertices from the graph G produces the null graph with paired domination number zero. We are therefore only interested in the γ_{pr} -stability of graphs with paired domination number at least 4. If G is a graph with $\gamma_{pr}(G) \geq 4$ where x and y are adjacent vertices in G , then $D = V(G) \setminus \{x, y\}$ belongs to the set $NI(G)$ and $\gamma_{pr}(G - D) = \gamma_{pr}(K_2) = 2 < \gamma_{pr}(G)$. This yields the following result.

Observation 4.2. Every isolate-free graph G of order n with $\gamma_{pr}(G) \geq 4$ satisfies $st_{\gamma_{pr}}^-(G) \leq n - 2$.

Proposition 4.3. *If T is a spanning tree of a connected graph G such that $\gamma_{pr}(T) = \gamma_{pr}(G)$, then $st_{\gamma_{pr}}^-(T) \geq st_{\gamma_{pr}}^-(G)$.*

Proof. Let S be a $st_{\gamma_{pr}}^-$ -set of T . Thus, S is a set in $NI(T)$ with $|S| = st_{\gamma_{pr}}^-(T)$ such that $\gamma_{pr}(T - S) < \gamma_{pr}(T)$. Since $\gamma_{pr}(G - S) \leq \gamma_{pr}(T - S)$ and $\gamma_{pr}(T) = \gamma_{pr}(G)$, the set S is a non-isolating set of G such that $\gamma_{pr}(G - S) < \gamma_{pr}(G)$. Hence, $st_{\gamma_{pr}}^-(G) \leq |S| = st_{\gamma_{pr}}^-(T)$. \square

The following result shows that to determine the γ_{pr} -stability of a graph G , it is not sufficient to only examine spanning trees T of G satisfying $\gamma_{pr}(T) = \gamma_{pr}(G)$.

Proposition 4.4. *For $k \geq 1$ an integer, there exist connected graphs G such that $st_{\gamma_{pr}}^-(T) - st_{\gamma_{pr}}^-(G) = k$ for every spanning tree T of G with $\gamma_{pr}(T) = \gamma_{pr}(G)$.*

Proof. For $k \geq 1$, let F be obtained from two vertex disjoint copies of $K_{2,k+1}$ by identifying a vertex of degree $k + 1$ from each copy. Let u be the resulting identified vertex of degree $2(k + 1)$, and let w_1 and w_2 be the two vertices of degree $k + 1$ in F . Further, let v_i be a common neighbor (of degree 2) of u and w_i for $i \in [2]$. Let G be obtained from F by adding a leaf neighbor x_i to w_i for $i \in [2]$. Thus, $diam(G) = 6$ and $x_1w_1v_1w_2v_2x_2$ is a shortest path in G of length 6. The graph G satisfies $\gamma_{pr}(G) = 4$. We remark that only connected graphs of $diam(G) \leq 3$ have $\gamma_{pr}(G) = 2$. Therefore, $st_{\gamma_{pr}}^-(G) \geq 3$. Moreover, the set $S = \{w_1, x_1, x_2\}$ is a non-isolating set of minimum cardinality satisfying $\gamma_{pr}(G - S) = 2 < \gamma_{pr}(G)$, and so $st_{\gamma_{pr}}^-(G) = 3$. However, the vertex u must have degree 2 in every spanning tree T of G for which $\gamma_{pr}(T) = \gamma_{pr}(G) = 4$, implying that the vertices w_1 and w_2 each have $k + 1$ leaf neighbors in T . This implies that every non-isolating set of T that decreases the paired domination number contains at least $k + 3$ vertices. The set $S = N_T[w_1]$ is a non-isolating set of minimum cardinality satisfying $\gamma_{pr}(T - S) = 2 < \gamma_{pr}(T)$, and so $st_{\gamma_{pr}}^-(T) \leq |S| = k + 3$. Consequently, $st_{\gamma_{pr}}^-(T) = k + 3$, and so $st_{\gamma_{pr}}^-(T) - st_{\gamma_{pr}}^-(G) = k$. \square

Proposition 4.5. *If S is a $st_{\gamma_{pr}}^-$ -set of a connected isolate-free graph G with $\gamma_{pr}(G) \geq 4$, then $\gamma_{pr}(G - S) = \gamma_{pr}(G) - 2$.*

Proof. Let S be a $st_{\gamma_{pr}}^-$ -set of G . Suppose, to the contrary, that $\gamma_{pr}(G - S) \leq \gamma_{pr}(G) - 4$. By the connectivity of G , there exists a vertex $u \in S$ that has a neighbor in the set $V(G) \setminus S$. We now consider the set $S' = S \setminus \{u\}$. Let D be a γ_{pr} -set of $G - S$. If u has a neighbor in D , then D is a γ_{pr} -set of $G - S'$, implying that $\gamma_{pr}(G - S') \leq |D| = \gamma_{pr}(G - S) \leq \gamma_{pr}(G) - 4$, contradicting our choice of the set S . Hence, u has no neighbor in D . Let v be an arbitrary neighbor of u that belongs to $V(G) \setminus S$. The set $D \cup \{u, v\}$ is a PD-set of $G - S'$ with u and v paired, and with the pairings of the vertices of D unchanged from their pairings in $G - S$. Hence, $\gamma_{pr}(G - S') \leq |D| + 2 \leq \gamma_{pr}(G) - 2$, once again contradicting our choice of the set S . \square

5 Paths and cycles

It is well known (see, for example, [11]) that for $n \geq 3$ we have $\gamma_{pr}(C_n) = \gamma_{pr}(P_n) = 2\lceil \frac{n}{4} \rceil$. In this section, we determine the paired domination stability of paths and cycles. The proofs require a detailed case analysis, which is straightforward albeit tedious. We therefore omit the proofs in this section. The γ_{pr}^- -stability of a path P_n and a cycle C_n on n vertices is given by the following result.

Theorem 5.1. *If G is a path P_n , for $n \geq 2$, or a cycle C_n , for $n \geq 3$, then*

$$st_{\gamma_{pr}}^-(G) = \begin{cases} 1 & \text{when } n \equiv 1 \pmod{4} \\ 2 & \text{when } n \equiv 2 \pmod{4} \\ 3 & \text{when } n \equiv 3 \pmod{4} \\ 4 & \text{when } n \equiv 0 \pmod{4}. \end{cases}$$

Next we determine the γ_{pr}^+ -stability of a path P_n . For $n \leq 10$ with $n \neq 8$ and for $n = 13$, no non-isolating set of vertices in a path P_n exists whose removal increases the

paired domination number, and hence, by definition, $st_{\gamma_{pr}^+}(P_n) = \infty$ for such values of n . It is therefore only of interest to determine the γ_{pr}^+ -stability of a path P_n , where $n \geq 8$ and $n \notin \{9, 10, 13\}$.

Theorem 5.2. For $n \geq 8$ and $n \notin \{9, 10, 13\}$,

$$st_{\gamma_{pr}^+}(P_n) = \begin{cases} 1 & \text{when } n \pmod{4} \in \{0, 3\} \\ 2 & \text{when } n \pmod{4} \in \{1, 2\}. \end{cases}$$

As a consequence of Theorems 5.1 and 5.2, the paired domination stability of a path is determined.

Corollary 5.3. For $n \geq 2$,

$$st_{\gamma_{pr}^+}(P_n) = \begin{cases} 1 & \text{when } n \pmod{4} \in \{0, 1, 3\} \text{ and } n \notin \{3, 4, 7\} \\ 2 & \text{when } n \equiv 2 \pmod{4} \\ 3 & \text{when } n \in \{3, 7\} \\ 4 & \text{when } n = 4. \end{cases}$$

We next consider the γ_{pr}^+ -stability of a cycle C_n . As shown in Theorem 5.1, the γ_{pr}^- -stability of a path and a cycle of the same order are equal. This is not always the case for the γ_{pr}^+ -stability of a path and a cycle. For example, $st_{\gamma_{pr}^+}(P_{12}) = 1$ and $st_{\gamma_{pr}^+}(C_{12}) = 2$. Analogously as in the case of paths, for small values of the order of a cycle the γ_{pr}^+ -stability is infinite. Namely, for $n \leq 14$ with $n \neq 12$ and $n = 17$ we have that $st_{\gamma_{pr}^+}(C_n) = \infty$. The following result determines the γ_{pr}^+ -stability of a cycle of large order.

Theorem 5.4. For $n \geq 12$ and $n \notin \{13, 14, 17\}$,

$$st_{\gamma_{pr}^+}(C_n) = \begin{cases} 2 & \text{when } n \equiv 0 \pmod{4} \\ 3 & \text{when } n \pmod{4} \in \{2, 3\} \\ 4 & \text{when } n \equiv 1 \pmod{4}. \end{cases}$$

As a consequence of Theorems 5.1 and 5.4, the paired domination stability of a cycle is determined.

Corollary 5.5. For $n \geq 3$,

$$st_{\gamma_{pr}^+}(C_n) = \begin{cases} 1 & \text{when } n \equiv 1 \pmod{4} \\ 2 & \text{when } n \pmod{4} \in \{0, 2\} \text{ and } n \notin \{4, 8\} \\ 3 & \text{when } n \equiv 3 \pmod{4} \\ 4 & \text{when } n \in \{4, 8\}. \end{cases}$$

6 Trees

In this section, we first determine the γ_{pr} -stability of trees in the family \mathcal{F}_Δ and a new family \mathcal{E}_Δ .

Lemma 6.1. For $\Delta \geq 2$, if $T \in \mathcal{F}_\Delta$, then $st_{\gamma_{pr}^-}(T) = 2\Delta$.

Proof of Lemma 6.1. Let T be an arbitrary tree in the family $\mathcal{F}_{k,\Delta}$ for some $k \geq 2$ and $\Delta \geq 2$. We show that $\text{st}_{\gamma_{\text{pr}}}^-(T) = 2\Delta$. The family $\mathcal{F}_{k,2}$ consists of all paths P_{4k} where $k \geq 2$. Therefore by Theorem 5.1, we have $\text{st}_{\gamma_{\text{pr}}}^-(T) = 4 = 2\Delta$ for each $T \in \mathcal{F}_{k,2}$, which yields the desired result. Hence, we may assume that $\Delta \geq 3$. We show, by induction on $k \geq 2$, that every tree T in the family $\mathcal{F}_{k,\Delta}$ satisfies $\text{st}_{\gamma_{\text{pr}}}^-(T) = 2\Delta$.

Suppose $k = 2$, and so $T \in \mathcal{F}_{2,\Delta}$ (where recall that $\Delta \geq 3$). The tree T can therefore be constructed from two vertex disjoint double stars T_1 and T_2 , where $T_i \cong S(\Delta - 1, \Delta - 1)$ for $i \in [2]$, by selecting leaves w_1 and w_2 of T_1 and T_2 , respectively, and adding the edge w_1w_2 to $T_1 \cup T_2$. Let x_i and y_i be the two vertices of T_i that are not leaves, where x_iw_i is an edge. We note that $y_1x_1w_1w_2x_2y_2$ is a path in T . We note that $\gamma_{\text{pr}}(T) = 4$ and the set $\{x_1, x_2, y_1, y_2\}$ is a γ_{pr} -set of T .

Let S be a $\text{st}_{\gamma_{\text{pr}}}^-$ -set of G . Thus, S is a set in $\text{NI}(G)$ with $|S| = \text{st}_{\gamma_{\text{pr}}}^-(G)$ such that $\gamma_{\text{pr}}(T - S) = 2$. Let R be a γ_{pr} -set of $T - S$, and so R is a minimum PD-set of $T - S$ (of cardinality 2). Since $T[R] = P_2$, we note that $T - S$ is a tree of diameter at most 3. This implies that at most one of x_i and y_{3-i} belong to $T - S$ for $i \in [2]$. Thus, $|S \cap \{x_i, y_{3-i}\}| \geq 1$ for $i \in [2]$.

Suppose that $y_1 \in S$ and $x_2 \in S$. If $x_1 \in S$, then all leaf neighbors of y_1, x_1 and x_2 belong to S , while if $y_2 \in S$, then all leaf neighbors of y_1, y_2 and x_2 belong to S . In both cases, $|S| \geq 3\Delta - 2 > 2\Delta$.

Suppose that $y_1 \in S$ and $x_2 \notin S$. If $y_2 \in S$, then all leaf neighbors of y_1 and y_2 belong to S , implying that $|S| \geq 2\Delta$. If $y_2 \notin S$, then $x_1 \in S$, implying that S contains all leaf-neighbors of y_1 and x_1 , and so $|S| \geq 2\Delta - 1$. However if in this case $|S| = 2\Delta - 1$, implying that $\text{diam}(T - S) \geq 4$, a contradiction. Hence, $|S| \geq 2\Delta$.

Suppose that $y_1 \notin S$ and $x_2 \in S$. Since $T - S$ is a tree, $y_2 \in S$ and all leaf neighbors of y_2 and x_2 belong to S , implying that $|S| \geq 2\Delta - 1$. However if in this case $|S| = 2\Delta - 1$, then S contains x_2 and all leaf neighbors of y_1 , implying that $\text{diam}(T - S) \geq 4$, a contradiction. Hence, $|S| \geq 2\Delta$. Therefore, in all three cases we have $|S| \geq 2\Delta$, as desired. This proves the base case when $k = 2$.

For the inductive hypothesis, let $k \geq 3$ and assume that if $T' \in \mathcal{F}_{k',\Delta}$ where $2 \leq k' < k$, then $\text{st}_{\gamma_{\text{pr}}}^-(T') = 2\Delta$. We now consider a tree T in the family $\mathcal{F}_{k,\Delta}$. Therefore, the tree T can be constructed from k vertex disjoint double stars H_1, \dots, H_k , where $H_i \cong S(\Delta - 1, \Delta - 1)$ for $i \in [k]$, by selecting one leaf y_i from each double star H_i and adding $k - 1$ edges between vertices in $\{y_1, \dots, y_k\}$ in such a way that the resulting graph is a tree with maximum degree Δ . Let w_i and x_i be the two (adjacent) vertices of H_i that are not leaves for $i \in [k]$, where y_i is a leaf neighbor of x_i for $i \in [k]$. We note that $\gamma_{\text{pr}}(T) = 2k$ and the set $\cup_{i=1}^k \{w_i, x_i\}$ is the unique γ_{pr} -set of T .

Let U be the graph of order k whose vertices correspond to the k double stars H_1, \dots, H_k where two vertices are adjacent in U if and only if the corresponding double stars are joined by an edge in T . We call U the underlying graph of T . By construction, the graph U is a tree, noting that T is a tree. Let $V(U) = \{u_1, \dots, u_k\}$ where u_i is the vertex of U corresponding to the double star H_i for $i \in [k]$. Renaming the double stars if necessary, we may assume that u_1 is a leaf in U , and that H_1 is joined to H_2 in T . Thus, $y_1y_2 \in E(T)$ and $y_1y_j \notin E(T)$ for $j \in [k] \setminus [2]$. We note that $w_1x_1y_1y_2x_2w_2$ is a path in T . Let $T' = T - V(H_1)$. By construction, the tree T' belongs to the family $\mathcal{F}_{k',\Delta}$ where $k' = k - 1 \geq 2$. By induction, we have $\text{st}_{\gamma_{\text{pr}}}^-(T') = 2\Delta$.

Let S be a $\text{st}_{\gamma_{\text{pr}}}^-$ -set of T . Thus, S is a set in $\text{NI}(T)$ with $|S| = \text{st}_{\gamma_{\text{pr}}}^-(T)$ such that $\gamma_{\text{pr}}(T - S) \leq \gamma_{\text{pr}}(T) - 2 = 2k - 2$. Let Q be a γ_{pr} -set of $T - S$, and so $|Q| \leq 2k - 2$.

Let $Q' = Q \cap V(T')$ and $S' = S \cap V(T')$. For $i \in [k]$, let $Q_i = Q \cap V(H_i)$ and $S_i = S \cap V(H_i)$. We proceed further with the following claim.

Claim 6.2. $|S| \geq 2\Delta$.

Proof of Claim 6.2. Suppose, to the contrary, that $|S| \leq 2\Delta - 1$.

Subclaim 6.2.1. $|Q_1| \geq 2$.

Proof of Subclaim 6.2.1. Suppose, to the contrary, that $|Q_1| \leq 1$. Suppose that $Q_1 = \emptyset$. In this case, $V(H_1) \setminus \{y_1\} \subseteq S_1$. If $y_1 \in S_1$, then $|S_1| = 2\Delta > |S|$, a contradiction. Hence, $y_1 \notin S_1$, and so $2\Delta - 1 \geq |S| \geq |S_1| = 2\Delta - 1$, implying that $S = S_1$ and $|S| = 2\Delta - 1$. In this case, a γ_{pr} -set of $T - S$ contains at least one of y_1 and y_2 . Since the set $\cup_{i=2}^k \{w_i, x_i\}$ is the unique γ_{pr} -set of T' , a γ_{pr} -set of $T - S$ is therefore not a γ_{pr} -set of T' , and so $\gamma_{pr}(T - S) \geq \gamma_{pr}(T') + 2 = 2(k - 1) + 2 = 2k$, a contradiction. Hence, $|Q_1| \geq 1$.

By supposition, $|Q_1| \leq 1$. Consequently, $|Q_1| = 1$, implying that $Q_1 = \{y_1\}$ and $V(H_1) \setminus \{x_1, y_1\} \subseteq S_1$, and so $|S_1| \geq 2\Delta - 2$. If $x_1 \in S_1$, then $|S_1| = 2\Delta - 1$ and we end up in the previous case, which leads to a contradiction. Hence, $x_1 \notin S_1$ and $x_1 \notin Q_1$, implying that $y_2 \in Q$ with the vertices y_1 and y_2 paired in Q , and $|S_1| = 2\Delta - 2$. By supposition, $|S| \leq 2\Delta - 1$. If $|S| = 2\Delta - 2$, then $S = S_1$ and $\gamma_{pr}(T - S) \geq \gamma_{pr}(T') + 2 = 2k$, a contradiction. Hence, $|S| = 2\Delta - 1$, and so the set S contains a vertex $v' \in V(T') \setminus \{y_2\}$. However noting that $\Delta \geq 3$, every non-isolating set of vertices of $T' - y_2$ that decreases the paired domination number cannot contain only one vertex, implying that $\gamma_{pr}(T - S) \geq |\{y_1, y_2\}| + \gamma_{pr}(T' - y_2) = 2 + \gamma_{pr}(T') = 2k$, a contradiction. \square

Subclaim 6.2.2. $\{x_1, y_1\} \subseteq Q$.

Proof of Subclaim 6.2.2. Suppose, to the contrary, that $y_1 \notin Q_1$, implying that $S' \in \text{NI}(T')$. Recall that S is a $\text{st}_{\gamma_{pr}}^-$ -set of T and $|S'| \leq |S| \leq 2\Delta - 1$. However, $\text{st}_{\gamma_{pr}}^-(T') = 2\Delta$. Therefore, $\gamma_{pr}(T' - S') \geq \gamma_{pr}(T') = 2(k - 1)$. Hence, $\gamma_{pr}(T - S) = \gamma_{pr}(T' - S') + |Q_1| \geq 2(k - 1) + 2 = 2k$, a contradiction. Hence, $y_1 \in Q_1$.

Suppose, to the contrary, that $x_1 \notin Q_1$. Thus, all $\Delta - 2$ leaf-neighbors of x_1 belong to the set S_1 . By Claim 6.2.1, we have $|Q_1| \geq 2$. Hence, the set Q_1 contains w_1 and one of its leaf-neighbor w'_1 . We now consider the set $S^* = S \setminus S_1$. Since $S^* \in \text{NI}(T)$ and $(Q \setminus \{w'_1\}) \cup \{x_1\}$ is a PD-set of $T - S^*$, we have $\gamma_{pr}(T - S^*) \leq |Q| = \gamma_{pr}(T - S)$, contradicting our choice of the set S . Hence, $x_1 \in Q_1$. \square

Subclaim 6.2.3. $w_1 \notin Q_1$.

Proof of Subclaim 6.2.3. Suppose, to the contrary, that $w_1 \in Q_1$. Hence, $\{w_1, x_1, y_1\} \subseteq Q_1$, and so $S \cap V(H_1) = \emptyset$ by the minimality of S . Thus, $S = S'$ and therefore $|S'| \leq 2\Delta - 1$.

We show firstly that x_1 and y_1 are paired in Q . Suppose, to the contrary, that x_1 and y_1 are not paired in Q . This implies that $y_2 \in Q$, and that y_1 and y_2 are paired in Q . Suppose that $x_2 \notin S$, implying that $S' \in \text{NI}(T')$. By the minimality of the set Q , we have $x_2 \notin Q$. Thus, the set $Q' \cup \{x_2\}$ is a PD-set of $T' - S'$, and so $|Q'| + 1 = |Q' \cup \{x_2\}| \geq \gamma_{pr}(T' - S') \geq \gamma_{pr}(T') = 2(k - 1)$. Hence, $|Q| = |Q_1| + |Q'| \geq 3 + (2k - 3) = 2k = \gamma_{pr}(T)$, a contradiction. Hence, $x_2 \in S$. We now consider the set $S^* = S \setminus \{x_2\}$. We note that S^* is a non-isolating set of vertices of T , and the set Q is a PD-set of $T - S^*$. Thus,

$\gamma_{\text{pr}}(T - S^*) \leq |Q| \leq 2k - 2$, which contradicts our choice of the set S . Hence, x_1 and y_1 are paired in Q .

Since x_1 and y_1 are paired in Q , the vertex w_1 is paired with one of its leaf neighbors, say w'_1 . By the minimality of Q we note that $Q_1 = \{w_1, w'_1, x_1, y_1\}$. If $x_2 \in Q$, then the set $Q \setminus \{w'_1, y_1\}$ is a PD-set of $T - S$ (with w_1 and x_1 paired), contradicting the minimality of Q . Hence, $x_2 \notin Q$. This in turn implies that $y_2 \notin Q$. If $y_2 \in S$, then once again we contradict the minimality of Q . Therefore, $y_2 \notin S$. We remark, though, that possibly $x_2 \in S$. Recall that by our earlier observations, $S = S'$.

Let $S'' = S \setminus \{x_2\}$. Thus, if $x_2 \notin S$, then $S'' = S$, while if $x_2 \in S$, then $S'' = S \setminus \{x_2\}$. The set S'' is a non-isolating set of T' such that $|S''| \leq |S| \leq 2\Delta - 1$. As observed earlier, $y_2 \notin Q'$ and $x_2 \notin Q'$. The set $Q' \cup \{y_2, x_2\}$ is a PD-set of $T' - S''$, implying that $|Q'| + 2 \geq \gamma_{\text{pr}}(T' - S'') \geq \gamma_{\text{pr}}(T') = 2(k - 1)$. Hence, $|Q'| \geq 2k - 4$, and so $|Q| = |Q_1| + |Q'| \geq 4 + (2k - 4) = 2k$, contradicting the fact that $|Q| \leq 2k - 2$. \square

Proof of Claim 6.2, continued: By Claim 6.2.3, $w_1 \notin Q_1$. This implies that $Q_1 = \{x_1, y_1\}$. The set S_1 therefore consists of the $\Delta - 1$ leaf neighbors of w_1 , and so $|S_1| = \Delta - 1$. This is true for every leaf in the tree U . Hence, if u_i is a leaf in U for some $i \in [k]$, then in the corresponding double star H_i of T we have $Q_i = \{x_i, y_i\}$ and $|S_i| = \Delta - 1$. Further, the set S_i consists of the $\Delta - 1$ leaf neighbors of w_i . In particular, $|Q_1| = 2$ and $|S_1| = \Delta - 1$. Since the underlying tree U of T has order $k \geq 3$, there are at least two leaves in U . Thus, u_p is a leaf in U for some $p \in [k] \setminus \{1\}$, implying that $|Q_p| = 2$ and $|S_p| = \Delta - 1$.

If $|Q_i| \geq 2$ for all $i \in [k]$, then $|Q| \geq 2k$, a contradiction. Hence, $|Q_q| \leq 1$ for some $q \in [k]$. By our earlier observations, u_q is not a leaf in the tree U , and so $q \notin \{1, p\}$. If $|Q_q| = 0$, then $\{w_q, x_q\} \subseteq S_q$, and so $|S_q| \geq 2$ (in fact, $|S_q| \geq 2\Delta - 1$) and $|S| \geq |S_1| + |S_p| + |S_q| \geq (\Delta - 1) + (\Delta - 1) + 2 = 2\Delta$, a contradiction. Hence, $|Q_q| = 1$, implying that $Q_q = \{y_q\}$ and $w_q \in S_q$, and so $|S_q| \geq 1$. Since the paired dominating number is an even integer and $|Q| \leq 2k$, there exists $r \in [k] \setminus \{1, p, q\}$ such that $|Q_r| = 1$. Therefore, $Q_r = \{y_r\}$ and $|S_r| \geq 1$. Hence, $|S| \geq |S_1| + |S_p| + |S_q| + |S_r| \geq (\Delta - 1) + (\Delta - 1) + 1 + 1 = 2\Delta$, a contradiction. This completes the proof of Claim 6.2. \square

Proof of Lemma 6.1, continued: By Claim 6.2, we have $|S| \geq 2\Delta$. By our choice of the set S , this implies that $\text{st}_{\gamma_{\text{pr}}}^-(T) = |S| \geq 2\Delta$. Conversely, if we consider the set $S = V(H_1)$, then $S \in \text{NI}(T)$ satisfies $|S| = 2\Delta$ and $\gamma_{\text{pr}}(T - S) = \gamma_{\text{pr}}(T') = 2k - 2 < \gamma_{\text{pr}}(T)$, and so $\text{st}_{\gamma_{\text{pr}}}^-(T) \leq 2\Delta$. Consequently, $\text{st}_{\gamma_{\text{pr}}}^-(T) = 2\Delta$. This completes the proof of Lemma 6.1. \square

We determine next the γ_{pr}^+ -stability of a tree in the family \mathcal{F}_Δ .

Lemma 6.3. *For $\Delta \geq 2$, if $T \in \mathcal{F}_\Delta$, then $\text{st}_{\gamma_{\text{pr}}}^+(T) \leq \Delta - 1$.*

Proof. Let T be an arbitrary tree in the family $\mathcal{F}_{k,\Delta}$ for some $k \geq 2$ and $\Delta \geq 2$. We use the same notation as in the proof of Lemma 6.1. In particular, $\gamma_{\text{pr}}(T) = 2k$ and H_1 corresponds to a leaf u_1 in the underlying tree U of T . Moreover, y_1y_2 is the edge joining H_1 and H_2 in T . Also, w_i and x_i are the support vertices in the double star H_i and $w_ix_iy_i$ is a path in H_i for $i \in [k]$. Let L be the set of $\Delta - 2$ leaf neighbors of x_1 in T , and let $S = L \cup \{x_1\}$. We resulting set $S \in \text{NI}(T)$ and the forest $T - S$ has two components, say F_1 and F_2 where $w_1 \in V(F_1)$ and $y_1 \in V(F_2)$. Moreover, $\gamma_{\text{pr}}(T - S) = \gamma_{\text{pr}}(F_1) + \gamma_{\text{pr}}(F_2) = 2 + 2k > \gamma_{\text{pr}}(T)$. Therefore, $\text{st}_{\gamma_{\text{pr}}}^+(T) \leq |S| = \Delta - 1$. \square

Recall that by Proposition 3.1, for $\Delta \geq 3$, if $T \in \mathcal{H}_\Delta$, then $\text{st}_{\gamma_{\text{pr}}}(T) = 2\Delta - 1$. Further we remark that $\text{st}_{\gamma_{\text{pr}}}^+(T) = \infty$. We next define another family of trees T with maximum degree Δ such that $\text{st}_{\gamma_{\text{pr}}}^-(T) = 2\Delta - 1$. For integers $\Delta \geq 3$ and $\Delta - 1 \geq k \geq 3$, let $E_{k,\Delta}$ be a graph obtained from the path P_2 with vertices u and v and the disjoint union of $2k$ double stars $S(\Delta - 1, \Delta - 1)$ by selecting one leaf from each double star and identifying half of the selected leaves with the vertex v and the other half of the selected leaves with the vertex u (see Figure 4). Let

$$\mathcal{E}_\Delta = \bigcup_{k \geq 3} E_{k,\Delta}.$$




Figure 4: A tree $E_{k,5}$ from the family \mathcal{E}_5 .

If T is a tree from the family \mathcal{E}_Δ , then $\text{st}_{\gamma_{\text{pr}}}^-(T) = 2\Delta - 1$. Moreover, if T is isomorphic to the graph $E_{k,\Delta}$, then $\text{st}_{\gamma_{\text{pr}}}^+(T) = k(\Delta - 1)$. In contrast to the family \mathcal{H}_Δ , the trees from the family \mathcal{E}_Δ have finite γ_{pr}^+ -stability.

7 Proof of Theorem 2.2

In this section we present a proof of Theorem 2.2, which we restate below.

Theorem 2.2. *If T is a tree with maximum degree Δ satisfying $\gamma_{\text{pr}}(T) \geq 4$, then the following hold.*

- (a) $\text{st}_{\gamma_{\text{pr}}}^-(T) \leq 2\Delta$, with equality if and only if $T \in \mathcal{F}_\Delta$.
- (b) $\text{st}_{\gamma_{\text{pr}}}(T) \leq 2\Delta - 1$, and this bound is sharp for all $\Delta \geq 2$.

Proof. We first prove the statement given in part (a). Since $\gamma_{\text{pr}}(T) \geq 4$, we have $\Delta \geq 2$. If $\Delta = 2$, then G is a path P_n of order $n \geq 5$. In this case, the family $\mathcal{F}_{k,\Delta} = \{P_n : n \equiv 0 \pmod{4} \text{ and } n \geq 8\}$, and Theorem 5.1 and Lemma 6.1 imply the desired result. Suppose, therefore, that $\Delta \geq 3$. The sufficiency of part (a) follows from Lemma 6.1. To prove the necessity, let T be a tree with maximum degree $\Delta \geq 3$ satisfying $\gamma_{\text{pr}}(T) \geq 4$. Let $d = \text{diam}(T)$, and so $d \geq 4$. Let $P : v_0 v_1 \dots v_d$ be a diametral path in G . Thus, v_0 and v_d are leaves in T and $d(v_0, v_d) = \text{diam}(G)$. We now consider the tree T rooted at the vertex v_d . Let D be a γ_{pr} -set of T .

Suppose that there is a child u_1 of v_2 that is a support vertex in T where $u_1 \neq v_1$. Let u_0 be a leaf neighbor of u_1 . Since every PD-set of T contains all support vertices, we have $\{v_1, u_1\} \subset D$. Renaming vertices if necessary, we may assume that u_0 and u_1 are paired in D . Thus, if S consists of the vertex u_1 and all leaf neighbors of u_1 , then $S \in \text{NI}(T)$ and $\gamma_{\text{pr}}(T - S) \leq |D| - 2 = \gamma_{\text{pr}}(T) - 2$. Hence, $\text{st}_{\gamma_{\text{pr}}}^-(T) \leq |S| \leq \Delta < 2\Delta - 1$, and the desired result follows. Assume, therefore, that every child of v_2 different from v_1 is a leaf.

Suppose that there is a γ_{pr} -set, $D_{2,3}$, of T such that v_2 and v_3 are paired in $D_{2,3}$. Necessarily, $v_1 \in D_{2,3}$ and v_1 is paired in $D_{2,3}$ with one of its leaf neighbors. Let S consist of the vertex v_1 and all of its leaf neighbors. Thus, $S \in \text{NI}(T)$ and $\gamma_{\text{pr}}(T - S) \leq |D_{2,3}| - 2 = \gamma_{\text{pr}}(T) - 2 < \gamma_{\text{pr}}(T)$, implying that $\text{st}_{\gamma_{\text{pr}}}^-(T) \leq |S| \leq \Delta < 2\Delta - 1$, once again implying the desired result. Therefore, we may assume that in every γ_{pr} -set of T the vertices v_2 and v_3 are not paired.

Suppose that there is a γ_{pr} -set, D_3 , of T which contains a neighbor of v_3 different from v_2 . In this case, if S consists of the vertex v_2 and all its descendants, then $|S| \leq 2\Delta - 1$, $S \in \text{NI}(T)$ and $\gamma_{\text{pr}}(T - S) \leq |D_3| - 2 = \gamma_{\text{pr}}(T) - 2 < \gamma_{\text{pr}}(T)$, noting that the set $D_3 \setminus S$ is a PD-set of $T - S$ and, by the minimality of D_3 we have $|D_3 \cap S| = 2$. Thus, $\text{st}_{\gamma_{\text{pr}}}^-(T) \leq |S| \leq 2\Delta - 1$, and the desired result follows. Hence, we may assume that every γ_{pr} -set of T contains the vertex v_2 but no other vertex in $N[v_3]$. In particular, $N[v_3] \cap D = \{v_2\}$.

Suppose that $d_T(v_1) < \Delta$ or $d_T(v_2) < \Delta$. Thus, $d_T(v_1) + d_T(v_2) \leq 2\Delta - 1$. In order to dominate the vertex v_0 , we have $v_1 \in D$. By our earlier assumptions, $v_2 \in D$ and every child of v_2 different from v_1 is a leaf. Thus by the minimality of the set D , the vertex v_1 is the only descendant of v_2 that belongs to the set D , and the vertices v_1 and v_2 are paired in D . Hence, if $S = N[v_2] \cup N[v_1]$, then $S \in \text{NI}(T)$ and $|S| = d_T(v_1) + d_T(v_2) \leq 2\Delta - 1$. Further, $D \setminus \{v_1, v_2\}$ is a PD-set of $T - S$, and so $\gamma_{\text{pr}}(T - S) \leq |D| - 2 = \gamma_{\text{pr}}(T) - 2$, implying that $\text{st}_{\gamma_{\text{pr}}}^-(T) \leq |S| \leq 2\Delta - 1$, yielding the desired result. Hence, we may assume that $d_T(v_1) = d_T(v_2) = \Delta$.

Suppose that $d_T(v_3) \geq 3$, and let u_2 be a child of v_3 different from v_2 . If u_2 is a leaf, then v_3 belongs to every γ_{pr} -set of T , while if u_2 is not a leaf, then from the structure of the rooted tree T the vertex u_2 can be chosen to belong to some γ_{pr} -set of T . In both cases, we contradict our earlier assumption that every γ_{pr} -set of T contains the vertex v_2 but no other vertex in $N[v_3]$. Hence, $d_T(v_3) = 2$. We now let $S = N[v_1] \cup N[v_2]$, and so $S \in \text{NI}(T)$ and $|S| = d_T(v_1) + d_T(v_2)$. By our earlier observations, $|S| = 2\Delta$ and $\gamma_{\text{pr}}(T - S) = \gamma_{\text{pr}}(T) - 2 < \gamma_{\text{pr}}(T)$, implying that $\text{st}_{\gamma_{\text{pr}}}^-(T) \leq |S| = 2\Delta$. This proves the desired upper bound.

We show next that if we have equality in the upper bound in part (a), then $T \in \mathcal{F}_\Delta$. Let $\text{st}_{\gamma_{\text{pr}}}^-(T) = 2\Delta$. By our earlier observations, we have that every child of v_2 different from v_1 is a leaf. Further, $d_T(v_1) = d_T(v_2) = \Delta$ and $d_T(v_3) = 2$. We now re-root the tree T at the vertex v_0 , thereby interchanging the roles of v_0 and v_d . Identical arguments as before show that every child of v_{d-2} different from v_{d-1} is a leaf. Further, $d_T(v_{d-1}) = d_T(v_{d-2}) = \Delta$ and $d_T(v_{d-3}) = 2$. In particular, $d \geq 6$.

Suppose that $d = 6$, and so $v_{d-3} = v_3$. In this case, the tree T is determined and $\gamma_{\text{pr}}(T) = 4$. Letting $S = (N[v_1] \cup N[v_2]) \setminus \{v_3\}$, we have $S \in \text{NI}(T)$ and $|S| = d_T(v_1) + d_T(v_2) - 1 = 2\Delta - 1$. Further, $\gamma_{\text{pr}}(T - S) = 2 < \gamma_{\text{pr}}(T)$. Therefore, $\text{st}_{\gamma_{\text{pr}}}^-(T) \leq |S| = 2\Delta - 1$, a contradiction. Hence, $d \geq 7$, and so $v_{d-3} \neq v_3$.

We now consider the tree $T' = T - (N[v_1] \cup N[v_2])$. If $\gamma_{\text{pr}}(T') = 2$, then by our earlier observations, we have $d = 7$ and $T' \cong S(\Delta - 1, \Delta - 1)$ where v_{d-1} and v_{d-2} are the two (adjacent) vertices in T' that are not leaves. Therefore, $T \in T_{2,\Delta}$, and so $T \in T_\Delta$. Hence, we may assume that $\gamma_{\text{pr}}(T') \geq 4$, for otherwise the desired characterization follows. In particular, $d \geq 8$. As observed earlier, $d_T(v_{d-1}) = d_T(v_{d-2}) = \Delta$, implying that $\Delta(T') = \Delta$ and $\text{st}_{\gamma_{\text{pr}}}^-(T') \leq 2\Delta$.

Let D be a γ_{pr} -set of T . Since every PD-set of T contains the set of support vertices, we note that $v_1, v_2 \in D$. By the minimality of D , no leaf-neighbor of v_1 or v_2 belongs to

D. If $v_3 \in D$, then $v_4 \in D$ (with v_3 and v_4 paired in D). However in this case, we can replace v_3 in D with an arbitrary neighbor of v_4 that does not belong to D . Hence, we can choose the γ_{pr} -set D of T so that $v_3 \notin D$. The resulting set D when restricted to $V(T')$ is a PD-set of T' , implying that $\gamma_{\text{pr}}(T') \leq |D| - 2 = \gamma_{\text{pr}}(T) - 2$. Conversely, every PD-set of T' can be extended to a PD-set of T by adding to it the vertices v_1 and v_2 (with v_1 and v_2 paired), and so $\gamma_{\text{pr}}(T) \leq \gamma_{\text{pr}}(T') + 2$. Consequently, $\gamma_{\text{pr}}(T) = \gamma_{\text{pr}}(T') + 2$.

Suppose that $\text{st}_{\gamma_{\text{pr}}}^-(T') < 2\Delta$. Let S' be a $\text{st}_{\gamma_{\text{pr}}}^-$ -set of T' . Thus, S is a set in $\text{NI}(T')$ with $|S'| = \text{st}_{\gamma_{\text{pr}}}^-(T') < 2\Delta$ such that $\gamma_{\text{pr}}(T - S') < \gamma_{\text{pr}}(T')$. If D' is a γ_{pr} -set of $T' - S'$, then $D' \cup \{v_1, v_2\}$ is a PD-set of $T - S$, and so $\gamma_{\text{pr}}(T - S') \leq |D'| + 2 = \gamma_{\text{pr}}(T - S') + 2 < \gamma_{\text{pr}}(T') + 2 = \gamma_{\text{pr}}(T)$. Hence, $S' \in \text{NI}(T)$ and $\gamma_{\text{pr}}(T - S') < \gamma_{\text{pr}}(T')$, implying that $\text{st}_{\gamma_{\text{pr}}}^-(T) \leq |S'| = \text{st}_{\gamma_{\text{pr}}}^-(T') < 2\Delta$, a contradiction. Therefore, $\text{st}_{\gamma_{\text{pr}}}^-(T') = 2\Delta$.

Hence, the tree T' satisfies $\Delta(T') = \Delta$, $\gamma_{\text{pr}}(T') \geq 4$ and $\text{st}_{\gamma_{\text{pr}}}^-(T') = 2\Delta$. Proceeding by induction, we have $T' \in \mathcal{F}_\Delta$. Thus, T' is constructed from the disjoint union of k' double stars each isomorphic to $S(\Delta - 1, \Delta - 1)$, by selecting one leaf from each double star and adding $k' - 1$ edges between these selected leaves to produce a tree with maximum degree Δ . The resulting tree T' satisfies $\gamma_{\text{pr}}(T') = 2k'$ with the $2k'$ support vertices forming a γ_{pr} -set of T' .

By construction of T' , the tree T' contains the vertex v_4 but not the vertex v_3 . Suppose that v_4 is a support vertex in T' , implying by construction of T' that v_4 is a vertex of degree Δ in T' . Let $S = (N[v_1] \cup N[v_2]) \setminus \{v_3\}$. We note that $S \in \text{NI}(T)$ and $|S| = 2\Delta - 1$. Let D' be the (unique) γ_{pr} -set of T' , and so D' is the set of $2k'$ support vertices in T' . In particular, we note that $v_4 \in D'$. The set D' is a PD-set of $T - S$, and so $\gamma_{\text{pr}}(T - S) \leq |D'| = \gamma_{\text{pr}}(T') = \gamma_{\text{pr}}(T) - 2$. Therefore, $\text{st}_{\gamma_{\text{pr}}}^-(T) \leq |S| = 2\Delta - 1$, a contradiction. Hence, v_4 is a leaf of T' , and so v_4 is a leaf in one of the k' double stars in the construction of T' . Selecting the leaf v_4 from this double star and selecting the leaf v_3 from the double star induced by $N[v_1] \cup N[v_2]$, which is isomorphic to $S(\Delta - 1, \Delta - 1)$, and adding back the edge v_3v_4 we re-construct the tree T , showing that $T \in \mathcal{F}_\Delta$. This completes the proof of part (a).

Part (b) now follows readily from part (a). If $T \in \mathcal{F}_\Delta$ for some $\Delta \geq 2$, then by Lemmas 6.1 and 6.3, we have $\text{st}_{\gamma_{\text{pr}}}(T) \leq \Delta - 1$. Hence, we may assume that $T \notin \mathcal{F}_\Delta$ for any $\Delta \geq 2$, for otherwise the bound in part (b) is immediate. With this assumption, the upper bound in part (b) follows immediately from part (a) noting that $\text{st}_{\gamma_{\text{pr}}}^-(T) \leq \text{st}_{\gamma_{\text{pr}}}(T) \leq 2\Delta - 1$. That the bound is tight for all $\Delta \geq 2$ follows from Proposition 3.1. \square

8 Proof of Theorem 2.3

In this section we present a proof of Theorem 2.3, which we restate below.

Theorem 2.3. *If G is a connected graph with $\gamma_{\text{pr}}(G) \geq 4$, then $\text{st}_{\gamma_{\text{pr}}}^-(G) \leq 2\Delta(G)$, and this bound is sharp.*

Proof. Let G be a connected graph with $\gamma_{\text{pr}}(G) \geq 4$ and let $\Delta = \Delta(G)$. Since $\gamma_{\text{pr}}(G) \geq 4$, we have $\Delta \geq 2$. If $\Delta = 2$, then G is a path P_n or a cycle C_n , and by Theorem 5.1, we have $\text{st}_{\gamma_{\text{pr}}}^-(G) \leq 2\Delta$, with equality if and only if $n \equiv 0 \pmod{4}$. Assume, therefore, that $\Delta \geq 3$.

Let T be a spanning tree of G such that $\gamma_{\text{pr}}(T) = \gamma_{\text{pr}}(G)$. We note that such a tree exists by Lemma 4.1. Let S be a $\text{st}_{\gamma_{\text{pr}}}^-$ -set of T . Thus, S is a set in $\text{NI}(T)$ with $|S| = \text{st}_{\gamma_{\text{pr}}}^-(T)$ such that $\gamma_{\text{pr}}(T - S) < \gamma_{\text{pr}}(T)$. By Observation 4.2, we have

$|S| = \text{st}_{\gamma_{\text{pr}}}^-(T) \leq n - 2$. Since $S \in \text{NI}(T)$, every vertex in $T - S$, and therefore in the supergraph $G - S$, has degree at least 1. Hence, $S \in \text{NI}(G)$ and since $\gamma_{\text{pr}}(G - S) \leq \gamma_{\text{pr}}(T - S)$, we have $\gamma_{\text{pr}}(G - S) < \gamma_{\text{pr}}(G)$. Thus, $\text{st}_{\gamma_{\text{pr}}}^-(G) \leq |S| = \text{st}_{\gamma_{\text{pr}}}^-(T)$. By Theorem 2.2, we have $\text{st}_{\gamma_{\text{pr}}}^-(T) \leq 2\Delta(T)$. Noting that $\Delta(T) \leq \Delta(G)$, we therefore have that $\text{st}_{\gamma_{\text{pr}}}^-(G) \leq \text{st}_{\gamma_{\text{pr}}}^-(T) \leq 2\Delta(T) \leq 2\Delta(G) = 2\Delta$.

To show that the upper bound in Theorem 2.3 is tight, we present a family of graphs with maximum degree Δ and $\gamma_{\text{pr}}(G) \geq 4$ satisfying $\text{st}_{\gamma_{\text{pr}}}^-(G) = 2\Delta$. Our first family, \mathcal{G}_Δ , is constructed as follows. For $k \geq 2$ and $\Delta \geq 2$, let $G_{k,\Delta}$ be a graph obtained from k double stars $S(\Delta - 1, \Delta - 1)$ by choosing two leaves at distance 3 apart in each double star and adding k edges between the chosen leaves in such a way, that every chosen vertex has degree 2 in the resulting graph. Let \mathcal{G}_Δ be the family of all such graphs $G_{k,\Delta}$ for all $k \geq 2$. The graph $G_{2,6} \in \mathcal{G}_6$, for example, is illustrated in Figure 5. We note that $\gamma_{\text{pr}}(G_{k,\Delta}) = 2k$ and that set of $2k$ vertices of degree Δ is the unique γ_{pr} -set of $G_{k,\Delta}$. Furthermore, $\text{st}_{\gamma_{\text{pr}}}^-(G_{k,\Delta}) = 2\Delta$. \square




Figure 5: The graph $G_{2,6}$ from a class of graphs $G_{k,\Delta}$.

Recall that by definition we have $\text{st}_{\gamma_{\text{pr}}}(G) \leq \text{st}_{\gamma_{\text{pr}}}^-(G)$ for every graph G . Hence, as an immediate consequence of Theorem 2.3 we have Corollary 2.4. Recall its statement.


Corollary 2.4. *If G is a connected graph with $\gamma_{\text{pr}}(G) \geq 4$, then $\text{st}_{\gamma_{\text{pr}}}(G) \leq 2\Delta(G)$.*


It remains an open problem, however, to determine if the upper bound of Corollary 2.4 is best achievable for all values of possible value of $\Delta(G) = \Delta \geq 2$. If $\Delta = 2$ and G is a path, then $G \cong P_n$ where $n \geq 5$, and $\text{st}_{\gamma_{\text{pr}}}(G) \leq 2\Delta - 2$ by Corollary 5.3. If $\Delta = 2$ and G is a cycle, then $G \cong C_n$ where $n \geq 5$, and $\text{st}_{\gamma_{\text{pr}}}(G) \leq 2\Delta$ by Corollary 5.5, with equality if and only if $G = C_8$. Hence, the only connected graph G with maximum degree $\Delta = 2$ satisfying $\gamma_{\text{pr}}(G) \geq 4$ and $\text{st}_{\gamma_{\text{pr}}}(G) = 2\Delta$ is the 8-cycle, namely $G = C_8$. For $\Delta \geq 3$, we do not know of a connected graph G with maximum degree Δ satisfying $\gamma_{\text{pr}}(G) \geq 4$ and $\text{st}_{\gamma_{\text{pr}}}(G) = 2\Delta$.

By Corollary 5.5 and Proposition 3.1, for any given $\Delta \geq 2$, there do exists infinite families of connected graphs G with maximum degree Δ satisfying $\text{st}_{\gamma_{\text{pr}}}(G) = 2\Delta - 1$. Thus, if the upper bound of Corollary 2.4 can be improved to $\text{st}_{\gamma_{\text{pr}}}(G) \leq 2\Delta - 1$ in the case when $\Delta \geq 3$, then this bound would be tight.

ORCID iDs

Aleksandra Gorzkowska  <https://orcid.org/0000-0001-5335-7351>

Michael A. Henning  <https://orcid.org/0000-0001-8185-067X>

Monika Piłśniak  <https://orcid.org/0000-0002-3734-7230>

Elżbieta Tumidajewicz  <https://orcid.org/0000-0002-1413-2413>

References

- [1] M. Amraee, N. Jafari Rad and M. Maghasedi, Roman domination stability in graphs, *Math. Rep. (Bucur.)* **21** (2019), 193–204, http://imar.ro/journals/Mathematical_Reports/php/2019/Mrc19_2.php.
- [2] A. Aytaç and B. Atay Atakul, Exponential domination critical and stability in some graphs, *Int. J. Found. Comput. Sci.* **30** (2019), 781–791, doi:10.1142/s0129054119500217.
- [3] D. Bauer, F. Harary, J. Nieminen and C. L. Suffel, Domination alteration sets in graphs, *Discrete Math.* **47** (1983), 153–161, doi:10.1016/0012-365x(83)90085-7.
- [4] R. C. Brigham, P. Z. Chinn and R. D. Dutton, Vertex domination-critical graphs, *Networks* **18** (1988), 173–179, doi:10.1002/net.3230180304.
- [5] R. C. Brigham, T. W. Haynes, M. A. Henning and D. F. Rall, Bicritical domination, *Discrete Math.* **305** (2005), 18–32, doi:10.1016/j.disc.2005.09.013.
- [6] T. Burton and D. P. Sumner, Domination dot-critical graphs, *Discrete Math.* **306** (2006), 11–18, doi:10.1016/j.disc.2005.06.029.
- [7] W. J. Desormeaux, T. W. Haynes and M. A. Henning, Total domination changing and stable graphs upon vertex removal, *Discrete Appl. Math.* **159** (2011), 1548–1554, doi:10.1016/j.dam.2011.06.006.
- [8] W. J. Desormeaux, T. W. Haynes and M. A. Henning, Paired domination in graphs, in: T. W. Haynes, S. T. Hedetniemi and M. A. Henning (eds.), *Topics in Domination in Graphs*, Springer, Cham, volume 64 of *Dev. Math.*, pp. 31–77, 2020, doi:10.1007/978-3-030-51117-3_3.
- [9] O. Favaron, D. P. Sumner and E. Wojcicka, The diameter of domination k -critical graphs, *J. Graph Theory* **18** (1994), 723–734, doi:10.1002/jgt.3190180708.
- [10] T. W. Haynes, S. T. Hedetniemi and M. A. Henning (eds.), *Topics in Domination in Graphs*, volume 64 of *Developments in Mathematics*, Springer, Cham, 2020, doi:10.1007/978-3-030-51117-3.
- [11] T. W. Haynes and P. J. Slater, Paired-domination in graphs, *Networks* **32** (1998), 199–206, doi:10.1002/(sici)1097-0037(199810)32:3<199::aid-net4>3.0.co;2-f.
- [12] M. A. Henning and M. Krzykowski, Total domination stability in graphs, *Discrete Appl. Math.* **236** (2018), 246–255, doi:10.1016/j.dam.2017.07.022.
- [13] M. A. Henning and N. J. Rad, On total domination vertex critical graphs of high connectivity, *Discrete Appl. Math.* **157** (2009), 1969–1973, doi:10.1016/j.dam.2008.12.009.
- [14] M. A. Henning and A. Yeo, *Total Domination in Graphs*, Springer Monographs in Mathematics, Springer, New York, 2013, doi:10.1007/978-1-4614-6525-6.
- [15] N. Jafari Rad, E. Sharifi and M. Krzykowski, Domination stability in graphs, *Discrete Math.* **339** (2016), 1909–1914, doi:10.1016/j.disc.2015.12.026.
- [16] Z. Li, Z. Shao and S.-j. Xu, 2-rainbow domination stability of graphs, *J. Comb. Optim.* **38** (2019), 836–845, doi:10.1007/s10878-019-00414-0.
- [17] D. P. Sumner, Critical concepts in domination, *Discrete Math.* **86** (1990), 33–46, doi:10.1016/0012-365x(90)90347-k.
- [18] D. P. Sumner and P. Blitch, Domination critical graphs, *J. Comb. Theory Ser. B* **34** (1983), 65–76, doi:10.1016/0095-8956(83)90007-2.