

Dominating sets in triangulations on surfaces

Hong Liu

*Department of Mathematics, University of Illinois Urbana-Champaign, Urbana, Illinois
61801, USA*

Michael J. Pelsmajer *

*Department of Applied Mathematics, Illinois Institute of Technology, Chicago, Illinois
60616, USA*

Received 13 June 2010, accepted 27 January 2011, published online 30 March 2011

Abstract

A dominating set $D \subseteq V(G)$ of a graph G is a set such that each vertex $v \in V(G)$ is either in the set or adjacent to a vertex in the set. Matheson and Tarjan (1996) proved that any n -vertex plane triangulation has a dominating set of size at most $n/3$, and conjectured a bound of $n/4$ for n sufficiently large. King and Pelsmajer recently proved this for graphs with maximum degree at most 6. Plummer and Zha (2009) and Honjo, Kawarabayashi, and Nakamoto (2009) extended the $n/3$ bound to triangulations on surfaces.

We prove two related results: (i) There is a constant c_1 such that any n -vertex plane triangulation with maximum degree at most 6 has a dominating set of size at most $n/6 + c_1$. (ii) For any surface S , $t \geq 0$, and $\epsilon > 0$, there exists c_2 such that for any n -vertex triangulation on S with at most t vertices of degree other than 6, there is a dominating set of size at most $n(1/6 + \epsilon) + c_2$.

As part of the proof, we also show that any n -vertex triangulation of a non-orientable surface has a non-contractible cycle of length at most $2\sqrt{n}$. Albertson and Hutchinson (1986) proved that for n -vertex triangulation of an orientable surface other than a sphere has a non-contractible cycle of length $\sqrt{2n}$, but no similar result was known for non-orientable surfaces.

Keywords: Dominating set, triangulation, graphs on surfaces, non-contractible cycle, non-orientable surface.

Math. Subj. Class.: 05C69

*The author gratefully acknowledges support from NSA Grant H98230-08-1-0043 and the Swiss National Science Foundation, Grant No. 200021-125287/1.

E-mail addresses: hliu36@illinois.edu (Hong Liu), pelsmajer@iit.edu (Michael J. Pelsmajer)

1 Introduction

In this paper, we only consider graphs that are finite, undirected, and simple (no loops or multiple edges) except where specified otherwise.

A *dominating set* $D \subseteq V(G)$ of a graph G is a set such that each vertex $v \in V(G)$ is either in the set or adjacent to a vertex in the set. The *domination number* of G , denoted $\gamma(G)$, is defined as the minimum cardinality of a dominating set of G . Domination is very widely studied (see [3] for a recent monograph). A *triangulation* on a given surface is a graph embedded on the surface such that every face is bounded by a triangle.

In 1996, Matheson and Tarjan [8] proved $\gamma(G) \leq n/3$ for any n -vertex triangulated disc G (which includes all plane triangulations) and that this bound is sharp. Plummer and Zha [11] recently extended this bound to triangulations on the projective plane and proved $\gamma(G) \leq \lceil n/3 \rceil$ for triangulations on the torus or Klein bottle. Honjo, Kawarabayashi, and Nakamoto [4] obtained $\gamma(G) \leq n/3$ for triangulations on the torus and the Klein bottle and also for locally planar triangulations (triangulations of sufficiently high representativity) on every other surface. Matheson and Tarjan also gave an infinite class of plane triangulations with n vertices that requires $n/4$ vertices to be dominated. They conjectured that $\gamma(G) \leq n/4$ for every plane triangulation G with a finite number of exceptions, such as the octahedron, which has 6 vertices and domination number 2.

Conjecture 1.1 (Matheson and Tarjan [8]). *There exists n_0 such that any n -vertex plane triangulation with $n > n_0$ has a dominating set of size at most $n/4$.*

High degree vertices are helpful when constructing a small dominating set; this motivates the study of Conjecture 1.1 on graphs with no (or few) high degree vertices.

King and Pelsmayer [7] recently proved Conjecture 1.1 for plane triangulations with maximum degree at most 6. That is, they found n_0 such that $\gamma(G) \leq n/4$ for any n -vertex plane triangulation G with maximum degree at most 6 and $n > n_0$. (The proof has $n_0 = 4.5 \times 10^6$.)

Our theorems extend the previous results. First, we show that the degree restriction allows one to prove the following stronger upper bound, which verifies a conjecture by King and Pelsmayer [7].

Theorem 1.2. *There exists a constant c such that any n -vertex plane triangulation with maximum degree at most 6 has a dominating set of size at most $n/6 + c$.*

We prove Theorem 1.2 with $c = 1.05 \times 10^7$.

Maximum degree at most 6 implies that there are at most 12 vertices of degree less than 6, and all other vertices have degree 6. King and Pelsmayer conjectured that their result can be extended to plane triangulations with a bounded number of vertices with degree not equal to 6.

Conjecture 1.3 (King and Pelsmayer [7]). *For any constant t , there exists n_t such that any n -vertex plane triangulation with $n > n_t$ and at most t vertices of degree other than 6 has a dominating set of size at most $n/4$.*

They also gave examples [7] showing that the bound $n/4$ in Conjecture 1.3 cannot be improved beyond $n/6 + c$; the examples also imply that the bound in Theorem 1.2 is best possible.

One reason to restrict t rather than the maximum degree is to allow us to consider domination of similar degree-restricted triangulations on various surfaces. Indeed, it follows easily from Euler’s Formula that if S is any surface other than the plane (sphere), projective plane, torus, or Klein bottle, then there are no triangulations on S with maximum degree at most 6.

Theorem 1.4. *For any surface S , integer $t \geq 0$, and $\epsilon > 0$, there exists $c = c(S, t, \epsilon)$ such that any n -vertex triangulation on S , with at most t vertices of degree other than 6, has a dominating set of size at most $n(\frac{1}{6} + \epsilon) + c$.*

We prove Theorem 1.4 with $c = O((g^3 + gt^2)/\epsilon)$, where g is the genus of S . Since n is a trivial upper bound for domination number, Theorem 1.4 is only meaningful when $c(S, t, \epsilon) < n$, in which case $t = O(\sqrt{n})$.

Since $n(\frac{1}{6} + \epsilon) + c \leq \frac{n}{4}$ whenever $n(\frac{1}{12} - \epsilon) \geq c$, Conjecture 1.3 follows by setting $\epsilon < \frac{1}{12}$ (for example, let $\epsilon = 0.08$) and $n_t = c/(\frac{1}{12} - \epsilon)$.

In the proof of Theorem 1.4, we need to know that every n -vertex triangulation on a surface has a “small” non-contractible cycle. Albertson and Hutchinson [1] showed that for any fixed orientable surface, there is a non-contractible cycle of length at most $\sqrt{2n}$. We prove a similar bound for non-orientable surfaces.

Theorem 1.5. *Any n -vertex triangulation on a non-orientable surface has a non-contractible cycle of length less than or equal to $2\sqrt{n}$.*

Remark 1.6. The upper bound of $\sqrt{2n}$ in [1] was improved to $O(\sqrt{n/g} \log g)$ in [5], where g is the genus of the orientable surface. However, using this result in our proof will not reduce c below $c = O((g^3 + gt^2)/\epsilon)$ in the orientable case.

We will use the cycles to reduce a surface triangulation to sphere (or plane) triangulations, as described in Section 2. A closely related, well-known method of reducing a graph to a planar graph is to delete a small set of vertices (a *planarizing set*). Deleting a planarizing set does not necessarily yield a triangulation, so it is not clear whether it could be used to prove Theorem 1.4. Even if this was possible, it probably would not yield a stronger result, since the best known bound on the size of a minimum planar set is $O(\sqrt{gn})$ [2, 6], which is not much smaller than the total number of vertices involved in at most $O(g)$ non-contractible cycles that appear during the proof, and which is the main bottleneck in lowering the value of c .

By the following construction, the order of magnitude of the bound in Theorem 1.5 is best possible: Take an icosahedron, identify opposite points, and then triangulate each of the resulting 10 faces with a triangular grid of k^2 triangles. We get a projective planar triangulation with $10k^2$ faces, and hence $15k^2$ edges and $n = 5k^2 + 2$ vertices. Its shortest non-contractible cycle has length $3k$, which is nearly $\frac{3}{\sqrt{5}}\sqrt{n}$, or about $1.34164\sqrt{n}$.

Thus, the correct bound has order \sqrt{n} , and the constant multiple is between 1.34 and 2.

Question 1.7. What is the smallest constant that could replace 2 in Theorem 1.5?

In Section 3 we explain how to reduce the proof of Theorem 1.4 to finding small dominating sets for triangulations of spheres that contain certain pre-specified subsets of vertices. We show how to find such a dominating set in Section 4, and prove Theorem 1.2 as well. The proof of Theorem 1.2 follow the basic structure of the proof in [7]: we construct

two kinds of dominating sets and show that one must be small enough. The argument in Section 4 that finishes the proof of Theorem 1.4 is similar except that one of the sets only dominates part of the graph; we apply induction to dominate the rest of the graph.

Before that, we review some standard definitions and then introduce some definitions needed for our paper.

2 Definitions and preliminaries

For any graph G , let $V(G)$ and $E(G)$ denote the set of vertices and edges, respectively. For a vertex v in a graph G , $\deg(v)$ and $\deg_G(v)$ each denote the *degree* of v in G . $\Delta(G)$ is the maximum degree in G . For any $U \subseteq V(G)$, let $G[U]$ denote the subgraph of G induced by U , which has vertex set U and edge set $\{uv \in E(G) : u, v \in U\}$.

The *length* of a path, cycle, or walk W is the number of edges and it is denoted by $|W|$. Since a walk W may repeat edges, $|W|$ may exceed $|E(W)|$. If W is a cycle then $|V(W)| = |W|$ and if W is a path then $|V(W)| = |W| + 1$. A *chord* of a cycle or walk W is an edge not in $E(W)$ with endpoints in W .

The *distance* between vertices u and v , denoted $d(u, v)$, is the minimum length of a u, v -path. The distance between two sets (where each set could be a single vertex) is the minimum length of a path with one endpoint in each set. For a vertex v in a graph G and any integer $i \geq 0$, let $N_i(v) = \{u \in V(G) : d(v, u) = i\}$ and let $N_i[v] = \{u \in V(G) : d(v, u) \leq i\}$. Let G_i be the subgraph of G induced by $N_i[v]$, that is, $V(G_i) = N_i[v]$ and $E(G_i)$ contains the edges in G that have both endpoints in $N_i[v]$.

A graph is *connected* if there is a u, v -path for every pair of vertices u, v . A vertex is a *cut-vertex* if its removal increases the number of components of the graph. A graph with more than two vertices is *2-connected* if it is connected and it has no cut-vertices. A *block* is a maximal subgraph with no cut-vertices. Each block is either 2-connected, a single edge and its endpoints, or an isolated vertex. For any connected graph G , its *block-cutpoint tree* T is defined such that the blocks and cut-vertices of G are the vertices of T , and a block B and vertex v of G are adjacent in T if and only if B contains v .

For a graph G and vertex set $U \subseteq V(G)$, a *Steiner tree* for U is a minimum-size tree that contains U .

By a *surface*, we generally mean a 2-manifold without boundary. For $g \geq 0$, let S_g denote the orientable surface of orientable genus g (a sphere with g handles). For $g \geq 1$, let N_g denote the non-orientable surface of non-orientable genus g (a sphere with g cross-caps). By the classification theorem for surfaces, these are all the possible surfaces. For both $S = S_g$ and $S = N_g$, g is called the *genus* of S .

Given a graph G on a surface S such that every face is homeomorphic to a disk, $|V(G)| - |E(G)| + |F(G)|$ is the *Euler characteristic* of S , where $F(G)$ is the set of faces. It is $2 - 2g$ for $S = S_g$ and $2 - g$ for $S = N_g$; thus, it is independent of the choice of G . For any G we have the *degree-sum formula* $\sum_v \deg(v) = 2|E(G)|$. For triangulations we have $2|E(G)| = 3|F(G)|$, so the Euler characteristic equals $|V(G)| - |E(G)|/3 = \frac{1}{6} \sum_v (6 - \deg(v))$.

A closed curve on a surface is either *one-sided* or *two-sided*, and a surface is *orientable* if and only if it has no one-sided closed curves. A two-sided closed curve can be *contractible*, *surface-separating*, or neither; closed curves which are neither are called *essential*. A *plane graph* is a graph drawn in the plane without crossings; a *planar graph* is a graph that can be drawn as a plane graph. With respect to graph embeddings, the sphere

is usually interchangeable with the plane. See [9] for further background on graphs on surfaces.

Suppose that C is a non-contractible cycle of G on the surface S . We can obtain a triangulation G' on a related surface S' , which we call the C -derived graph of G , as follows. Cut the surface along C , and copy C locally onto each side of the cut so that each vertex and edge is doubled. This creates a triangulated surface with one hole if C is one-sided, and two holes if C is two-sided. Attach a disk to each hole to create a surface without boundary; this is S' . Add a vertex to each disk with edges to every vertex on the disk boundary; this yields G' , which is a triangulation of S' .

If C is one-sided, then C is replaced by a cycle C_1 of length $2|C|$, so $|V(G')| = |V(G)| + |C| + 1$, $|E(G')| = |E(G)| + 3|C|$, and $|F(G')| = |F(G)| + 2|C|$, where $F(G)$ is the set of faces of the embedded graph G . In this case, the Euler characteristics of S and S' differ by one. If C is two-sided, then C is replaced by cycles C_1, C_2 of length $|C|$, so $|V(G')| = |V(G)| + |C| + 2$, $|E(G')| = |E(G)| + 3|C|$, and $|F(G')| = |F(G)| + 2|C|$. In this case, the Euler characteristics of S and S' differ by two. If C is two-sided, then C may be *surface-separating*, in which case the Euler characteristic of S' equals the sum of the Euler characteristics of its two components.

Since the Euler characteristic of S_g is $2 - 2g$ and the Euler characteristic of N_g is $2 - g$, there are only certain possibilities for S' , which we summarize in Table 1.

Case	S	C	S'	
1	S_g	2-sided, non-separating	S_{g-1}	
2	S_g	2-sided, surface-separ.	$S_{k_1} \cup S_{k_2}$	$k_1, k_2 \geq 1$ and $k_1 + k_2 = g$
3	N_g	1-sided, non-separating	N_{g-1}	(only if $g \geq 2$)
4	N_g	1-sided, non-separating	$S_{(g-1)/2}$	(only if g is odd)
5	N_g	2-sided, non-separating	N_{g-2}	
6	N_g	2-sided, non-separating	$S_{(g-2)/2}$	(only if g is even)
7	N_g	2-sided, surface-separ.	$N_{k_1} \cup N_{k_2}$	$k_1, k_2 \geq 1$ and $k_1 + k_2 = g$
8	N_g	2-sided, surface-separ.	$S_{k_1/2} \cup N_{k_2}$	$k_1/2, k_2 \geq 1$ and $k_1 + k_2 = g$

Table 1: All possible C -derived surfaces S' , given the non-contractible cycle C on surface S with genus $g \geq 1$

The genus of S_g is also called *orientable genus* and the genus of N_g is also called *non-orientable genus*. Although it is somewhat unusual, it will be convenient for us to say that the *non-orientable genus of $S_{g/2}$ is g* (where g is even). (See Table 1 for motivation.)

An *outerplane* graph is a plane graph with all of its vertices incident to its outer face. For any graph G on a surface S , its *dual* is a graph on S with a vertex in each face of G , such that whenever two faces share an edge in G , the vertices corresponding to those faces are joined by an edge of the dual. The *weak dual* of an outerplane graph G is the dual minus the vertex in the outer face of G ; the weak dual is a forest, and it is a tree if G is 2-connected. The outer face of a connected outerplane graph H is bounded by a closed walk; without loss of generality, we may always assume that the walk is oriented counterclockwise, so that the exterior of H is always to the right.

The next two definitions allow us to state and work with a lemma from [7] (Lemma 4.6 in this paper) in the situation where the maximum degree is not bounded by 6.

Definition 2.1. Consider a closed walk $W = v_0, v_1, \dots, v_m = v_0$ (indexed by the cyclic

group Z_m) that bounds a connected outerplane subgraph H of a plane triangulation G , oriented counterclockwise, so that the exterior of H is always to the right.

For each $i \in Z_m$, let $\text{rdeg}_i(W)$ be the number of edges incident to v_i from the right—more specifically, if the edges incident to v_i are ordered such that they are counterclockwise near v_i , then count the ones that are after $v_i v_{i-1}$ and before $v_i v_{i+1}$. (The edges counted by $\text{rdeg}_i(W)$ lie in the exterior of H , and if v_i is not a cut-vertex of H , then $\text{rdeg}_i(W)$ counts every edge incident to v_i which lies in the exterior of H .)

The *outer degree sequence* of W is the cyclic sequence $\text{rdeg}_1(W), \dots, \text{rdeg}_m(W)$ indexed by Z_m .

We name a few special types of cyclic sequences that arise in the proof.

Definition 2.2. The cyclic sequence d_1, \dots, d_m (indexed by Z_m) is

Type A if $d_i = 2$ for all $i \in Z_m$,

Type B if $d_i = 3$ for some $i \in Z_m$ and $d_j = 2$ for all $j \in Z_m \setminus \{i\}$,

Type C if $d_i = 4$ for some $i \in Z_m$ and $d_j = 2$ for all $j \in Z_m \setminus \{i\}$,

Type D if $d_i = d_{i+1} = 3$ for some $i \in Z_m$ and $d_j = 2$ for all $j \in Z_m \setminus \{i, i + 1\}$,

Type E if $d_i = 3$ and $d_j = 1$ for some $i, j \in Z_m$ and $d_k = 2$ for all $k \in Z_m \setminus \{i, j\}$.

The following definition is from [7], albeit slightly renamed:

Definition 2.3. Let w, ℓ, k be integers with $w \geq 3$, $\ell \geq 1$, and $0 \leq k < w$. A (w, ℓ, k) -cylinder, (w, ℓ) -cylinder, or *triangulated cylinder* is any plane graph constructed as follows.

Fix an integer k with $0 \leq k < w$. Start with the Cartesian product of a w -cycle and a path of length ℓ . The vertices can be labeled $z_{a,b}$ with a in the cyclic group Z_w and $0 \leq b \leq \ell$. For each $0 \leq b < \ell$, add an edge from $z_{a,b}$ to $z_{a+1,b+1}$ if $0 \leq a < k$, and add an edge from $z_{a,b}$ to $z_{a-1,b+1}$ if $k < a \leq w$. All triangles (except those of the form $z_{0,b_0} z_{1,b_1} z_{2,b_2}$ when $w = 3$) are 3-faces.

The parameter w is the *width* and ℓ is the *length*. The cycles induced by $\{z_{a,0} : a \in Z_w\}$ and $\{z_{a,\ell} : a \in Z_w\}$ bound w -faces; these cycles are called the *boundary cycles*. The *interior* vertices are the vertices not in either boundary cycle.

Suppose that H is a (w, ℓ, k) -cylinder with $0 < k < w$ and let $V(H) = \{y_{a,b} : z \in Z_w, 0 \leq b \leq \ell\}$. Let $z_{w-a,b} = y_{a,b}$ for each $y_{a,b} \in V(H)$. With these new vertex names, we see that H is also a $(w, \ell, w - k)$ -cylinder. Thus, we may always assume that $0 \leq k \leq w/2$ for any (w, ℓ, k) -cylinder.

Note that a (w, ℓ) -cylinder has exactly $w(\ell + 1)$ vertices.

We consider one infinite graph: let G_∞ be the infinite 6-regular triangular grid (see Figure 1 on the left). There is a pattern of vertices from G_∞ that uses every seventh vertex (see the right side of Figure 1) and that dominates G_∞ ; let $D_\infty \subseteq V(G_\infty)$ be this (infinite) set of vertices.

3 Proof of Theorem 1.4

Given a triangulation G of an arbitrary surface, we can modify the graph by repeatedly picking a non-contractible cycle C and replacing its component G_C with the C -derived




Figure 1: G_∞ , with a dominating set D_∞ that contains every seventh vertex

graph of G_C . This continues until we have a graph G' that is the disjoint union of triangulations of spheres. Vertices of degree other than 6 in G' either came from vertices like that in G or from vertices that at some point were in one of these cycles C . We wish to keep track of these vertices and the way that they cluster.

Definition 3.1. For any graph G , we will always use the notation $U, U_0, \overline{U_0}$ for vertex sets and d_U for an integer that satisfy:

1. U is the disjoint union of U_0 and $\overline{U_0}$,
2. $\{v \in V(G) : \deg(v) \neq 6\} \subseteq U$,
3. every component of $G[U]$ contains at least one vertex of U_0 , and
4. for each $v \in \overline{U_0}$, there is a $u \in U_0$ with $d(v, u) \leq d_U$.

(For a graph called G^* , we use the notation $U^*, U_0^*, \overline{U_0^*}, d_{U^*}$ instead, etc.)

Note that for any triangulation G of a surface, we can satisfy Definition 3.1 by letting $U = U_0 = \{v \in V(G) : \deg(v) \neq 6\}$, $\overline{U_0} = \emptyset$, and $d_U = 0$.

Suppose that G is a triangulation of a surface with $U, U_0, \overline{U_0}, d_U$ that satisfy Definition 3.1, and C is a non-contractible cycle in G . The C -derived graph G' has two components G_1, G_2 if C is surface-separating and one component if C is non-separating. We wish to modify $U, U_0, \overline{U_0}, d_U$ so that Definition 3.1 is satisfied for the C -derived graph and for its components.

Lemma 3.2. *Suppose that G is a triangulation of a surface with $U, U_0, \overline{U_0}, d_U$ that satisfy Definition 3.1, C is a non-contractible cycle in G , and G' is the C -derived graph. If C is 2-sided, let C_1, C_2 be the new cycles that replace C and let v_1, v_2 be the new vertices in the disks bounded by C_1, C_2 . If C is 1-sided, then let C_1 be the new cycle that replaces C , and let v_1 be the new vertex in the disk bounded by C_1 .*

If C is 2-sided, then Definition 3.1 is satisfied for G' by $U' = (U \setminus V(C)) \cup V(C_1) \cup V(C_2) \cup \{v_1, v_2\}$, $U'_0 = (U_0 \setminus V(C)) \cup \{v_1, v_2\}$, $\overline{U'_0} = (\overline{U_0} \setminus V(C)) \cup V(C_1) \cup V(C_2)$, and $d'_{U'} = d_U + 1$. If G' has two components G_1, G_2 , then for $i = 1, 2$, Definition 3.1 is satisfied for G_i by $U^i = U' \cap V(G_i)$, $U^i_0 = U'_0 \cap V(G_i)$, $\overline{U^i_0} = \overline{U'_0} \cap V(G_i)$, and $d^i_{U'} = d'_{U'}$.

If C is 1-sided, then Definition 3.1 is satisfied for G' by $U' = (U \setminus V(C)) \cup V(C_1) \cup \{v_1\}$, $U'_0 = (U_0 \setminus V(C)) \cup \{v_1\}$, $\overline{U'_0} = (\overline{U_0} \setminus V(C)) \cup V(C_1)$, and let $d'_{U'} = d_U + 1$.

Proof. Suppose that C is 2-sided and $U', U'_0, \overline{U'_0}, d'_{U'}$ are as stated above. The properties (1) and (2) of Definition 3.1 are clearly satisfied for G' . (3) is also satisfied, since any component of $G'[U']$ that intersects C_1 (C_2) must also contain v_1 (v_2). Since property (4)

holds for G and $U, U_0, \overline{U_0}, d_U$, every vertex v of $\overline{U_0} \setminus V(C)$ is connected to a vertex of U_0 by a path in G with at most d_U edges. Hence, a minimal path P in G from v to $U_0 \cup V(C)$ has at most d_U edges. $E(P)$ gives us a path in G' from v to U'_0 or $C_1 \cup C_2$, and in the latter case it is one more edge away from v_1 or v_2 , which is in U'_0 . Since $d'_U = d_U + 1$, property (4) holds true for G' and $U', U'_0, \overline{U'_0}, d'_U$.

Hence, $U', U'_0, \overline{U'_0}, d'_U$ satisfy Definition 3.1 for G' . If C is surface-separating, then intersecting the sets with each component G_1, G_2 clearly gives sets that satisfy Definition 3.1 for each component with d'_U .

The same sort of argument works if C is 1-sided. (In this case, C cannot be surface-separating.) □

Remark 3.3. Still using the terminology of Lemma 3.2: Note that $|U'_0| - |U_0| \leq 2$ if C is 2-sided, and $|U'_0| - |U_0| \leq 1$ if C is 1-sided. Also, note that $|\overline{U'_0}| - |\overline{U_0}| \leq 2|C|$ (in both cases).

Moreover, suppose that we have a dominating set D' of G' that contains U' . D' contains $V(C_1) \cup V(C_2) \cup \{v_1, v_2\}$ if C is 2-sided and D' contains $V(C_1) \cup \{v_1\}$ if C is 1-sided. If we replace these vertices in D' by $V(C)$, we get a vertex set D which is a dominating set for G and contains U . Recall that G' has $|C| + 1$ or $|C| + 2$ more vertices than G , depending on whether C is 1-sided or 2-sided. Then $|D'| - |D| = |V(G')| - |V(G)|$.

Now consider graphs for which each component is a triangulation of a surface. If we repeatedly find a non-contractible cycle C , get the C -derived graph, and apply Lemma 3.2, we will end up with a graph G^* with $U^*, U^*_0, \overline{U^*_0}, d^*_U$ that satisfies Definition 3.1.

Let g_0, g_1, g_2 be the number of surface-separating, (non-separating) 1-sided, and non-separating 2-sided cycles C used during that process. Let $\sum |C|$ denote the sum of cycle-sizes, taken over all cycles used during the process. The total number of 2-sided cycles used is $g_0 + g_2$, so $|V(G^*)| - |V(G)| = \sum |C| + 2g_0 + g_1 + 2g_2$. By Remark 3.3, $|U^*_0| - |U_0| \leq 2g_0 + g_1 + 2g_2$ and $|\overline{U^*_0}| - |\overline{U_0}| \leq 2 \sum |C|$.

Suppose that D^* is a dominating set of G^* that contains U^* . Then by repeatedly applying Remark 3.3, we get a dominating set D of G that contains U , such that $|D^*| - |D| = |V(G^*)| - |V(G)|$.

Now, we are ready to start the proof of Theorem 1.4.

Suppose that we are given a triangulation G of an arbitrary surface S with n vertices, at most t of which have degree other than 6. We will modify the graph and surface until we have a graph G^* which is the disjoint union of triangulations of spheres, by repeatedly picking a minimum non-contractible cycle C in any current component H that is on a non-spherical surface, and replacing H by its C -derived graph H' . Let g_0, g_1, g_2 and $\sum |C|$ be as defined above.

Let the *genus-sum* of a graph be the sum of the genera of its components. Observe that the process ends precisely when the genus-sum reach zero.

According to Table 1: When C is surface-separating, the number of components increases by one, but the genus-sum is unchanged. When C is non-separating, the orientable genus-sum decreases by 1 if C is 2-sided, and the non-orientable genus-sum decreases by j if C is j -sided.

If $G = S_g$, then the cycle C is 2-sided in each step, so $g_1 = 0$ and it takes g steps to reduce the genus-sum to zero, so $g_2 = g$. If $G = N_g$, we must have $g_1 + 2g_2 = g$ in order for the genus-sum to be reduced to zero. In each step where C is surface-separating, the

number of components with non-zero genus is increased by one, so $g_0 \leq g - 1$, and in the end we have $g_0 + 1 \leq g$ components.

We may assume that $|U_0| = t$, $|\overline{U_0}| = 0$, and $d_U = 0$, by the comment following Definition 3.1. Then, we may conclude that $|\overline{U_0^*}| \leq 2 \sum |C|$, $|U_0^*| \leq t + 2g_0 + g_1 + 2g_2$, $|V(G^*)| = n + \sum |C| + 2g_0 + g_1 + 2g_2$, $|D^*| = |D| + \sum |C| + 2g_0 + g_1 + 2g_2$, and $d_U^* \leq g_0 + g_1 + g_2 \leq 2g - 1$.

We still need to find a dominating set D^* of G^* that contains U^* . To do this, we find a dominating set for each of its components H , which are triangulations of spheres such that $U^* \cap V(H)$, $U_0^* \cap V(H)$, $\overline{U_0^*} \cap V(H)$, and d_U^* satisfy Definition 3.1.

Lemma 3.4. *Let G be a triangulation of the sphere with n vertices and suppose that $U, U_0, \overline{U_0}, d_U$ satisfies Definition 3.1. Then G has a dominating set D that contains U such that*

$$|D| \leq \frac{n}{6} + 3(|U_0| - 1)(2\sqrt{3n} + 2d_U + 9) + \frac{3}{2}|\overline{U_0}| + \frac{1}{3}.$$

We defer the proof of Lemma 3.4 until later; For now, we assume that it is true. Applying Lemma 3.4 to each component of G^* , we get dominating sets for each component whose union D^* is a dominating set for G^* . Hence,

$$|D^*| \leq \sum_H \left(\frac{|V(H)|}{6} + 3(|U_0^H| - 1)(2\sqrt{3|V(H)|} + 2d_U^* + 9) + \frac{3}{2}|\overline{U_0^H}| + \frac{1}{3} \right),$$

where the sum is taken over all components H of G^* . Note that $\sum |V(H)| = |V(G^*)|$, $\sum |U_0^H| = |U_0^*|$, $\sum |\overline{U_0^H}| = |\overline{U_0^*}|$, and $\sum (|U_0^H| - 1)(2\sqrt{3|V(H)|}) \leq \sum (|U_0^H| - 1) \sum (2\sqrt{3|V(H)|})$. There are $g_0 + 1$ components H of G^* , so

$$\sum \sqrt{|V(H)|} \leq (g_0 + 1) \sqrt{(\sum |V(H)|)/(g_0 + 1)} = \sqrt{(g_0 + 1)|V(G^*)|},$$

and we get

$$|D^*| \leq \frac{|V(G^*)|}{6} + 3(|U_0^*| - g_0 - 1) \left(2\sqrt{3(g_0 + 1)|V(G^*)|} + 2d_U^* + 9 \right) + \frac{3}{2}|\overline{U_0^*}| + \frac{1}{3}(g_0 + 1).$$

Note that $|U_0^*| - g_0 - 1 \leq t + g_0 + g_1 + 2g_2 - 1$. Also, $|D^*| = |D| + \sum |C| + 2g_0 + g_1 + 2g_2$, $d_U^* \leq 2g - 1$, $|U_0^*| \leq t + 2g_0 + g_1 + 2g_2$, $|\overline{U_0^*}| \leq 2 \sum |C|$, and $g_0 \leq g - 1$, so

$$|D| \leq \frac{|V(G^*)|}{6} + 3(t + g + g_1 + 2g_2 - 2) \left(2\sqrt{3g|V(G^*)|} + 4g + 7 \right) + 2 \sum |C| - g_1 - 2g_2 + \frac{1}{3}.$$

Also, $|V(G^*)| = n + \sum |C| + 2g_0 + g_1 + 2g_2$. To continue, we need an upper bound for $\sum |C|$.

As stated in Section 1, every n -vertex triangulation of a non-spherical surface has a non-contractible cycle C with $|C| \leq \sqrt{2n}$ if the surface is orientable and $|C| \leq 2\sqrt{n}$ if the surface is non-orientable.

Let $f(m) = f^{(1)}(m) = \lfloor m + \sqrt{2m} + 2 \rfloor$ for $m \geq 3$. For $i > 1$ and $m \geq 3$, let $f^{(i)}(m) = f(f^{(i-1)}(m))$. Given a triangulation of an orientable surface with at most m vertices, there is a non-contractible cycle C such that the C -derived graph has at most $f(m)$ vertices. If G is embedded on S_g , then there are $g_2 + g_0$ cycles considered during the process that produces G^* , and at every stage of the process, every component is orientable. Therefore, $|V(G^*)| \leq f^{(g_2+g_0)}(n)$, where $n = |V(G)|$.

The size of each C in the process depends on the size of the component H containing C , which could be part of a graph at any stage in the process before the end (when G^* has been obtained). The last C considered in the process cannot be surface-separating, so each time we consider a new cycle C , at most $g_2 - 1$ non-separating cycles C have already been considered. Also, surface-separating cycles produce two components which each have fewer vertices than their source component. Therefore, the number of vertices in a component H that contains any of the cycles C is at most $f^{(g_2-1)}(n)$. Since there are $g_2 + g_0$ cycles C considered during the entire process, we obtain $\sum |C| \leq (g_2 + g_0)\sqrt{2f^{(g_2-1)}(n)}$. Since $g_2 = g$ and $g_2 + g_0 \leq 2g - 1$, we get $|V(G^*)| \leq f^{(2g-1)}(n)$ and $\sum |C| \leq (2g - 1)\sqrt{2f^{(g-1)}(n)}$.

Let $F(m) = F^{(1)}(m) = \lfloor m + 2\sqrt{m} + 1 \rfloor$ for $m \geq 3$. For $i > 1$ and $m \geq 3$, let $F^{(i)}(m) = F(F^{(i-1)}(m))$. Note that $F(m) \geq f(m)$ for all $m \geq 3$. Given a triangulation of a non-orientable surface with at most m vertices, there is a non-contractible cycle C such that the C -derived graph has at most $F(m)$ vertices. If G is embedded on N_g , then there are $g_1 + g_2 + g_0$ cycles C considered during the process, and there can be orientable and non-orientable components during the process. Since $f(m) \leq F(m)$, we have $|V(G^*)| \leq F^{(g_1+g_2+g_0)}(n)$. As before, taking the C -derived graph does not increase the size of any component H which contains some non-contractible cycle from later in the process, if C is surface-separating cycles or if C is the last non-separating cycle considered. Continuing as before, in this case we can obtain $\sum |C| \leq (g_1 + g_2 + g_0)2\sqrt{F^{(g_1+g_2-1)}(n)}$. Since $g_1 + g_2 + g_0 \leq 2g - 1$ and $g_1 + g_2 - 1 \leq g - 1$, we get $|V(G^*)| \leq F^{(2g-1)}(n)$ and $\sum |C| \leq (2g - 1)2\sqrt{F^{(g-1)}(n)}$.

Using these bounds for $|V(G^*)|$ and $\sum |C|$, we can rewrite our bound for $|D|$ in terms of $f^{(i)}(n)$ when G is embedded on an orientable surface, and in terms of $F^{(i)}(n)$ when G is embedded on an orientable surface.

Thus, we need to find bounds on $f^{(i)}(n)$ and $F^{(i)}(n)$.

Claim: $f^{(i)}(n) \leq n + i\sqrt{2n} + i^2 + 3i - 2$ and $F^{(i)}(n) \leq n + 2i\sqrt{n} + i^2$.

Proof by induction: For $i = 1$, $f^i(n) = f(n) \leq n + \sqrt{2n} + 1 + 3 - 2$, as desired.

Suppose that $i \geq 2$. By induction, $f^{(i-1)}(n) \leq n + (i-1)\sqrt{2n} + (i-1)^2 + 3(i-1) - 2 = n + i\sqrt{2n} - \sqrt{2n} + i^2 + i - 4$. Therefore, $f^{(i)}(n) = \lfloor f^{(i-1)}(n) + \sqrt{2f^{(i-1)}(n)} + 2 \rfloor$ is at most

$$n + i\sqrt{2n} - \sqrt{2n} + i^2 + i - 4 + \sqrt{2 \left[n + i\sqrt{2n} - \sqrt{2n} + i^2 + i - 4 \right]} + 2.$$

This is at most the claimed upper bound $n + i\sqrt{2n} + i^2 + 3i - 2$ if and only if

$$\sqrt{2 \left[n + i\sqrt{2n} - \sqrt{2n} + i^2 + i - 4 \right]} \leq \sqrt{2n} + 2i.$$

Equivalently (squaring both sides), this is

$$2 \left[n + i\sqrt{2n} - \sqrt{2n} + i^2 + i - 4 \right] \leq 2n + 4i\sqrt{2n} + 4i^2.$$

which is true if and only if

$$0 \leq (2i + 2)\sqrt{2n} + 2i^2 - 2i + 8,$$

which is true.

To prove $F^{(i)}(n) \leq n + 2i\sqrt{n} + i^2$, observe that $F(n) = \lfloor (\sqrt{n} + 1)^2 \rfloor$ and that the desired bound can be rewritten as $F^{(i)}(n) \leq (\sqrt{n} + i)^2$.

When $i = 1$, we have $F^{(1)}(n) = F(n) \leq (\sqrt{n} + 1)^2$, as desired. Suppose that $i \geq 2$. By induction, $F^{(i-1)}(n) \leq (\sqrt{n} + i - 1)^2$. Then $F^{(i)}(n) = (\sqrt{F^{(i-1)}(n)} + 1)^2 \leq ((\sqrt{n} + i - 1) + 1)^2 = (\sqrt{n} + i)^2$. Thus the claim is proved.

Note that $\sqrt{f^{(i)}(n)} \leq \sqrt{F^{(i)}(n)} \leq \sqrt{n} + i$. Now we are ready to finish the proof of Theorem 1.4.

First, consider the case that G is a triangulation of S_g ($g \geq 1$). We have

$$|V(G^*)| \leq f^{(2g-1)}(n) \leq n + (2g - 1)\sqrt{2n} + (2g - 1)^2 + 3(2g - 1) - 2,$$

$$\sqrt{|V(G^*)|} \leq \sqrt{f^{(2g-1)}(n)} \leq \sqrt{n} + 2g - 1,$$

and

$$\sum |C| \leq (2g - 1)\sqrt{2f^{(g-1)}(n)} \leq (2g - 1)\sqrt{2}(\sqrt{n} + g - 1).$$

Together with

$$|D| \leq \frac{|V(G^*)|}{6} + 3(t + g + g_1 + 2g_2 - 2) \left(2\sqrt{3g|V(G^*)|} + 2g + 7 \right) + 2 \sum |C| - g_1 - 2g_2 + \frac{1}{3},$$

$g_0 \leq g - 1$, $g_1 = 0$, and $g = g_2$, we can get $|D| \leq \frac{n}{6} + a\sqrt{n} + b$, where

$$a = \frac{\sqrt{2}}{6}(2g - 1) + 6\sqrt{3g}(t + 3g - 2) + 2\sqrt{2}(2g - 1)$$

and

$$b = 3(t + 3g - 2) \left(2\sqrt{3g}(2g - 1) + 4g + 7 \right) + 2\sqrt{2}(2g - 1)(g - 1) + \frac{2}{3}g^2 + \frac{7}{3}g - \frac{1}{3}.$$

Note that a and b depend only on g and t , with $a = O(g^{1/2}(g+t))$ and $b = O(g^{3/2}(g+t))$.

Next, consider the case that G is a triangulation of N_g . We have

$$|V(G^*)| \leq F^{(2g-1)}(n) \leq n + 2(2g - 1)\sqrt{n} + (2g - 1)^2,$$

$$\sqrt{|V(G^*)|} \leq \sqrt{n} + 2g - 1,$$

and

$$\sum |C| \leq (2g - 1)2\sqrt{F^{(g-1)}(n)} \leq 2(2g - 1)(\sqrt{n} + g - 1).$$

Using the same bound for $|D|$ with $g_0 \leq g - 1$ and $g_1 + 2g_2 = g$, we again can get $|D| \leq \frac{n}{6} + a\sqrt{n} + b$, but this time with

$$a = \frac{1}{3}(2g - 1) + 6\sqrt{3g}(t + 2g - 2) + 2(2g - 1)$$

and

$$b = 3(t + 2g - 2) \left(2\sqrt{3g}(2g - 1) + 4g + 7 \right) + 4(2g - 1)(g - 1) + \frac{2}{3}g^2 - \frac{2}{3}g + \frac{1}{2}.$$

Again, a and b depend only on g and t , with $a = O(g^{1/2}(g + t))$ and $b = O(g^{3/2}(g + t))$.

If G is a triangulation of the sphere S_0 , a direct application of Lemma 3.4 with $|U_0| = t$, $|\overline{U}_0| = 0$, and $d_U = 0$ gives $|D| \leq \frac{n}{6} + a\sqrt{n} + b$ with $a = 6\sqrt{3}(t - 1)$ and $b = 27(t - 1) + \frac{1}{3}$. So $a = O(g^{1/2}(g + t))$ and $b = O(g^{3/2}(g + t))$ is valid for all surfaces.

To prove Theorem 1.4, it suffices to have $\frac{n}{6} + a\sqrt{n} + b \leq n(\frac{1}{6} + \epsilon) + c$, for some $c = c(S_g, t, \epsilon)$. Let $c = a^2/(4\epsilon) + b$. Then the previous inequality is equivalent to $0 \leq n\epsilon - a\sqrt{n} + a^2/(4\epsilon)$, which is true since the right side equals $(\sqrt{n\epsilon} - \frac{a}{2\sqrt{\epsilon}})^2$.

Since $a^2 = O(g^3)$ and $b = O(g^{5/2})$ for $g \geq t$, and $a^2 = O(gt^2)$ and $b = O(g^{3/2}t)$ for $t \geq g$, we get $a^2 + b = O(a^2) = O(g^3 + gt^2)$. This yields $c = O(a^2/\epsilon) = O((g^3 + gt^2)/\epsilon)$.

Remark 3.5. There must be a constant c' such that $c = (c'(g^3 + gt^2)/\epsilon) (1 + o(1))$. This constant is different for S_g and N_g .

Using the stronger Hutchinson bound of $|C| \leq O(\sqrt{n/g} \log g)$ instead of $|C| \leq \sqrt{2n}$ for the orientable case would improve c' , but not $O((g^3 + gt^2)/\epsilon)$.

The argument for the non-orientable case would also work for S_{2g} , where g is non-orientable genus. But the constant c' we would get is not as good as the current c' for the orientable case.

4 Proof of Lemma 3.4 and Theorem 1.2

For convenience, we restate Lemma 3.4 and Theorem 1.2 as a single lemma.

Lemma 4.1. *Let G be a triangulation of the sphere with n vertices and suppose that $U, U_0, \overline{U}_0, d_U$ satisfies Definition 3.1. Then G has a dominating set D that contains U such that*

$$|D| \leq \frac{n}{6} + 3(|U_0| - 1)(2\sqrt{3n} + 2d_U + 9) + \frac{3}{2}|\overline{U}_0| + \frac{1}{3}.$$

Moreover, if $\Delta(G) \leq 6$ and $U = U_0 = \{v \in V(G) : \deg(v) \neq 6\}$, then

$$|D| \leq \frac{n}{6} + 1.05 \times 10^7.$$

By the comment following Definition 3.1, when $\Delta(G) \leq 6$ we may assume that $U = U_0 = \{v \in V(G) : \deg(v) \neq 6\}$ and $d_U = 0$; hence Lemma 4.1 does indeed imply Theorem 1.2.

From Euler’s formula it follows that

$$\sum_{u \in U} (\deg(u) - 6) = -12. \tag{4.1}$$

G has minimum degree at least 3 since otherwise G must be a triangle, in which case any one vertex gives us a sufficiently small dominating set. Then, by Equation 4.1, $|U| \geq 4$. Note that if $\Delta(G) \leq 6$ and $U = \{v \in V(G) : \deg(v) \neq 6\}$, then by Equation 4.1, $|U| \leq 12$.

Let T_0 be a Steiner tree for U_0 in G ; that is, let T_0 be a tree in G such that $U_0 \subseteq V(T_0)$ and T_0 is of minimum size. Let $U'_0 = U_0 \cup \{v \in V(T_0) : \deg_{T_0}(v) \neq 2\}$.

Suppose that $|U_0| > 1$. Let L_0 be the set of leaves in T_0 ; then $|L_0| \geq 2$ and $L_0 \subseteq U_0$. One can prove by induction that a tree with $k \geq 2$ leaves has at most $k - 2$ vertices of degree greater than 2. Then we have $|U'_0| \leq 2|L_0| - 2 \leq 2|U_0| - 2$.

Since U'_0 contains $\{v \in V(T_0) : \deg_{T_0}(v) \neq 2\}$, $E(T_0)$ can be partitioned by the maximal paths in T_0 with no internal vertices in U'_0 (and endpoints in U'_0). There are $|U'_0| - 1$ such paths; let P_0 be such a path of maximum length. Then $|P_0|(|U'_0| - 1) \geq |E(T_0)|$. Since $|V(T_0)| = |E(T_0)| + 1$ and $|U'_0| \leq 2|U_0| - 2$, we get

$$|V(T_0)| \leq (2|U_0| - 3)|P_0| + 1.$$

(If $\Delta(G) \leq 6$ and $U = U_0 = \{v \in V(G) : \deg(v) \neq 6\}$, then $|U_0| \leq 12$, so $|V(T_0)| \leq 21|P_0| + 1$.)

If $|U_0| \leq 1$, then by $|U| \geq 4$ and Definition 3.1(3), we must have $|U_0| = 1$. In this case, $|V(T_0)| = 1$.

Since every component of $G[U]$ contains a vertex of U_0 and T_0 contains U_0 , $T_0 \cup G[U]$ is connected. Let T be a spanning tree in $T_0 \cup G[U]$. So, T contains U and $|V(T)| \leq |V(T_0)| + |\overline{U_0}|$. (If $|U_0| \leq 1$, $|V(T)| = 1 + |\overline{U_0}|$.)

Next, we define G' : make two copies of each edge of T and, for each vertex $v \in V(T)$, make $\deg_T(v)$ copies of v . Draw these all near the original edges and vertices, and create incidences in the natural way so that we obtain a plane graph with one face f_T that contains T (before T is deleted), and the other faces are all 3-faces (that correspond to the faces of G). (See Figure 2 for an example.) Note the boundary of f_T is a cycle. For convenience, let us reembed G' in the plane such that f_T is the outer face. Ignoring f_T , we have a triangulated disc. Let V'_T be the vertices in G' copied from $V(T)$; then $G' - V'_T = G - V(T)$.




Figure 2: An example of constructing G' near a portion of T

The following lemma was originally stated only for the case where G' is constructed from a graph G that has maximum degree at most 6, but it applies (with the exact same proof) whenever G' is a triangulated disk and all its interior vertices have degree equal to 6.

Lemma 4.2. [King and Pelsmajer [7]] *Suppose that G' is a triangulated disk and all its interior vertices have degree equal to 6. Then G' can be mapped to G_∞ such that vertices are sent to vertices, edges to edges, and interior 3-faces to 3-faces, such that adjacent 3-faces in G' are mapped to distinct 3-faces in G_∞ .*

Let g be a map from G' to G_∞ as described in Lemma 4.2. Note that g is not necessarily injective. There is a pattern of vertices from G_∞ that uses every seventh vertex (see the

right side of Figure 1). Let $D_\infty \subseteq V(G_\infty)$ be the (infinite) set of vertices indicated in the figure. For each vertex $v \in V(G')$, $g(v) \in D_\infty$ or $g(v)$ is adjacent to a vertex in D_∞ . If $v \notin V'_T$ then the seven vertices of $N[v]$ map to the seven vertices of $N[g(v)]$; as this includes one vertex of D_∞ , v is dominated by $g^{-1}(D_\infty) = \{v \in V(G') : g(v) \in D_\infty\}$. Therefore G' is dominated by the union of $g^{-1}(D_\infty)$ and V'_T , or equivalently the union of $g^{-1}(D_\infty) - V'_T$ and V'_T . Let $D' = g^{-1}(D_\infty) - V'_T$ and let D be the union of D' and $V(T)$. By the construction of G' from G , D is a subset of $V(G)$ that dominates every vertex of G .

In Subsection 4.1, we find upper bounds for $|D|$ in terms of n and $|V(T)|$. In [7], where the maximum degree is at most 6, such a bound is much easier to find.

4.1 Upper bounds for $|D|$

Consider any distinct $v_1, v_2 \in D'$ with $g(v_1) \neq g(v_2)$. Since $D' \subseteq G' - V'_T$, $N[v_1]$ and $N[v_2]$ are 7-vertex subsets of G' . Then, by Lemma 4.2, g maps $N[v_1]$ and $N[v_2]$ bijectively to $N[g(v_1)]$ and $N[g(v_2)]$, respectively. According to the right side of Figure 1, since $g(v_1) \neq g(v_2)$, $N[g(v_1)]$ and $N[g(v_2)]$ are disjoint. Then $N[v_1]$ and $N[v_2]$ must also be disjoint. Therefore, if $v_1, v_2 \in D'$ and $N[v_1] \cap N[v_2] \neq \emptyset$, then $g(v_1) = g(v_2)$.

Lemma 4.3. *If $g(v_1) = g(v_2)$ in G_∞ where $v_1, v_2 \in V(G')$ are distinct vertices, and $u \in N[v_1] \cap N[v_2]$, then $\deg(u) > 6$ and there are at least 5 edges between uv_1 and uv_2 in the rotation at u in the drawing of G' .*

Proof. If v_1, v_2 are adjacent in G' , then they are mapped to adjacent (hence distinct) vertices by Lemma 4.2. So v_1 and v_2 are not adjacent, and $u \notin \{v_1, v_2\}$. The 3-faces of G' that are incident to u will either form a path or a 6-cycle in the dual of G' . According to the map g , the images of the faces under g will again be consecutive around $g(u)$, so the number of these faces between v_1 and v_2 must be $6i$ for some positive integer i . The number of edges between uv_1 and uv_2 in the rotation at u is $6i - 1 \geq 5$, and thus $\deg(u) \geq (6i - 1) + 2 \geq 7$. (See Figure 3.) □




Figure 3: Two vertices v_1, v_2 with $g(v_1) = g(v_2)$ and $u \in N[v_1] \cap N[v_2]$

Let t' be the number of vertices in G' of degree greater than 6, and for each vertex $v \in D'$ let t'_v be the number of vertices in $N[v]$ with degree greater than 6.

Consider any $v \in D'$, and suppose that $u \in N[v] \cap N[v']$ for some $v' \in D'$ with $v \neq v'$. By the paragraph preceding Lemma 4.3, $g(v) = g(v')$. Then, by Lemma 4.3, the degree of u is greater than 6. Therefore, vertices of degree at most 6 in $N[v]$ are not in $N[v']$ for any $v' \in D'$ with $v \neq v'$. Since $v \in D' \subseteq G' - V'_T$, $N[v]$ is a 7-vertex subset of G' . Hence, $N[v]$ contains $7 - t'_v$ vertices of degree at most 6, which are not in $N[v']$ for any $v' \in D'$ with $v \neq v'$. Therefore, $\sum_{v \in D'} (7 - t'_v) \leq |V(G')| - t'$. We can write

$$|D'| = \frac{1}{6} \sum_{x \in D'} 6 \leq \frac{1}{6} \left(|V(G')| - t' + \sum_{v \in D'} (t'_v - 1) \right)$$

and soon, we will bound $\sum_{v \in D'} (t'_v - 1)$.

If $\Delta(G) \leq 6$, then any vertex with degree greater than 6 in G' must be in V'_T and must, as a vertex in G , be a leaf of T . If we also have $U = U_0 = \{v \in V(G) : \deg(v) \neq 6\}$, then T is a Steiner tree for U , so every leaf of T has degree less than 6. Therefore, if $\Delta(G) \leq 6$ and $U = U_0 = \{v \in V(G) : \deg(v) \neq 6\}$, then t' and every t'_v is zero, and the results of the previous paragraph simplify to $7|D'| \leq |V(G')|$, or $|D'| \leq \frac{1}{7}|V(G')|$.

To bound $\sum_{v \in D'} (t'_v - 1)$, we make an auxiliary plane graph H : Let $V(H)$ be the set of all vertices in G' that have degree greater than 6. For each $v \in D'$, let $B[v]$ be the union of the six triangles incident to v in the embedding of G' in the plane (including the interior and boundary of each triangle). Then each $B[v]$ is a hexagon with t'_v vertices of H on its boundary. For distinct $x, y \in D'$, the interiors of the hexagons $B[x]$ and $B[y]$ are disjoint; the hexagons may intersect on their boundaries. Now, for each $v \in D'$ with $t'_v \geq 2$, draw a $(t'_v - 1)$ -leaf star in $B[v]$ on the t'_v vertices of $V(H)$ in $B[v]$, such that the edges are drawn on the interior of $B[v]$. (See Figure 4 for an example.) Let $E(H)$ be the set of all such edges. Then H is a plane graph with t' vertices and $\sum_{v \in D'} (t'_v - 1)$ edges.




Figure 4: An example of a star drawn in $B[v]$ when $t'_v = 4$

Lemma 4.4. H is an outerplanar graph with no multiple edges.

Proof. Assume there are two edges in H sharing the same endpoints, say x, y , which means there are vertices $u, v \in D'$ such that $B[u]$ and $B[v]$ each contain both x and y . Let Q be G' restricted to the quadrilateral $xuyv$ and its interior, as shown in Fig 5. By Lemma 4.3, there are at least 5 edges incident to x between the two edges ux and vx in the quadrilateral, so $\deg_Q(x) \geq 7$; similarly $\deg_Q(y) \geq 7$. All the vertices in the interior of G' have degree 6, and $\deg_Q(u) \geq 2, \deg_Q(v) \geq 2$, so we have $\sum_{z \in Q} \deg(z) \geq 6(|V(Q)| - 4) + 2 + 2 + 7 + 7 = 6|V(Q)| - 6$. Also, $\sum_{z \in Q} \deg(z) = 2|E(Q)|$, so $|E(Q)| \geq 3|V(Q)| - 3$. However, Q is planar, so Euler's formula yields $|E(Q)| \leq 3|V(Q)| - 6$. This is a contradiction, so H has no multiple edges.

The vertices of degree greater than 6 in G' are all on the boundary of G' , so all these vertices are incident to the outer face of H as well. These are the vertices of H , so H is outerplanar. \square

By Lemma 4.4, $|E(H)| \leq 2|V(H)| - 3$, so $\sum_{v \in D'} (t'_v - 1) \leq 2t' - 3$. Then $|D'| \leq \frac{1}{6}(|V(G')| - t' + 2t' - 3) = \frac{1}{6}(|V(G')| + t' - 3)$. Since $|V(G')| = n - |V(T)| + |V'_T|$ and $|D| = |D'| + |V(T)|$, we obtain $|D| \leq \frac{1}{6}(n + 5|V(T)| + |V'_T| + t' - 3)$. Clearly, $t' \leq |V'_T|$, so $|D| \leq \frac{1}{6}(n + 5|V(T)| + 2|V'_T| - 3)$. When we defined G' , we made $\deg_T(v)$ copies of v for all $v \in V(T)$, so $|V'_T| = \sum_{v \in T} \deg_T(v) = 2|E(T)| = 2|V(T)| - 2$. Therefore,

$$|D| \leq \frac{1}{6}(n + 9|V(T)| - 7).$$




Figure 5: The quadrilateral $xuyv$ and its interior in H

Recall that if $\Delta(G) \leq 6$ and $U = U_0 = \{v \in V(G) : \deg(v) \neq 6\}$, then $|D'| \leq \frac{1}{7}|V(G')|$. Since $|V(G')| = n - |V(T)| + |V'_T|$ and $|V'_T| = 2|V(T)| - 2$, in this case we have $|D| = |V(T)| + |D'| \leq |V(T)| + \frac{1}{7}(n - |V(T)| + 2|V(T)| - 2)$, or

$$|D| \leq \frac{1}{7}(n + 8|V(T)| - 2).$$

Thus we have found upper bounds for $|D|$ in terms of n and $|V(T)|$. When this bound does not suffice, we will need to find a different dominating set.

4.2 When G contains a large triangulated cylinder

Within this subsection, we assume that $|U_0| > 1$.

Recall that P_0 is a longest path in T_0 such that no internal vertices are in U'_0 . Let v_1, v_2 be the endpoints of the path P_0 . Let x be a middle vertex of P_0 , that is, let x be a vertex on P_0 of distance $\lfloor |P_0|/2 \rfloor$ from an endpoint of P_0 . Let $N_i(x)$ be the set of vertices of G with distance exactly i from x , let $N_i[x]$ be the set of vertices of G with distance at most i from x , and let G_i be the graph induced by $N_i[x]$.

The $d_U = 0$ case of the following result was obtained in [7], using mostly the same proof.

Lemma 4.5. $N_i(x)$ does not intersect U for $i < \lfloor |P_0|/2 \rfloor - d_U$.

Proof. Suppose that j is the smallest index such that $N_j(x)$ intersects U_0 . Since each $u \in \overline{U_0}$ has distance at most d_U to a vertex of U_0 , vertices of $N_i(x)$ with $i < j - d_U$ do not intersect U .

Let $u \in U_0 \cap N_j(x)$ and let Q be an x, u -path of length j (which is in G_j). There is a unique x, u -path in T_0 ; without loss of generality, assume that it contains v_1 (rather than v_2). By the choice of P_0 , deleting the interior of its x, v_1 -subpath from T_0 gives a 2-component graph that contains U_0 . We could then add Q to that graph to obtain a connected graph that contains U_0 , and let T'_0 be a spanning tree of it. Then, $|V(T'_0)| \leq |V(T_0)| - (\lfloor |P_0|/2 \rfloor - 1) + (j - 1)$. But $|V(T_0)| \leq |V(T'_0)|$ since T_0 is a Steiner tree for U_0 , so $j \geq \lfloor |P_0|/2 \rfloor$. \square

Let r be minimum such that G_r is not a triangulated hexagon. Then G_{r-1} accounts for $1 + \sum_{i=1}^{r-1} 6i$ distinct vertices, so $n > 1 + 6r(r-1)/2 > 3(r-1)^2$.

Next, we seek to understand the structure of G_i for values $i \geq r$. Our immediate goal is Lemma 4.8; Lemmas 4.6 and 4.7 help us obtain it. In [7] the result stated in Lemma 4.8 was obtained for graphs of maximum degree at most 6, using a different proof.

From Subsection 2, recall the definitions *outer degree sequence* of a walk that bounds a connected outerplane subgraph of G and *type A, B, C, D, and E cyclic sequences*.

Lemma 4.6 (King and Pelsmajer [7]). *Suppose that x is a vertex in a plane triangulation G , and r is the minimum such that G_r (the graph induced by $N_r[x]$) is not a triangulated hexagon. If every vertex in $N_r[x]$ has degree 6, then $N_r[x]$ contains a cycle C' of length at most $2r + 1$ of type A, B, C, D, or E. Moreover, if C' is type A or E then $|C'| \in \{2r, 2r + 1\}$.*

In [7], walks of type C and D appear when “ $q = 1$ ”, type B appear when “ $q = 2$ ” and type A and E arise when “ $q = 3$ ”.

Lemma 4.7. *Suppose that $W = v_0, v_1, \dots, v_m = v_0$ is a walk (indexed by the cyclic group Z_m) that bounds an outerplane subgraph H of a plane triangulation G such that W is oriented counterclockwise, that is, with the exterior of H always to its right. Suppose that W is type A, B, C, D, or E and suppose that every vertex of W has degree 6 in G .*

Then W is a cycle of G . Furthermore, the neighbors of W on the interior of H form a connected outerplane graph H' , bounded by a counterclockwise walk W' with length and type specified by Table 3.

Proof. If H has only one vertex v , then $\deg(v) = \text{rdeg}_0(W)$. If H contains a leaf v , which is the i th vertex of W , then $\deg(v) = 1 + \text{rdeg}_i(W)$. Since $\deg(v) = 6$ for all $v \in V(H)$ and $\text{rdeg}_i(W) \leq 4$ for all $i \in Z_m$, both cases give a contradiction.

Suppose that H contains a 2-connected leaf-block B with cut-vertex v , with v appearing as the i th and j th vertex of W as it enters and exits B . Then $\deg_B(v) \geq 2$ and $\deg_H(v) \geq \deg_B(v) + 1$, so $\text{rdeg}_i(W) + \text{rdeg}_j(W) \leq \deg(v) - \deg_H(v) \leq 6 - 3 = 3$. Then the outer degree sequence of W must be type E with $\{\text{rdeg}_i(W), \text{rdeg}_j(W)\} = \{1, 2\}$. Also, the three previous inequalities must be equalities; in particular, $\deg_H(v) = 3$. Therefore, v is incident to a single-edge (cut-edge) block of H , and it and B are the only two blocks of H that contain v .

Since the previous argument applies to any 2-connected leaf-block of H , there is at most one 2-connected leaf-block in H . Since H has no leaves, the block-cutpoint tree of H has at most one leaf. However, any nontrivial tree has at least two leaves, so the block-cutpoint tree must have only one vertex. That is, H must be 2-connected. Then W is a cycle.

Since W is a cycle, each vertex v_i in W is incident to exactly two edges in W and $\text{rdeg}_i(W)$ edges on the exterior of W . Also $\deg(v_i) = 6$, so v_i is incident to exactly $4 - \text{rdeg}_i(W)$ edges on the interior of W . (See Table 2, ignoring the last column.)

If W is of type C and v_i is its vertex with $\text{rdeg}_i(W) = 4$, then v_i is incident to no edges on its interior, so $v_{i-1}v_iv_{i+1}$ is a triangle on the interior of H . Removing v_i from W yields a cycle of length $m - 1$ of type D that bounds the outerplane graph $H - v_i$. For convenience, rename W , m , and H to be these values instead. Now W is of type A, B, D, or E. Note that every value of $4 - \text{rdeg}_i(W)$ is 2 except for at most one 3 and up to two 1s.

Suppose that W has at least one chord on its interior. (A *chord* of W is an edge of $G - E(W)$ with endpoints in W .) By the choice of H , we may assume without loss of generality that $E(H)$ is the union of $E(W)$ and all chords of W that lie in the interior of W . Then the weak dual of H is a nontrivial tree, with at least two leaf-faces. Let v_iv_j be

$rdeg_i(W)$	number of such i	$4 - rdeg_i(W)$	$3 - rdeg_i(W)$
3	at most two	1	0
4	at most one	0	(none)
1	at most one	3	2
2	$m, m - 1$ or $m - 2$	2	1

Table 2: The number of edges incident to v_i on the exterior of W , the frequency of each case, the number of edges incident to v_i on the interior of W , and the number of edges of W' that form a triangle with v_i (when W is of type A, B, C, D, or E and $m = |W|$)

a chord incident to a leaf-face f of the weak dual of H , and let B be the boundary cycle of f . For any edge xy in B , there is a triangle xyz in G in $B \cup f$. B is chordless, so either z is in f or $B = xyz$. If $B = xyz$, then f is a face of G ; letting $\{x, y\} = \{v_i, v_j\}$, z is incident to zero edges on the interior of W — a contradiction. Hence, z is in f . It follows that every vertex of B , including v_i and v_j , is incident to an edge in f . At most one vertex of W is incident to more than two edges on its interior. Since v_i and v_j are each incident to an edge in f and the edge v_iv_j , without loss of generality we can assume that v_i is incident to no other edges in the interior of W . Let v_iv_jz be the unique triangle of G that contains v_iv_j and is not in $B \cup f$. Then v_iz must be an edge of W . As before, v_iv_jz does not bound a leaf-face of the weak dual of H , because then z would be incident to zero edges on the interior of W . Therefore v_jz must be a chord of W . Hence, v_j is the unique vertex of W that is incident to 3 edges on the interior of W . Recall that H has a leaf-face $f' \neq f$. Applying the same argument to f' , we find that f' is also incident to v_j , and that v_j is incident to an edge in f' . This edge is not a chord and it is not in f , so v_j is incident to four edges in the interior of W — a contradiction. Thus, we have shown that W has no chords in its interior.

For each $i \in Z_m$, the vertices adjacent to v_i in the interior of W form a path Q_i , such that for each edge xy in Q_i , v_ixy bounds a face of G . Let W' be the closed walk obtained by concatenating these paths. By construction, W' bounds an outerplane graph, oriented in the counterclockwise direction. Q_i has $4 - rdeg_i(W)$ vertices, so its length is $3 - rdeg_i(W)$. See Table 2 on the right. Thus, each v_i in W yields one edge in W' , except if $rdeg_i(W)$ is 3 or 1, in which case v_i yields 0 or 2 edges in W' . This shows that W' has the length specified by Table 3 when W is type A, B, D, or E.

Type of W	length of W'	Type of W'
A	m	A
B	$m - 1$	B
C	$m - 3$	C
D	$m - 2$	C
E	m	E

Table 3: How the type and length of W determine the type and length of W' (where $m = |W|$)

For each v'_i in W' , $rdeg_i(W')$ is the number of paths Q_i that contains v'_i . Thus, when Q_i and Q_{i+1} have length at least one, they meet at a vertex v'_i with $rdeg_i(W') = 2$; when Q_i has length two (when W is type E) it yields on vertex v'_i with $rdeg_i(W') = 1$; when

$|Q_i| = |Q_{i+1}| = 0$ (type D) or $|Q_i| = 0$ (type B), there is a vertex v'_i with $\text{rdeg}_i(W') = 4$ or $\text{rdeg}_i(W') = 3$. Altogether, this shows that W' has the type indicated in Table 3, when W is type A, B, D, or E.

When W is type C, then it was replaced by a walk of type D and length $m - 1$, so by the previous two paragraphs, W' should have length $(m - 1) - 2$ and be type C, as indicated in Table 3. \square

When $N_{3r+1}[x]$ does not intersect U , we will produce a *triangulated cylinder* (recall definition from Subsection 2).

Lemma 4.8. *Suppose that G is a plane triangulation, with P_0 , x , and r defined as before. If $N_{3r+1}[x]$ does not intersect U , then G contains a (w, ℓ) -cylinder H with no interior vertices in U , such that $w \in \{2r, 2r + 1\}$ and $\ell \geq 2(\lfloor |P_0|/2 \rfloor - r - d_U - 1)$.*

Proof. Since $N_{3r+1}[x]$ does not intersect U , every vertex in $N_{3r+1}[x]$ has degree 6. By Lemma 4.6, $N_r[x]$ contains a cycle W_0 of length at most $2r + 1$ of type A, B, C, D, or E. W_0 bounds an outerplanar subgraph of G , and it can be oriented counterclockwise. For any $i \geq 1$, let N_i^* be the set of vertices in the interior of W_0 at distance exactly i from W_0 . Let j be minimum such that $N_j^* = \emptyset$ or N_j^* intersects U . Since W_0 is in $N_r[x]$, $j > 0$. Then we can repeatedly apply Lemma 4.7 for $i = 0, \dots, j - 1$ with $W = W_i$, which proves that W_i is a cycle of type A, B, C, D, or E and produces the closed (nonempty) walk $W_{i+1} = W'$ on vertex set N_{i+1}^* . Therefore, W_j is nonempty, so N_j^* must contain a vertex of U .

According to Table 3, if W_0 is type B, C, or D, then for all $1 \leq i \leq 2r + 1$, every W_i is type B or C. Then $|W_{i+1}| \leq |W_i| - 1$ for all $0 \leq i \leq j - 1$, so $|W_j| \leq |W_0| - j$. W_j is nonempty and $|W_0| \leq 2r + 1$, so $0 < 2r + 1 - j$. Since W_0 is in $N_r[x]$, $N_i^* \subseteq N_{r+i}[x]$ for all $i \geq 1$. Then $N_j^* \subseteq N_{r+j}[x] \subseteq N_{3r+1}[x]$. However, $N_{3r+1}[x]$ does not intersect U , so this is a contradiction. Thus we may assume that W_0 is type A or E, in which case W_i is the same type and length as W_0 for all $1 \leq i \leq j$, and $|W_0| \in \{2r, 2r + 1\}$ by Lemma 4.6. Let $w = |W_0|$. W_i is a w -cycle for all $0 \leq i \leq j - 1$.

For any $i \geq 1$, let N_{-i}^* be the set of vertices in the exterior of W_0 at distance exactly i from W_0 , and let k be the minimum such that $N_{-k}^* = \emptyset$ or N_{-k}^* contains a vertex of degree not equal to 6. Now, reembed G in the plane such that the interior and exterior of W_0 are switched and repeat the previous argument. This yields w -cycles W_{-i} on N_{-i}^* for all $0 \leq i \leq k - 1$. The cycles W_i for $-(k - 1) \leq i \leq j - 1$ and the edges between consecutive cycles form a (w, ℓ) -cylinder with $w = |W_0|$ and $\ell = j + k - 2$.

N_j^* and N_{-k}^* each contain a vertex of U , so $N_{r+j}[x]$ and $N_{r+k}[x]$ do too, since W_0 is in $N_r[x]$. By Lemma 4.5, $r + j \geq \lfloor |P_0|/2 \rfloor - d_U$ and $r + k \geq \lfloor |P_0|/2 \rfloor - d_U$. Then $\ell \geq j + k - 2 \geq 2(\lfloor |P_0|/2 \rfloor - r - d_U - 1)$. \square

The following results are proved in [7].

Lemma 4.9 (King and Pelsmajer [7]). *Suppose that G is a plane triangulation and that $\Delta(G) \leq 6$. If H is a (w, ℓ) -cylinder in G with ℓ maximal, then $G - V(H)$ has at most $w(w - 1)$ vertices.*

Lemma 4.10 (King and Pelsmajer [7]). *Suppose that H is a (w, ℓ) -cylinder. Then H has $w(\ell + 1)$ vertices and there is a set of $\lceil \frac{\ell}{7} \rceil(w + 2)$ vertices on H that dominates all vertices on the interior of H .*

We will see that the size $\lceil \frac{\ell}{7} \rceil (w + 2)$ is efficient enough — that is, it uses roughly one-sixth of the vertices on the (w, ℓ) -cylinder or less — when $w \geq 12$. However, we must prove a new result for the cases when w is small.

Lemma 4.11. *If H is a (w, ℓ) -cylinder and $3 \leq w \leq 12$, then H contains a set of at most $\frac{|V(H)|}{6} + 12 = \frac{1}{6}w(\ell + 1) + 12$ vertices that dominates the interior vertices of H .*

Proof. Let $Z = \{z_{a,b} : a \in Z_w, 0 \leq b \leq \ell\}$ be the vertex set of H , and note that $|Z| = w(\ell + 1)$. Let $Z' = \{z_{a,b} : a \in Z_w, 0 < b < \ell\}$, the vertices on the interior of H .

For all integers w, k with $3 \leq w \leq 12$ and $0 \leq k < w$, we will give an integer $m = m(w, k)$ and a set $S \subseteq \{z_{a,b} : a \in Z_w, 0 \leq b < m\}$ such that (i) every $z_{a,b} \in Z'$ is dominated by some $z_{c,d} \in Z$ such that $z_{c,d \bmod m} \in S$, and (ii) $|S| \leq \min(\frac{1}{6}mw, 12)$. If we have such m and S , then $S_Z = \{z_{a,b} \in Z : z_{a,b \bmod m} \in S\}$ is a set of size at most $|S| \lceil (\ell + 1)/m \rceil$ in Z that dominates Z' . Since $|S| \lceil (\ell + 1)/m \rceil \leq |S|(\frac{\ell}{m} + 1) \leq \frac{1}{6}w\ell + 12$, this will finish the proof.

Thus, it remains to find such m, S for all w, k such that $3 \leq w \leq 12$ and $0 \leq k < w$. Recall that we may assume that $0 \leq k \leq w/2$. In each case of the proof, we describe S via a figure where $z_{a,b}$ is located on row a and column b , showing rows 0 to w (row 0 and row w are identified) and columns 0 to at least m . When $b = 0 \equiv w \pmod w$, then $z_{a,b}$ is shown twice, but once as a hollow dot (for example, in Figure 6). The figure will make it clear that S_Z dominates Z' as desired.

We begin with two general cases: when w is a multiple of 2 or 3.

Consider any case where w is a multiple of 3. No matter what k is, any three consecutive rows of Z' can be dominated by taking every other vertex on the middle row. (For example, see Figure 6 on the left). Thus, we have m, S with $m = 2$ and $|S| = w/3$. Clearly S_Z dominates Z' and $|S| = \frac{1}{6}mw \leq 12$, so this suffices.




Figure 6: Dominating sets when $w = 6$: an example with $m = 2$ and $k = 2$, with $3m$ columns shown (left) and an example with $m = 3$ and $k = 3$, with $2m$ columns shown (right)

Next, consider any case where w is even. Let $m = 3$, and S will contain one vertex from each even-indexed row. Then $|S| = \frac{1}{2}w = \frac{1}{6}mw \leq 6$. Clearly, S_Z will dominate all vertices in each even-indexed row of Z' . It remains to show that for any k , S can be constructed so that S_Z dominates all vertices in all odd-indexed rows of Z' .

For each row $2i$, S contains either $z_{2i,0}$, $z_{2i,1}$, or $z_{2i,2}$, and this “offset” determines the entire pattern on that row in S_Z . (For example, see Figure 6 on the right.) Once the offset

is chosen for row $2i$, two of the three possible offsets for row $2i + 2 \pmod w$ will ensure that all vertices in row $2i + 1 \pmod w$ of Z' are dominated. Thus, starting with $z_{2,1}$ in S_Z for row 2, the offsets for rows $2i$ with $1 \leq i \leq w/2 - 1$ can be chosen so that all vertices in rows $3, \dots, w - 3$ of Z' are dominated. Finally, of the three possible offsets for row 0, two will ensure that all vertices in row $w - 1$ of Z' are dominated, and two will ensure that all vertices in row 1 of Z' are dominated; hence, there is an offset for row 0 so that both rows are dominated.

It remains to consider the cases $w = 5, 7, 11$, for all $0 \leq k < w$. We consider each of these cases separately, giving a figure that shows S in $\{z_{a,b} : 0 \leq b < m\}$ in the appendix and noting that $|S| \leq 12$ and $\frac{|S|}{mw} \leq \frac{1}{6}$ in Table 4. This completes the proof of Lemma 4.11. \square

w	k	Figure	$ S $	m	$ S /(mw)$
5	0	8	4	5	4/25
5	1	8	5	7	1/7
5	2	8	5	6	1/6
7	0	9	7	7	1/7
7	1	9	8	7	8/49
7	2	10	8	7	8/49
7	3	10	8	7	8/49
11	0	11	12	7	12/77
11	1	11	12	7	12/77
11	2	11	9	5	9/55
11	3	12	12	7	12/77
11	4	12	12	7	12/77
11	5	12	12	7	12/77

Table 4: All cases with $w = 5, 7, 11$

At this point, we break the argument into two proofs.

4.3 Finishing the proof of Theorem 1.2

Suppose that $\Delta(G) \leq 6$ and $U = U_0 = \{v \in V(G) : \deg(v) \neq 6\}$. Then $T = T_0, |\overline{U_0}| = d_U = 0, |U_0| = |U| \leq 12$, and there is a dominating set D of G with $|D| \leq \frac{n+8|V(T)|-2}{7}$. Also, $|U_0| > 1$ because $|U| \geq 4$, so we have P_0 and r with $n > 3(r - 1)^2$, and Lemma 4.5 applies.

Let $c = 1.05 \times 10^7$. If $|D| \leq n/6 + c$ then we are done, so we may assume that $\frac{1}{7}(n + 8|V(T)| - 2) > n/6 + c$, or equivalently, $n < 48|V(T)| - 42c - 12$. $|V(T)| = |V(T_0)| \leq (2|U_0| - 3)|P_0| + 1 \leq 21|P_0| + 1$, so $n < 1008|P_0| - 42c + 36$ and $|P_0| > \frac{n-36+42c}{1008}$. And since $c = 1.05 \times 10^7$, $|P_0| > \frac{42c-36}{1008} > 42 \times 10^4$.

Claim: Every vertex in $N_{3r+1}[x]$ has degree 6.

By Lemma 4.5, it is true if $3r + 1 < \lfloor |P_0|/2 \rfloor$. Suppose that it is false. Then $3r + 1 \geq \lfloor |P_0|/2 \rfloor \geq (|P_0| - 1)/2$, so $r \geq (|P_0| - 3)/6$. Since $n > 3(r - 1)^2$, we get $n > \frac{1}{12}(|P_0|^2 - 18|P_0| + 81)$. Then $n < 1008|P_0| - 42c + 36$ gives $12(1008|P_0| - 42c + 36) > |P_0|^2 - 18|P_0| + 81$, or equivalently, $12114 + (-504c + 351)/|P_0| > |P_0|$. Which contradicts $|P_0| > 42 \times 10^4$. Thus the claim is proved.

Now we may apply Lemma 4.8 to obtain a (w, ℓ, k) -cylinder H with ℓ maximized. By Lemmas 4.8 and 4.9, $w = 2r$ or $w = 2r + 1$, $\ell \geq |P_0| - 2r - 3$, and $n \leq w(\ell + 1) + w(w - 1) = w(\ell + w)$. By Lemma 4.10, H contains a set S_H of at most $\lceil \frac{\ell}{7} \rceil(w + 2)$ vertices that dominates its interior. $V(G) - V(H)$ dominates itself and the boundary of H , so if we add $V(G) - V(H)$, we get a set that dominates G ; it has size at most $\lceil \frac{\ell}{7} \rceil(w + 2) + n - w(\ell - 1)$. We are done if this is at most $\frac{n}{6} + c$, so we may assume that $\lceil \frac{\ell}{7} \rceil(w + 2) + n - w(\ell - 1) > \frac{n}{6} + c$. Since $\lceil \frac{\ell}{7} \rceil \leq \frac{\ell + 6}{7}$ and $n \leq w(\ell + w)$, we get $w(\ell - 1) - \frac{(\ell + 6)(w + 2)}{7} + c < \frac{5}{6}n \leq \frac{5}{6}w(\ell + w)$. It follows that $(w - 12)\ell < 35w^2 + 78w + 72 - 42c$.

First, consider the case $w \geq 13$. Then $\ell < (35w^2 + 78w + 72 - 42c)/(w - 12) = 35w + 498 + \frac{6048 - 42c}{w - 12}$. Since $c > 6048$ and $w \leq 2r + 1$, we have $\ell < 35w + 498 \leq 70r + 533$. Since $\ell \geq |P_0| - 2r - 3$, we have $|P_0| < 72r + 536$. With $r < 1 + \sqrt{n/3}$ and $|P_0| > \frac{n - 36 + 42c}{1008}$, we can obtain $n - 36 + 42c - 1008(24\sqrt{3n} + 608) < 0$. Let $f(x) = x^2 - 1008 \cdot 24\sqrt{3}x - 1008 \cdot 608 - 36 + 42c$; then $f(\sqrt{n}) < 0$. Since $f(x)$ is a quadratic function with a positive quadratic term, $f(x)$ must have two roots. Therefore $(1008 \cdot 24\sqrt{3})^2 - 4(-1008 \cdot 608 + 42c - 36) > 0$, which contradicts $c = 1.05 \times 10^7$ (but not by much, which explains our choice of c). This completes the case $w \geq 13$.

Next, suppose that $3 \leq w \leq 12$. By Lemma 4.11, H contains a set of size S that dominates all vertices on its interior of H with $|S| \leq |V(H)|/6 + 12$. If we add $V(G) - V(H)$ and all the vertices of the boundary of H to S , we get a set that dominates G ; its size is at most $|V(H)|/6 + 12 + w(w - 1) + 2w$. Since $w \leq 12$ and $H \subseteq G$, we have $|V(H)|/6 + 12 + w(w - 1) + 2w \leq n/6 + 12 + 132 + 24 < n/6 + c$, as desired. \square

4.4 Finishing the proof of Lemma 3.4

Recall that there is a dominating set D of G with $|D| \leq \frac{n + 9|V(T)| - 7}{6}$.

If $|U_0| = 1$, then $|V(T)| = 1 + |\overline{U_0}|$, so

$$|D| \leq \frac{n + 9|\overline{U_0}| + 2}{6} = \frac{n}{6} + \frac{3}{2}|\overline{U_0}| + \frac{1}{3},$$

as desired. Thus, we may assume that $|U_0| > 1$.

We have P_0 and r with $n > 3(r - 1)^2$. Also, by Lemma 4.5, every vertex in $N_i[x]$ has degree 6 and is not in U if $i < \lfloor |P_0|/2 \rfloor - d_U$.

Case 1 $|P_0| < 2\sqrt{3n} + 2d_U + 9$.

Then, since $|V(T_0)| \leq (2|U_0| - 3)|P_0| + 1$ and $|V(T)| \leq |V(T_0)| + |\overline{U_0}|$, we have $|V(T)| < (2|U_0| - 3)(2\sqrt{3n} + 2d_U + 9) + |\overline{U_0}| + 1$. G has a dominating set D with $|D| \leq \frac{n + 9|V(T)| - 7}{6}$. Then

$$|D| < \frac{n}{6} + \frac{3}{2}(2|U_0| - 3)(2\sqrt{3n} + 2d_U + 9) + \frac{3}{2}|\overline{U_0}| + \frac{1}{3}.$$

Case 2 $|P_0| \geq 2\sqrt{3n} + 2d_U + 9$.

Since $n > 3(r - 1)^2$, we get $\sqrt{3n} > 3(r - 1)$, which yields $|P_0| > 6r + 2d_U + 3$, so $|P_0| \geq 6r + 2d_U + 4$. Then $\lfloor |P_0|/2 \rfloor - d_U > 3r + 1$, so by Lemma 4.5, every vertex in $N_{3r+1}[x]$ has degree 6 and is not in U . By Lemma 4.8, G has a (w, ℓ) -cylinder H with no interior vertices in U , such that $w \in \{2r, 2r + 1\}$ and $\ell \geq 2(\lfloor |P_0|/2 \rfloor - r - d_U - 1)$. Since $\lfloor |P_0|/2 \rfloor \geq \frac{1}{2}(|P_0| - 1)$ and $\sqrt{3n} > 3(r - 1)$, we get

$$\ell > 2 \left(\frac{1}{2}(2\sqrt{3n} + 2d_U + 9 - 1) - \frac{1}{3}\sqrt{3n} - 1 - d_U - 1 \right) = 4(1 + \sqrt{n/3}).$$

By Lemma 4.11, the interior of the triangulated cylinder can be dominated by a set S_Z of at most $\frac{w(\ell+1)}{6} + 12$ vertices if $3 \leq w \leq 12$. In this case, $12 \leq 4w$, so we get $|S_Z| < \frac{w\ell}{6} + 5w$. If $w \geq 13$, then by Lemma 4.10, the interior of the triangulated cylinder can be dominated by a set S_Z of at most $\lceil \frac{\ell+1}{7} \rceil (w+2)$ vertices. Since $\lceil \frac{\ell+1}{7} \rceil \leq \frac{\ell}{7} + 1$, in this case we have $|S_Z| \leq (\frac{\ell}{7} + 1)(w+2) = \frac{w\ell}{6} - \frac{(w-12)\ell}{42} + (w+2) < \frac{w\ell}{6} + 2w$. Thus, we have $|S_Z| < \frac{w\ell}{6} + 5w$ for all $w \geq 3$.

In order to apply induction on n , we delete the $w(\ell - 1)$ interior vertices of (w, ℓ) -cylinder H . Let C_1 and C_2 be the boundary cycles of H ; these now bound holes in the surfaces. Recall that every w -cycle on the cylinder has the exact same pattern of turns: none, or exactly one right turn and exactly one left turn which are at the same places around each cycle. Thus we can identify C_1 and C_2 such that corresponding turns are matched to each other. Thus, when two vertices of degree 6 are identified, the resulting vertex will have degree 6. In this way, identify each pair of corresponding vertices v_1, v_2 from C_1, C_2 to get a new vertex v^* and a new w -cycle C^* . This creates a new plane (or sphere) triangulation G^* with $n - w\ell$ vertices.

Still using v_1, v_2, v^* to represent corresponding vertices on C_1, C_2, C^* , we define disjoint subsets $U_0^*, \overline{U_0^*}$ of $V(G^*)$ as follows: If v_1 or v_2 is in U_0 , then put v^* in U_0^* ; also, let $U_0^* \setminus V(C^*) = U_0 \setminus V(H)$. Note that $|U_0^*| \leq |U_0|$. If v_1 or v_2 is in $\overline{U_0}$ and $v^* \notin U_0^*$, then put v^* in $\overline{U_0^*}$; also, let $\overline{U_0^*} \setminus V(C^*) = \overline{U_0} \setminus V(H)$. Note that $|\overline{U_0^*}| \leq |\overline{U_0}|$. Let $U^* = U_0^* \cup \overline{U_0^*}$; note that $U^* \setminus V(C^*) = U \setminus V(H)$ and that for each $v^* \in U^* \cap V(C^*)$, v_1 or v_2 is in $V(C) \cap U$.

We wish to show that Definition 3.1 is satisfied by G^* with $U^*, U_0^*, \overline{U_0^*}, d_U$.

A vertex of degree not equal to 6 in G^* is also a vertex of degree other than 6 in G if it is not in $V(C^*)$, and if $v^* \in V(C^*)$ does not have degree 6 then at least one of v_1 or v_2 does not have degree 6. Therefore every vertex of degree not equal to 6 in G^* is in U^* .

Any vertex $x \in \overline{U_0^*}$ corresponds to a vertex $y \in \overline{U_0}$ (either $x = v^* \in V(C^*)$ and $y \in \{v_1, v_2\}$, or $x \notin V(C^*)$ and $y = x$). There is a path P in G from y to U_0 of length at most d_U . The vertices of P that lie in H can be replaced by vertices on C^* in a natural way so that we get a walk in G^* from x to U_0^* of length at most $|P|$. Therefore, any vertex in $\overline{U_0^*}$ has distance at most d_U in G^* to U_0^* .

Since U does not intersect the interior of H , each component of $G[U]$ becomes a connected subgraph of $G^*[U^*]$ that contains at least one vertex of U_0^* . Each component of $G^*[U^*]$ is the union of some of these subgraphs, so it also intersects U_0^* .

Definition 3.1 is satisfied for G^* with $U^*, U_0^*, \overline{U_0^*}, d_U$, so we can apply induction. We get a dominating set D^* that contains U^* such that

$$|D^*| \leq \frac{n - w\ell}{6} + 3(|U_0^*| - 1)(2\sqrt{3(n - w\ell)} + 2d_U + 9) + \frac{3}{2}|\overline{U_0^*}| + \frac{1}{3}.$$

Since $|U_0^*| \leq |U_0|$ and $|\overline{U_0^*}| \leq |\overline{U_0}|$, we get

$$|D^*| \leq \frac{n - w\ell}{6} + 3(|U_0| - 1)(2\sqrt{3(n - w\ell)} + 2d_U + 9) + \frac{3}{2}|\overline{U_0}| + \frac{1}{3}.$$

Temporarily set x so that $\sqrt{n} - \sqrt{n - w\ell} = xw$. Then $\sqrt{n} - xw = \sqrt{n - w\ell}$, so $n + x^2w^2 - 2xw\sqrt{n} = n - w\ell$. Then $x^2w + \ell = 2x\sqrt{n}$. Since $\ell > 4(1 + \sqrt{n/3})$ and $x^2w > 0$, we get $4(1 + \sqrt{n/3}) < 2x\sqrt{n}$. Then $x > 2\sqrt{1/3}$, so $\sqrt{n} - \sqrt{n - w\ell} > 2\sqrt{1/3}w$

and $2\sqrt{3(n - w\ell)} < 2\sqrt{3n} - 4w$. It follows that

$$|D^*| < \frac{n - w\ell}{6} + 3(|U_0| - 1)(2\sqrt{3n} - 4w + 2d_U + 9) + \frac{3}{2}|U_0| + \frac{1}{3}.$$

G is dominated by the union of $D^* - V(C^*)$, S_Z , and $V(C_1) \cup V(C_2)$; this set D has size at most $|D^*| + (\frac{w\ell}{6} + 5w) + 2w$. Since $|U_0| \geq 2$, we have $3(|U_0| - 1)(-4w) + 5w + 2w \leq -5w < 0$. Thus, we get

$$|D| < \frac{n}{6} + 3(|U_0| - 1)(2\sqrt{3n} + 2d_U + 9) + \frac{3}{2}|U_0| + \frac{1}{3}$$

Thus, whether $|U_0| \leq 1$, or whether we are in one of the two cases where $|U_0| > 1$, we obtain a dominating set D for G of the desired size. This finishes the proof of the Lemma 3.4. \square

5 Small non-contractible cycles in non-orientable surfaces

In this section we prove Theorem 1.5.

First we prove it for any triangulation G on the projective plane N_1 . Given G on N_1 , let C be a minimum-length non-contractible cycle. C must be one-sided. Cut along C and double C alongside the cut, as in the C -derived construction from Section 2, but do not add the disk, nor the extra vertex that goes in the disk. This yields a triangulated disk G' bounded by a cycle C' of length $2|C|$. Label the vertices of C' in clockwise order, as $v_0, v_1, \dots, v_{|C|} = v'_0, v'_1, \dots, v'_{|C|} = v_0$; then v_j and v'_j (for any $0 \leq j \leq |C|$) are copies of the same vertex in C . Let $m = \lfloor |C|/2 \rfloor$, let $x = v_m$, and for all $j \geq 0$, let V_j be the set of vertices v in G' such that the distance $d(v, x) = j$.

Lemma 5.1. V_j contains a path P_j from v_{m-j} to v_{m+j} of length at least $2j$, for all j with $0 \leq j \leq m$.

Proof. Any path P in G' between opposite vertices v_j, v'_j of C corresponds to non-contractible cycle in C ; then by the choice of C , the length of P is at least $|C|$. Recall that for any vertices u, v , any u, v -walk contains a u, v -path.

Suppose that W is a v_i, v_j -walk (or path) in G' with $0 \leq i < j \leq m$. Then $v_0, \dots, v_i, W, v_j, \dots, v_{|C|} = v'_0$ is a v_0, v'_0 -walk in G' , which contains a v_0, v'_0 -path P in G' . Since the length of P must be at least $|C|$, the length of W must be at least $j - i$.

For $0 \leq j \leq m$, we can apply the previous observation where the indices are $(m - j, m)$ or $(m, m + j)$ and conclude that $v_{m \pm j} \notin V_i$ for any $i < j$. Since $v_m, v_{m-1}, \dots, v_{m-j}$ and $v_m, v_{m+1}, \dots, v_{m+j}$ are paths of length j , we have $v_{m \pm j} \in V_j$ for all $0 \leq j \leq m$. If we apply the same observation where the indices are $m - j$ and $m + j$, we can conclude that any v_{m-j}, v_{m+j} -walk (or path) in G' has length at least $2j$, for any $0 \leq j \leq m$.

Thus, it remains to show that G' contains a v_{m-j}, v_{m+j} -path P_j with $V(P_j) \subseteq V_j$, for all $0 \leq j \leq m$. We prove this by induction. It is trivial for $j = 0$ since $V_0 = \{x\}$. Assume that it is true for fixed j with $0 \leq j < m$. We must prove it for $j + 1$.

Without loss of generality, we may assume that P_j is v_{m-j}, v_{m+j} -path with vertices in V_j of minimum length. Then P_j is an *induced path*, that is, there is no edge between non-consecutive vertices of P_j .

P_j divides the triangulated disk into two faces; let f be the face that does not contain x . For any $y_k \in V_k$, there is a path y_k, y_{k-1}, \dots, y_0 with each $y_i \in V_i$. If y_k is in f , then

this path must intersect P_j at y_i with $0 \leq i < k$, which implies that $k > j$. Therefore, f contains no vertex of $\bigcup_{0 \leq i \leq j} V_i$.

Let $P_j = y_1, y_2, \dots, y_p$, where $v_{m-j} = y_1$ and $y_p = v_{m+j}$. A triangle in f with more than one endpoint on P_j must intersect $V(P_j)$ at two consecutive endpoints y_i, y_{i+1} , since P_j is an induced path and G' has no multiple edges. Every edge $y_i y_{i+1}$ of P_j is incident to exactly one triangle in f ; let z_i be its third vertex, which is in f . Let $z_0 = v_{m-j-1}$ and let $z_p = v_{m+j+1}$. (See Figure 7.) For each y_i in P_j , there is a set of triangles in f such that y_i is their only vertex in P_j , naturally ordered by the embedding near y_i ; removing y_i yields a z_{i-1}, z_i -walk in f . Concatenating these walks gives a z_0, z_p -walk W in f . Since every vertex of W is incident to a vertex of P_j , and W is in f , the vertices of W must be contained in V_{j+1} . W contains a z_0, z_p -path; let this be P_{j+1} . \square




Figure 7: Construction of P_{j+1} in f , when P_j is assumed to be of minimum length

Lemma 5.2. Any n -vertex triangulation on the projective plane has a non-contractible cycle of length less than or equal to $2\sqrt{n} - 1$.

Proof. Since the vertex sets V_0, \dots, V_m are disjoint, the vertex sets of the paths P_0, \dots, P_m are disjoint. Therefore,

$$n \geq \sum_{j=0}^m |V(P_j)| \geq \sum_{j=0}^m (2j + 1) = (m + 1)^2.$$

Then $2\sqrt{n} - 1 \geq 2m + 1 = 2\lfloor |C|/2 \rfloor + 1 \geq |C|$. \square

Lemma 5.3. Any n -vertex triangulation on a non-orientable surface with genus $g > 1$ has a non-contractible cycle of length less than or equal to $2\sqrt{n}$.

Proof. For any non-orientable surface N_g , it is known that the double cover of N_g is S_{g-1} . Let G be an n -vertex triangulation of N_g with $g > 1$, and let H be the double cover of G . Then H is a $2n$ -vertex triangulation on S_{g-1} , an orientable surface that is not the sphere. By [1], H has a non-contractible cycle C_H of length at most $\sqrt{2(2n)} = 2\sqrt{n}$.

C_H maps to a closed walk C_G on N_g . If C_G is contractible, then there is a homotopy in S_{g-1} from C_G to a point. It lifts to a homotopy in N_g from C_H to a point ([10, Lemma

54.2], for example). Then C_H is contractible in N_g , which is a contradiction. Therefore, C_G is a non-contractible closed walk in G of length at most $|C_H|$. C_G must contain a non-contractible cycle, which has length at most $|C_G| \leq |C_H| \leq 2\sqrt{n}$. \square

The previous two lemmas complete the proof of Theorem 1.5.

Acknowledgements

We would like to thank Erin W. Chambers for pointing us in the direction of double covers and Erika L.C. King for allowing us to use two figures from [7]. We would also like to thank the referees for their careful work, which led to critical improvements in the paper.

References

- [1] M. O. Albertson and J. P. Hutchinson, On the independence ratio of a graph, *J. Graph Theory* **2** (1978), 1–8.
- [2] H. N. Djidjev, On some properties of nonplanar graphs, *Compt. rend. Acad. bulg. Sci.* **37** (1984), 1183–1185.
- [3] T. W. Haynes, S. T. Hedetniemi and P. J. Slater, *Fundamentals of domination in graphs*, Monographs and Textbooks in Pure and Applied Mathematics **208**. Marcel Dekker, Inc., New York, 1998.
- [4] T. Honjo, K. Kawarabayashi and A. Nakamoto, Dominating sets in triangulations on surfaces, *J. Graph Theory* **63** (2010), 17–30.
- [5] J. P. Hutchinson, On short noncontractible cycles in embedded graphs, *SIAM. J. Discrete Math.* **1** (1988), 185–192.
- [6] J. P. Hutchinson and G. L. Miller, On deleting vertices to make a graph of positive genus planar, in: H. Wilf (ed.), *Discrete algorithms and complexity (Kyoto, 1986)*, Perspect. Comput. **15**, Academic Press, Boston, MA, 1987, 81–98.
- [7] E. L. C. King and Michael J. Pelsmajer, Dominating sets in plane triangulations, *Discrete Math.* **310** (2010), 2221–2230.
- [8] L. R. Matheson and R. E. Tarjan, Dominating Sets in Planar Graphs, *European J. Combin.* **17**(6) (1996), 565–568.
- [9] B. Mohar and C. Thomassen, *Graphs on Surfaces*, first ed., The Johns Hopkins University Press, Baltimore, MD, 2001.
- [10] J. R. Munkres, *Topology*, second ed., Prentice-Hall, Inc., Upper Saddle River, NJ, 2000.
- [11] M. Plummer and X. Zha, On certain spanning subgraphs of embeddings with applications to domination, *Discrete Math.* **309** (2009), 4784–4792.

Appendix: Figures for cases with $w = 5, 7, 11$




Figure 8: Dominating sets for $w = 5$, with $k = 0$ (left), $k = 1$ (center), and $k = 2$ (right)




Figure 9: Dominating sets for $w = 7$ with $k = 0$ (left) and $k = 1$ (right)




Figure 10: Dominating sets for $w = 7$ with $k = 2$ (left) and $k = 3$ (right)




Figure 11: Dominating sets for $w = 11$ with $k = 0$ (left), $k = 1$ (center), and $k = 2$ (right)




Figure 12: Dominating sets for $w = 11$ with $k = 3$ (left), $k = 4$ (center), and $k = 5$ (right)