

Compatibility of selected herbicides with entomopathogenic fungus *Beauveria bassiana* (Bals.) Vuill

Franci Aco CELAR¹, Katarina KOS²

Received December 22, 2011; accepted January 19, 2012.

Delo je sprejeto 22. decembra 2011, sprejeto 19. januarja 2012.

ABSTRACT

The *in vitro* effect of five commonly used herbicides viz., pyridate, fluzifop-P-butyl, foramsulfuron, tembotrione and S-metolachlor on mycelial growth of entomopathogenic fungus *Beauveria bassiana* (ATCC 74040) was evaluated each at different concentrations: 100, 75, 50, 25, 12.5, 6.25 and 0% of recommended field application rate on PDA agar plates at 15 and 25°C. The herbicides tested were classified in 1-4 scoring categories based on reduction in mycelial growth: 1 = harmless (<25% reduction), 2 = slightly harmful (25-50%), 3 = moderately harmful (51-75%), harmful (>75%) in toxicity tests. All the five herbicides had fungistatic effect to *B. bassiana* at varying intensities depending on their concentrations in medium. The present study showed that *B. bassiana* is very sensitive to the herbicides tested, particularly at recommended as well as lower field dosage. The selected herbicides foramsulfuron, tembotrione and S-metolachlor have strong fungistatic effect on mycelial growth (> 75% inhibition) at 15 °C and concentrations from 50 to 100%. Foramsulfuron has fungicidal effect at 100 % concentration. Foramsulfuron, tembotrione and S-metolachlor were less inhibitory at 25 than at 15 °C, but the temperature had no influence on reduction of mycelial growth at pyridate and fluzifop-P-butyl. Of the herbicides tested, pyridate and fluzifop-P-butyl showed less adverse effects and are probably compatible with *B. bassiana* in the field. However, extensive field studies complemented by parallel laboratory experiments should consider assessing the interaction between selected herbicides and *B. bassiana* isolates to evaluate their ecological impact in cropped environments.

Key words: *Beauveria bassiana*, herbicides, inhibition, mycelial growth, compatibility

IZVLEČEK

KOMPATIBILNOST IZBRANIH HERBICIDOV Z ENTOMOPATOGENO GLIVO *Beauveria bassiana* (Bals.) Vuill

V *in vitro* poskusih smo na PDA agarnih ploščah in temperaturah 15 ter 25 °C preučevali učinek petih pogosto uporabljenih herbicidov, in sicer piridata, fluzifop-P-butila, foramsulfurona, tembotriona in S-metolaklora na rast micelija entomopatogene glive *B. bassiana* (ATCC 74040) pri različnih koncentracijah: 100, 75, 50, 25, 12,5, 6,25 in 0 % priporočenega poljskega odmerka. Glede na inhibicijo rasti micelija smo po toksikoloških testih preučevane herbicide razvrstili v štiri razrede: 1 = neškodljiv (<25 % inhibicija), 2 = malo škodljiv (25 – 50 %), 3 = zmerno škodljiv (51 – 75 %), 4 = škodljiv (> 75 %). Vseh pet herbicidov ima fungistatičen učinek na glivo *B. bassiana*, na obseg pa vpliva njihova koncentracija v gojišču. Raziskava je pokazala, da je gliva *B. bassiana* zelo občutljiva na preizkušane herbicide, posebej pri priporočenih poljskih koncentracijah, pa tudi pri manjših odmerkih. Pri temperaturi 15 °C in koncentracijah od 50 do 100 % imajo herbicidi na podlagi foramsulfurona, tembotriona in S-metolaklora izrazit fungistatičen učinek (> 75% inhibicija), pri 100 % odmerku pa ima foramsulfuron celo fungicidni učinek. Foramsulfuron, tembotrion in S-metolaklor so bili manj inhibitorni pri 25 kot 15 °C. Temperatura ni vplivala na inhibicijo rasti micelija pri piridatu in fluzifop-P-butilu. Od vseh preizkušanih herbicidov sta imela piridat in fluzifop-P-butyl najmanj zaviralnih učinkov in bi jih lahko uporabljali na pridelovalnih površinah skupaj z glivo *B. bassiana*. Poleg laboratorijskih testov s herbicidi bi morali izvajati vzporedne poskuse na pridelovalnih površinah, da bi dejansko izvednotili njihov ekološki vpliv na glivo *B. bassiana*.

Ključne besede: *Beauveria bassiana*, herbicidi, inhibicija, rast micelija, kompatibilnost

¹ izr. prof. dr., Katedra za fitomedicino, kmetijsko tehniko, poljedelstvo, pašništvo in travništvo, Jamnikarjeva 101, SI-1111 Ljubljana, e-pošta: franci.celar@bf.uni-lj.si

² asist., univ. dipl. inž. agr., prav tam

1 INTRODUCTION

Entomopathogenic fungi have an important role in biological control of various harmful insects and mites (Keller, 1991). *Beauveria bassiana* (Bals.) Vuill. is the most studied and well known entomopathogenic fungus and is frequently used in commercially available mycoinsecticides (Inglis *et al.*, 2001).

Numerous organisms, physical and chemical factors of soil ecosystem and agrochemicals that are commonly used in crop production, influence entomopathogenic fungi present in soil. Their diversity and number is also affected by plant species present in crop rotation, cultural practices, intensity of soil use and fertilization with mineral and organic fertilizers (Hummel *et al.*, 2002; Klingen and Haukeland, 2006).

Pesticides are anthropogenic factor with synergistic or antagonistic influence on pests as well as their pathogens (entomopathogenic fungi) and through that on their efficiency (Benz, 1987). Optimally chosen pesticides can minimize their harmful effect on entomopathogenic fungi (Luz *et al.*, 2007; Sterk *et al.*, 1999). The efficacy of entomopathogenic fungus *B. bassiana* incorporated in the soil is affected by regular pesticide usage in agronomical practice, because it often leads to their accumulation in soil. It is important to know the compatibility of pesticides (including herbicides) with entomopathogenic fungi to be able to include mycoinsecticides with *B. bassiana* in integrated crop protection (Ambethgar *et al.*, 2009).

Non-target effects of pesticides on beneficial organisms are gaining their importance in developing active

substances for new insecticides and in re-registration of existing ones in European community. Side effects of pesticides (especially of fungicides) on entomopathogenic fungi and the influence of entomopathogenic fungi on other useful microorganisms are tested from microbiological point of view (Sterk *et al.*, 2003).

Due to the complexity of natural environment, expenses and duration of field experiments it is important to make some preliminary test *in vitro* where it is possible to control all the factors. Despite all that, the obtained results can not be directly used in agronomical practice. *In vivo* field test must follow in order to finally confirm or reject the outcome of laboratory tests (Mietkiewski *et al.*, 1997; Moorhouse *et al.*, 1992).

Data obtained by *in vitro* and some *in vivo* experiments suggest a general sensitivity of entomopathogenic fungi to some herbicides (Ambethgar *et al.*, 2009; Gardner and Storey, 1985; Harrison and Gardner, 1992; Keller, 1986; Mietkiewski *et al.*, 1989; Poprawski and Majchrowicz, 1995; Todorova *et al.*, 1998; Wardle and Parkinson, 1992).

We have observed a strong inhibitory effect of some tested herbicides on mycelium growth of *B. bassiana* in our previous laboratory experiments. Herbicides even had a stronger detrimental effect in comparison to fungicides (Celar *et al.*, 2011). Based on these results we have extended our survey to some other herbicides commonly used in Slovenian crop production and the results of this study are presented in this paper.

2 MATERIALS AND METHODS

Our test method is based on the guidelines for testing side-effects of pesticides on *B. bassiana* (Coremans-Pelseneer, 1994), but small plugs of mycelium were placed on the treated

medium with different herbicide concentrations, instead of spore suspension inoculation. Five herbicides were used in our essay (Table 1).

Table 1: Basic data about herbicides used in laboratory essay.**Preglednica 1:** Osnovni podatki o herbicidih, uporabljenih v laboratorijskem preizkušanju.

HERBICIDE	Active ingredient	a.i. %	Dose rate/ ha	Water per ha (l) (recommended)	FD* ml(g)/l	Manufacturer
Lentagran WP	pyridate	45	2 kg	200-400	2	Belchim Crop Protection
Fusilade Forte	fluazifop-P-butyl	15	2 l	200-300	2	Syngenta
Equip	foramsulfuron	2,25	2,5 l	200-300	2,5	Bayer CS
Laudis	tembotrione	4,4	1,5 l	200-300	1,5	Bayer CS
Dual Gold 960 EC	S-metolchlor	96	1,5 l	200-500	1,5	Syngenta

*100 % field dosage used in essay – herbicide concentration in medium

*100 % poljski odmerek v poskusu - koncentracija herbicida v gojišču

To isolate entomopathogenic fungus *B. bassiana* from a product Naturalis® (INTRACHEM Bio Italia S.p.A.) in pure culture, a standard dilution method on potato dextrose agar (PDA – Merck) medium was used. This mycoinsecticide has permission for use in Slovenia and contains *B. bassiana* isolate ATCC 74040.

Just before solidification of sterile PDA medium, herbicide was added in different concentrations; 100 % of recommended field dosage rate, 75 %, 50%, 25%, 12.5 %, and 6.25 %. Treated medium was then poured in sterile Petri dishes (9 cm diameter), 15 ml in each and cooled. Recommended water consumption for herbicides is between 200 and 500 L/ha, but for our essay common water consumption of 1.000 L/ha was used to prepare agar plates. This means that initial laboratory concentrations in agar plates were 2 to 5-times lower that it would be in actual field application suspension. E.g. dose rate of Lentagran is 2 kg/ha, our initial (100 %) concentration was 0,2 %. Control was sterile PDA medium without herbicides added. A small plug (Ø 5 mm) of *B. bassiana* inoculated in the dark on PDA plates at 25 °C for 14 days, was inverted in the center of prepared PDA plates with herbicide and control. Three repetitions were made for each treatment. Inoculated agar plates were then incubated in dark in growth chambers at 15 and 25 °C with 60 % relative air humidity.

After 7 and 14 days, mycelial growth was measured with image analyzer (Nikon NIS Elements BR 2.30).

The fungus-herbicides compatibility data were analyzed according to IOBC classification scheme (Sterk et al., 2003). The replicated fungus radial growth data were averaged and were expressed as percentage of growth inhibition in comparison to corresponding control following Hokkanen and Kotiluoto (1992).

$$I(\%) = \frac{C - H}{C} \cdot 100$$

Where: I, C, H stand for percentage of growth inhibition, growth of fungus in control and growth of fungus in herbicidal medium, respectively. The herbicides were further classified into the toxicity categories proposed by the IOBC working group: Class 1: harmless (<25% inhibition), Class 2: slightly harmful (25%-50%), Class 3: moderately harmful (51%-75%) and Class 4: harmful (>75%). The effect of the herbicides was scored as fungicidal if growth dropped totally, as otherwise it was taken as fungistatic.

All data were analysed using Student-Newman-Keuls test, $P < 0.05$ (Statgraphics Plus Professional 5.1; StatPoint Technologies, Inc.).

3 RESULTS AND DISCUSSION

Average relative mycelium growth rates of *B. bassiana* on agar plates containing different herbicides in six concentrations are presented in Figure 1 and 2. Herbicides based on foramsulfuron, tembotrione and S-metolachlor at 15°C and 50–100% concentration have a strong fungistatic effect (>75% of growth inhibition). Foramsulfuron at the highest tested concentration (100%) even had a fungicidal effect (the mycelium did not grow). The inhibitory effect of these three

herbicides on mycelium growth is decreasing with the decreasing concentration in agar plates, but there are no statistically significant differences in concentrations higher or equal to 50 %, where the inhibition rate reaches 90 to 100% (Figure 1, Table 2). Even 25% concentration of herbicides results in high growth inhibitory effect (82–90%). Similar, 12.5% concentration inhibits mycelium growth for 71-74%. Even at the lowest tested concentration (6.25%) the

inhibitory effect of these three herbicides is so strong, that they are assigned to class 2 (25–50%, slightly harmful). Other two tested herbicides, i.e. pyridate and fluzafop-P-butyl, had a much lower inhibitory effect on mycelium growth in comparison to previously mentioned three herbicides. Pyridate and fluzafop-P-

butyl were moderately harmful (51–75 % of inhibition, class 3) at higher concentrations (50–100%). However, at lower tested concentrations (6.25–25%) herbicides can be assigned to class 2 (slightly harmful) or class 1 (harmless) based on the inhibition rates, as in the case of fluzafop-P-butyl.




Figure 1: Mean relative mycelium growth of *B. bassiana* at different herbicides and concentrations according to control treatment in % (control treatment 100 %) at 15 °C after 14 days (different lowercase letters show significant differences in mean relative mycelial growth among different concentrations within each herbicide).

Slika 1: Povprečni relativni prirasti micelija glive *B. bassiana* pri različnih herbicidih in koncentracijah v primerjavi s kontrolo izraženi v % (kontrola je 100 %) pri 15 °C po 14 dneh (različne črke pomenijo statistično značilne razlike v priraščanju micelija pri različnih koncentracijah v okviru enega herbicida).




Figure 2: Mean relative mycelium growth of *B. bassiana* at different herbicides and concentrations according to control treatment in % (control treatment 100 %) at 25 °C after 14 days (different lowercase letters show significant differences in mean relative mycelial growth among different concentrations within each herbicide).

Slika 2: Povprečni relativni prirasti micelija glive *B. bassiana* pri različnih herbicidih in koncentracijah v primerjavi s kontrolo izraženi v % (kontrola je 100 %) pri 25 °C po 14 dneh (različne črke pomenijo statistično značilne razlike v priraščanju micelija pri različnih koncentracijah v okviru enega herbicida).

Similar results were obtained at 25°C, where herbicides based on foramsulfuron, tembotrione, and S-

metolachlor had a significant growth-inhibitory effect, especially at 50–100% concentrations (Figure 2, Table

2). All three previously mentioned herbicides are classified as harmless (class 1) at lowest tested concentration based on growth inhibition rates. The mycelium growth inhibition rates at lower tested concentrations (6.25–25%) are much lower at 25°C in comparison to those at 15°C.

Lower fungistatic effect of herbicides at higher temperature could be explained with the better vitality of *B. bassiana* at higher temperatures (its optimum temperatures for development are between 20 and 26°C). Average growth rates of *B. bassiana* at 25°C are up to 5-times higher compared to 15°C. However, this does not explain the results for pyridate and fluzifop-P-butyl, where the temperature had no significant effect on mycelium growth inhibition (Figure 1 and 2).

100% concentration of tested herbicides was calculated based on the 1000 L of water used for treatment of one hectare. It can be seen in Table 1, that the manufacturer recommends 2- to 5-times less water use (200–500 L). This means 2- to 5- times higher herbicide concentration for the treatment. This decision for amount of water used was made in order to make concentrations of individual herbicides more comparable. *B. bassiana* incorporated in the soil is never exposed to the herbicide concentration used for plant treatment. Our highest concentration of herbicide on agar plates (100%), means only 20–50% of the concentration used for plant treatment on the field (depends on the type of herbicide).

Table 2: Percent of mycelium growth inhibition of *B. bassiana* at different herbicides and concentrations according to control treatment at 15 and 25 °C after 14 days and classification in scoring categories based on reduction in mycelial growth.point mutations.

Preglednica 2: Odstotek inhibicije rasti micelija glive *B. bassiana* pri različnih herbicidih in koncentracijah v primerjavi s kontrolnim obravnavanjem pri 15 in 25 °C po 14 dneh ter razdelitev v posamezne razrede glede na obseg inhibicije.

		pyridate	X	fluzifop-p-butyl	X	foramsulfuron	X	tembotrion	X	S-metolachlor	X
100 %	15°C	59,17 d ¹	3	57,60 e*	3	100 e*	4	95,95 e	4	94,09 e	4
	25°C	64,65 d²	3	62,29 e*	3	96,46 e*	4	90,77 de	4	92,65 f	4
75 %	15°C	54,34 d	3	48,75 d	2	96,38 e*	4	96,53 e	4	92,02 de*	4
	25°C	47,97 c	2	55,23 de	3	87,11 de*	4	88,70 d	4	88,82 e*	4
50 %	15°C	50,44 d	3	46,06 d	2	93,76 e	4	91,44 e*	4	90,78 d*	4
	25°C	40,36 b	2	49,43 d	2	73,29 d	3	94,98 e*	4	94,46 g*	4
25 %	15°C	40,00 c	2	18,89 c	1	82,58 d*	4	83,47 d*	4	89,13 d*	4
	25°C	33,23 b	2	22,08 c	1	48,52 c*	2	62,68 c*	3	76,64 d*	4
12,5 %	15°C	24,72 b	1	5,66 b	1	69,07 c*	3	51,06 c*	3	74,25 c*	3
	25°C	31,98 b	2	9,81 b	1	27,79 b*	2	21,46 b*	1	55,20 c*	3
6,25 %	15°C	28,26 b*	2	5,84 b	1	37,17 b	2	29,32 b*	2	47,96 b*	2
	25°C	34,03 b*	2	11,57 b	1	12,55 ab	1	0,00 a*	1	18,66 b*	1
0 %		0,00 a		0,00 a		0,00 a		0,00 a		0,00 a	

Legend: X - scoring categories of inhibition: 1. class: harmless (<25%), 2. class: slightly harmful (25-50%), 3. class: moderately harmful (51-75%), and 4. class: harmful (>75%)

* significant difference of percent of mycelial growth inhibition between two temperatures (15 and 25 °C)

^{1, 2} different lowercase letters show significant differences in mycelial growth inhibition among different concentrations within each herbicide at one temperature

Legenda: X – razred inhibicije: 1. razred: neškodljiv (<25 %); 2. razred: malo škodljiv (25-50 %); 3. razred: zmerno škodljiv (51-75 %) in 4. razred: škodljiv (>75 %)

* statistično značilna razlika v % inhibicije med temperaturama (15 in 25°C)

^{1, 2} različne črke pomenijo statistično značilne razlike v inhibiciji med posameznimi koncentracijami pri enem herbicidu in eni temperaturi

Conditions established in soil after herbicide application (partial binding to clay and hummus particles, microbiological and chemical decomposition, rinsing, dissolving due to precipitation etc.) were simulated in this way. Beside all that, we were interested to see the

effect of small amounts of herbicide residuals on entomopathogenic fungus *B. bassiana*. The later is very important to be able to evaluate the influence of residuals of herbicides used in previous crop on efficacy of mycoinsecticide based on *B. bassiana* used in current

crop. Obtained results have confirmed that our decision was correct, since all of the studied herbicides had a significant fungistatic effect even at low concentrations.

If we take a look at the harmfulness classification (Table 2) of individual treatments we see that S-metolachlor, tembotrione and foramsulfuron are harmful for *B. bassiana* at 50% of field concentration (class 4), S-metolachlor even at 25% concentration. Herbicides pyridate and fluzazifop-P-butyl are moderately harmful (class 3) only at the highest tested concentrations. Laboratory experiments *in vitro* showed higher acceptability of pyridate and fluzazifop-P-butyl usage together with the use of *B. bassiana* in integrated crop production in comparison to S-metolachlor, tembotrione and foramsulfuron, that are inhibitory for mycelium growth even at low concentrations.

After collecting published results of many different experiments on *B. bassiana* and pesticides Klingen and

Haukeland (2006) observed that fungicides have the highest mycelium growth inhibitory effect, while insecticides and herbicides have a fungistatic effect. Detrimental effects of herbicides (especially terrestrial ones) on growth and sporulation of fungus *B. bassiana* was confirmed by many different researchers (Ambethgar, 2009; Gardner and Storey, 1985; Harrison and Gardner, 1992; Mietkiewski *et al.*, 1989; Poprawski in Majchrowicz, 1995; Todorova *et al.*, 1998; Wardle and Parkinson, 1992). In our study, effects of herbicides not commonly tested by other researchers were evaluated. Only Poprawski and Majchrowicz (1995) also observed fungicidal effect of metolachlor on *B. bassiana*. Similar herbicide from our study, S-metolachlor, also inhibited mycelium growth (93%). We have to stress, that it is a similar kind of herbicide, not exactly the same one and that concentrations used in our study were lower compared to those by Poprawski and Majchrowicz (1995).

4 CONCLUSIONS

Based on the results of this essay, conclusions that entomopathogenic fungus *B. bassiana* is very sensitive to the herbicides tested can be made. Particularly at recommended, but as well as at lower field dosage, herbicides have strong fungistatic or even fungicidal effect. Preliminary tests on agar plates (*in vitro*) in laboratory conditions have limited application value and

therefore can not be directly transformed in field practice. These results must be examined also in field trials (*in situ*). Besides of active ingredients the formulation of herbicide can also have fungistatic or fungicidal effect (Morjan *et al.*, 2002) meaning that results can not be generalized for all the products containing the same active ingredient.

5 ACKNOWLEDGEMENT

This work was founded by Phytosanitary Administration of the Republic of Slovenia, Ministry of Agriculture, Forestry and Food of R Slovenia. We would like to thank also to manufacturers Karsia d.o.o.,

Bayer CropScience Slovenija and Syngenta Agro Slovenia for free samples of herbicides used in our essay.

6 REFERENCES

- Ambethgar, V., Swamiappan, M., Rabindra, R. J., Rabindran, R. 2009. Influence of some herbicides on *in vitro* vegetative growth of *Beauveria bassiana* (Balsamo) Vuillemin. Resistant Pest Management Newsletter 19: 13-16.
- Benz, G. 1987. Environment. V: Epizootiology of Insect Diseases (ur. Fuxa, R. in Tanada, J.). Wiley, New York: 177-214.
- Celar, F. A., Sekne, Š., Mesec, D., Kos, K. 2011. Učinek herbicidov in fungicidov na rast micelija entomopatogene glive *Beauveria bassiana* (Bals.) Vuill. = Effect of selected herbicides and fungicides on mycelial growth of entomopathogenic fungus *Beauveria bassiana* (Bals.) Vuill. V: Maček, J., Trdan, S. (ur.). Zbornik predavanj in referatov 10. slovenskega posvetovanja o varstvu rastlin, Podčetrtek, 1.-2. marec 2011. Ljubljana: Društvo za varstvo rastlin Slovenije = Plant Protection Society of Slovenia: 171-175.
- Coremans-Pelseneer, J., 1994. Laboratory tests on the entomopathogenic fungus *Beauveria*. IOBC/WPRS Bull. 17,10: 147-154.
- Gardner, W. A., Storey, G. K. 1985. Sensitivity of *Beauveria bassiana* to selected herbicides. J. Econ. Entomol. 78: 1275-1279.
- Harrison, R.D., Gardner, W.A. 1992. Fungistasis of *Beauveria bassiana* by selected herbicides in soil. J. Entomol. 27: 233-238.
- Hokkanen, H.M.T and R. Kotiluoto. 1992. Bioassay of the side effects of pesticides on *Beauveria bassiana* and

- Metarhizium anisopliae: standardized sequential testing procedure. World Pesticides Research Science Bulletin, 11: 148-151.
- Hummel, R.L., Walgenbach, J.F., Barbercheck, M.E., Kennedy, G.G., Hoyt, G.D., Arellano, C. 2002. Effects of production practices on soil-borne entomopathogens in western North Carolina vegetable systems. Environmental Entomology 31: 84-91.
- Inglis, G.D., Goettel, M.S., Butt, T.M., Strasser, H. 2001. Use of Hyphomycetous Fungi for Managing Insect Pests. V: Fungi as Biocontrol Agents – Progress, Problems and Potential (ur. Butt, T.M., Jackson, C.W., Magan, N.). CABI Publishing, Walingford: 23-69.
- Keller, S. 1986. Investigations on the effect of herbicides on aphid pathogenic Entomophthoraceae. V: Ecology of Aphidophaga (ur. Hodek, I.). Academia, Prague and Dordrecht: 493-497.
- Keller, S. 1991. Les maladies fongiques des ravageur et leur importance pratique. Revue Suisse de viticulture, arboriculture, horticulture 23: 299-310.
- Klingen, I., Haukeland, S. 2006. The soil as a reservoir for natural enemies of pest insects and mites with emphasis on fungi and nematodes. V: An Ecological and Societal Approach to Biological Control (ur. Eilenberg, J., Hokkanen, H.M.T.). Springer, Netherlands: 145–211.
- Luz, C., Bastos, N., Nunes, R. 2007. In vitro susceptibility to fungicides by invertebrate-pathogenic and saprobic fungi. Mycopathologia 164: 39-47.
- Mietkiewski, R.T, Pell, J.K, Clark, S.J. 1997. Influence of pesticide use on the natural occurrence of entomopathogenic fungi in arable soils in the UK: field and laboratory comparisons. Biocontr. Sci. Technol. 7: 565–575.
- Mietkiewski, R., Sapiecha, A., Mietkiewska, Z. 1989. Growth of entomopathogenic fungi on a medium containing herbicides used in orcharding. Acta Mycology 25: 35-50.
- Moorhouse, E.R., Gillespie, A.T., Sellers, E.K., Chamley, A.K. 1992. Influence of fungicides and insecticides on the entomogenous fungus *Metarhizium anisopliae*, a pathogen of the vine weevil, *Othiorhynchus sulcatus*. Biocontrol Science and Technology 2: 49-58.
- Morjan, W.E., Pedigo, L.P., Levis, L.C. 2002. Fungicidal effects glyphosate and glyphosate formulations on four species of entomological fungi. Environmental Entomology 31: 1206-1212.
- Poprawski, T.J., Majchrowicz, I. 1995. Effects of herbicides on in vitro vegetative growth and sporulation of entomopathogenic fungi. Crop Protection 14: 81-87.
- Sterk, G., Hassan, S.A., Baillo, M., Bakker, F., Bigler, F., Blumel, S., Bogenschutz, H., Boller, E., Bromand, B., Brun, J., Calis, J.N.M., Coremans-Pelseneer, J., Duso, C., Garrido, A., Grove, A., Heimbach, U., Hokkanen, H., Jacas, J., Lewis, G., Moreth, L., Polgar, L., Roversti, L., Samsøe-Peterson, L., Sauphanor, B., Schaub, L., Staubli, A., Tuset, J.J., Vainio, A., Van de Veire, M., Viggiani, G., Vinuela, E., Vogt, H. 1999. Results of the seventh joint pesticide testing programme carried out by the IOBC/WPRS-Working Group 'Pesticides and Beneficial Organisms'. BioControl 44: 99-117.
- Sterk, G., Heuts, F., Merck, N., Bock, J. 2003. Sensitivity of non-target arthropods and beneficial fungal species to chemical and biological plant protection products: Results of laboratory and semi-field trials. V: 1st International Symposium on Biological Control of Arthropods, Honolulu, 14-18 januar 2002, USDA Forest Service, Forest Health Technology Enterprise Team: 306-313.
- Todorova, S.I., Coderre, D., Duchesne, R.M. and Côté, J.C. 1998. Compatibility of *Beauveria bassiana* with selected fungicides and herbicides. Biological Control 27: 427-433.
- Wardle, D.A., Parkinson, D. 1992. The influence of the herbicide glyphosate on interspecific interactions between four soil fungal species. Mycol. Res. 96: 180-186.