

DOI: 10.2478/v10014-009-0002-4

**Agrovoc descriptors:** *Juglans regia*, walnuts, fruits, phenolic compounds, aromatic compounds, ethanol, methanol, alcohols, tannins, organic acids, catechin

**Agris category code:** F60

## Extraction of phenolic compounds from green walnut fruits in different solvents

Jerneja JAKOPIČ<sup>1</sup>, Robert VEBERIČ<sup>2</sup>, Franci ŠTAMPAR<sup>3</sup>

Received September 11, 2006; accepted March 20, 2009  
Prispelo 11. septembra 2006; sprejeto 20. marca 2009

### ABSTRACT

The extractive efficiency of phenolic compounds from plant material is greatly depended on the solvent. In our research, methanol and ethanol were used for taking out the phenols from green walnut fruits. The total phenolics and some individual phenolics, such as gallic, chlorogenic, ellagic, sinapic and protocatechuic acid, (+)-catechin and juglone were detected. Total phenolic contents were determined spectrophotometric but individual phenols using HPLC. Amounts of total phenolics were higher when for extraction methanol was used compare to ethanol, in both cultivars, Elit and Franquette. Similar effect of the solvents was observed in the case some individual phenols, such as juglone, (+)-catechin, gallic, protocatechuic and chlorogenic acid. On the contrary, for ellagic and sinapic acid the extraction was better when ethanol was used compare to methanol.

**Key words:** *Juglans regia* L., ethanol, methanol, juglone, gallic acid, catechin

### EKSTRAKCIJA FENOLNIH SNOVI IZ ZELENIH PLODOV OREHA Z RAZLIČNIMI TOPILI

#### IZVLEČEK

Učinkovitost ekstrakcije fenolnih snovi iz rastlinskega tkiva je v veliki meri odvisna od topila. V našem poskusu smo primerjali ekstrakcijo fenolov iz zelenih plodov orehov z etanolom in metanolom. Določali smo skupne in nekatere posamezne fenolne spojine, kot so galna, klorogenska, elagna, sinapinska in protokatehulna kislina, ter (+)-katehin in juglon. Skupne fenole smo določili spektrofotometrično, posamezne pa s pomočjo HPLC sistema. Vsebnost skupnih fenolov je bila pri obeh sortah, 'Elit' in 'Franquette', večja pri ekstrakciji v metanolu kot v etanolu. Podobne rezultate smo dobili pri nekaterih posameznih fenolih, kot so juglon, (+)-katehin, galna, protokatehulna in klorogenska kislina. Nasprotno je bila ekstrakcija pri elagni in sinapinski kislini bolj uspešna v etanolu kot v metanolu.

**Ključne besede:** *Juglans regia* L., etanol, metanol, juglon, galna kislina, katehin

## 1 INTRODUCTION

Walnut (*Juglans regia* L.) naturally occurs in Slovenian forests. For many years the trees are also grown in orchards because of the delicious fruits, which can be eaten raw and are excellent for dessert as well as in baking and confectionery (Prasad, 2003). Before the endocarp hardens, the green fruits including green husks can be pickled in vinegar or sliced and steeped into

alcohol. The latter case of use is traditional in Slovenia as well as in Italy in preparation of a walnut liqueur (Štampar et al., 2006; Alamprese and Pompei, 2005).

Walnut fruits are rich in phenolic compounds (Prasad, 2003). Their contents depend on many environmental conditions, as well as genotype of different cultivars

<sup>1</sup> University of Ljubljana, Biotechnical Faculty, Agronomy Department, Chair for Fruit Growing, Jamnikarjeva 101, SI-1000 Ljubljana, Slovenia; e-mail: jerneja.jakopic@bf.uni-lj.si

<sup>2</sup> University of Ljubljana, Biotechnical Faculty, Agronomy Department, Chair for Fruit Growing, Jamnikarjeva 101, SI-1000 Ljubljana, Slovenia

<sup>3</sup> University of Ljubljana, Biotechnical Faculty, Agronomy Department, Chair for Fruit Growing, Jamnikarjeva 101, SI-1000 Ljubljana, Slovenia

(Colarič et al., 2005; Solar et al., 2006). Furthermore, the presence of phenols changes during the season (Solar et al., 2006). Phenolics are involved in growth and reproduction and provide plants with resistance to pathogens and predators (Bravo, 1998).

For extraction of polyphenols from plant matter different solvent systems have been used (Chavan et al., 2001) and their efficiency varies. Extraction yield depends on the solvent and method of extraction (Goli et al., 2004). Commonly used solvents for extracting various substances from plant material are water,

aqueous mixtures of ethanol, methanol and acetone (Sun and Ho, 2005).

We have analyzed the contents of the total and of some of the individual phenolic compounds in green walnut fruits, extracted with methanol or ethanol. Two cultivars, Elit and Franquette were involved in the research. In scientific researches, extraction of the phenolics is often performed in methanol, but when a walnut liqueur is made, green walnuts are steeped in ethanol. The aim of our study was to ascertain the influence of the solvents on the content of total as well as on some individual phenolic compounds.

## 2 MATERIAL AND METHODS

### 2.1 Plant material

Material for analyses was taken from two walnut cultivars, Elit and Franquette in an experimental orchard of the Biotechnical Faculty, located in Maribor (Slovenia). In the end of June, green walnut fruits were picked from three trees of each cultivar.

Two fresh fruits were sliced and ground to a fine powder and 100 mg of the sample was measured into a test tube. The sample was extracted with 5 ml methanol or ethanol in ultrasonic bath for 45 minutes. Then the samples were centrifuged for 7 minutes at 4200 rpm. The supernatant was filtered through polyamide filter Chromafil AO-45/25, transferred into vial prior analyses.

### 2.2 Total phenols content

Total phenolic content was assessed by using the Folin-Ciocalteu phenol reagent method (Singleton and Rossi, 1965). To 100  $\mu$ l of the samples, 5 ml of bidistilled water and 500  $\mu$ l of Folin-Ciocalteu reagent were added. After 30 sec to 8 min, 1.5 ml of sodium carbonate (20% w/v) was added. The extracts were stand for 30 min at 40°C. On the basis of measured absorbances at 765 nm, determination from calibration curve and considering dilutions, the total phenolic content was expressed as gallic acid equivalents (GAE) in milligrams per gram fresh walnut fruits.

### 2.3 Individual phenols content

The content of individual phenolics in extracts was analyzed on the Thermo Finningan Surveyor HPLC system equipped with photodiode array detector (PDA detector). A Chromsep HPLC column SS (250 x 4.6 mm, Hypersil 5 ODS) protected

with a Chromsep guard column SS (10 x 3 mm) was used. The system was controlled using the ChromQuest™ 4.0 Chromatography workstation software system. The chromatographic conditions (mobile phase, gradient program, temperature of column) were similar to those described by Schieber et al. (2001).

Chromatograms were observed at 280 nm. Identification of individual phenolic was qualitatively achieved using method of external standards and quantitatively comparing peak area on chromatograms of samples with those of diluted standard solutions.

### 2.4 Chemicals

For quantification of phenolics the following standards were used: juglone, gallic and protocatechuic acid from Merck, chlorogenic and ellagic acid from Sigma, (+)-catechin from Roth and sinapic acid from Fluka.

Following chemicals were used for the determination of total phenolics: Folic-Ciocalteu's phenol reagent and gallic acid from Sigma and sodium carbonate from Riedel-de Haën.

Acetonitrile, methanol and acetic acid were HPLC grade from Merck. The water used was bidistilled, purified with a Milli-Q water purification system (Millipore, Badford, USA).

### 2.5 Statistical analyses

Data were evaluated by one-way analysis of variance (ANOVA) using Statgraphics Plus 4.0 (Manugistics, Inc.; Rockville, USA) software. T-test was performed to determine significant differences at  $P < 0.05$ .

## 3 RESULTS AND DISSCUSION

Phenolic compounds were extracted from green walnut fruits, cultivars 'Elit' and 'Franquette'. Total phenolic content as well contents of seven individual phenolics, such as gallic, chlorogenic, ellagic, sinapic and protocatechuic acid, (+)-catechin and juglone, were detected. As extraction solvent, methanol or ethanol were compared.

Colarič et al. (2005) ascertained that green unripe walnuts are rich in individual phenolic compounds. The total phenolics content of green walnut fruits extracted in two solvents are presented in Table 1 for cultivars Elit and Franquette. In the case of ethanol, the total phenolic content ranged from 126.2 mg GAE per g in cultivar Elit to 135.3 mg GAE per g in cultivar Franquette. In the case of methanol more phenolic

compounds were extracted in both cultivars. There was significant difference between phenolic contents of the extracts using two mentioned solvents in both cultivars. Also other authors have established that the phenolics content extracts are strongly dependent on the type of the solvent as well as on the different concentrations of solvent (Turkmen et al., 2006; Yilmaz and Toledo, 2006). The extraction yield and extraction efficiency of caffeine and major catechins of green tea were higher with pure methanol comparing to pure ethanol (Perva-Uzunalić et al., 2006). Therefore, methanol proved to have a bit better characteristics as a solvent for polyphenols and anthocyanins from black currant and grape byproducts than ethanol (Lapornik et al., 2005). On the other hand, Wang and Halliwell (2001) reported that aqueous ethanol was superior to aqueous methanol and acetone for extraction of the flavonoids from tea. In extracting phenolic compounds from peanut skin, ethanol and methanol were more effective than water, with ethanol being the most efficient extraction solvent (Yu et al., 2005). Meanwhile, the methanol was the solvent with best results for phenols from pine sawdust, while in almond hulls ethanol was the best extraction solvent (Pinelo et al., 2004). Jung et al. (2006) also compared the influence of different solvents and they found out that the ethanol extracts contained higher amounts of total phenolics and flavonoids than water and methanol extracts from wild ginseng leaves.

**Table 1:** The total phenols (mg GAE/g FW) in green walnut fruits, extracted with methanol or ethanol in cultivars Elit in Franquette. Average means and standard errors are presented. *P*-values in last column of the table mark statistically significant differences between the solvents.

	Methanol	Ethanol	<i>P</i> -value
Franquette	161.07 ± 7.28	135.27 ± 2.42	0.0041
Elit	148.98 ± 4.74	126.20 ± 5.71	0.0119


In spite of the fact, that higher amounts of total phenolics were extracted in methanol comparing to ethanol, this is not the case for all the individual phenols. The highest values of the analyzed phenols

achieved juglone. Its extraction was more effective with methanol than with ethanol. Similar effects of the solvents were observed in the case of (+)-catechin, gallic, protocatechuic and chlorogenic acid (Fig. 1). Kallithara et al. (1995) also indicated that the methanol extraction is best for (+)-catechin, (-)-epicatechin and epigallocatechin, whereas the largest amount of gallic acid were yielded by 75 % of ethanol. In our research 96% ethanol was used, which has proved to be less effective solvent for gallic acid than methanol.

For other analyzed phenols, such as ellagic and sinapic acid, higher contents were gained with ethanol than with methanol. On the Fig.2 their contents are shown for cultivars Elit and Franquette. Also the individual anthocyanin extraction from grape and black currant marc depended on the use of solvent (Lapornik et al., 2005). Rødtjer et al. (2006) reported that quantification of the total amount of phenolics in the extracts showed that 70% solvent-water mixtures extracted the phenolics more efficiently and contained more complex mixtures of phenolic compounds than the pure solvent extracts did.

Many authors established that the extraction yield of phenols is greatly depending on the solvent polarity (Turkmen et al., 2006; Lapornik et al., 2005). There are some studies, where the influences of the solvent concentration are being studied. Thus, Yilmaz and Toledo (2006) found out that aqueous solutions of ethanol, methanol or acetone were better than a single-compound solvent system for the extraction of the total phenols from Muscadine seed power.

We can conclude that the efficiency of the phenolics extraction depends on the type of the solvent as well on the phenol, which is being isolated. For total phenolics extraction from green walnut fruits methanol was more efficient compare to ethanol. Similar was for some individual phenols, such as juglone, (+)-catechin, gallic, protocatechuic and chlorogenic acid. On the contrary, for ellagic and sinapic acid extraction was better when using ethanol.



**Figure 1:** The individual phenols (mg/g FW) in green walnut fruits, extracted with methanol or ethanol in cultivars Elit in Franquette. Average means and standard errors are presented. Asterisk marks statistically significant differences between the solvents at  $P$ -value < 0.05.

#### 4 REFERENCES

- Alamprese, C. and Pompei, C. (2005). Influence of processing variables on some characteristics of nocino liqueur. *Food Chemistry* 92:203-209.
- Bravo, L. (1998). Polyphenols: Chemistry, dietary sources, metabolism, and nutritional significance. *Nutrition Reviews* 56:317-333.
- Chavan, U. D., Shahidi, F. and Naczk, M. 2001. Extraction of condensed tannins from beach pea (*Lathyrus maritimus* L.) as affected by different solvents. *Food Chemistry* 75: 509–512.
- Colarič, M., Veberič, R., Solar, A., Hudina, M. and Stampar, F. (2005). Phenolic acids, syringaldehyde, and juglone in fruits of different cultivars of *Juglans regia* L. *Journal of Agricultural and Food Chemistry* 53:6390-6396.
- Goli, A. H., Barzegar, M. and Sahari, M. A. 2004. Antioxidant activity and total phenolic compounds of pistachio (*Pistachia vera*) hull extracts. *Food Chemistry* 92:521–525.

- Jung, C.H., Seog, H.M., Choi, I.W., Park, M.W. and Cho, H.Y. 2006. Antioxidant properties of various solvent extracts from wild ginseng leaves. *LWT* 39:266–274.
- Kallithara, S., Garcaviaguera, C., Bridle, P. and Bakker, J. 1995. Survey of solvents for the extraction of rape seed phenolics. *Phytochemical Analysis* 6:265-267.
- Lapornik, B., Prošek, M and Wondra, A.G. 2005. Comparison of extracts prepared from plant by-products using different solvents and extraction time. *Journal of Food Engineering* 71:214-222.
- Perva-Uzunalić, A., Škerget, M., Knez, Ž., Weinreich, B., Otto, F. and Grüner S. 2006. Extraction of active ingredients from green tea (*Camellia sinensis*): Extraction efficiency of major catechins and caffeine. *Food Chemistry* 96:597–605
- Pinelo, M., Rubilar, M., Sineiro, J. and Núñez, M.J. 2004. Extraction of antioxidant phenolics from almond hulls (*Prunus amygdalus*) and pine sawdust (*Pinus pinaster*). *Food Chemistry* 85:267-273.
- Prasad, R.B.N. 2003. Walnuts and pecans. p. 6071-6079. In: B. Caballero, L.C. Trugo and P.M. Finglas (eds.), *Encyclopaedia of food sciences and nutrition*, Academic Press, London.
- Rødtjer, A., Skibsted, L.H. and Andersen, M.L. 2006. Antioxidative and prooxidative effect of extracts made from cherry liqueur pomace. *Food Chemistry* 99:6-14.
- Schieber, A., Keller, P. and Carle, R. (2001). Determination of phenolic acids and flavonoids of apple and pear by high-performance liquid chromatography. *Journal of Chromatography A* 910:265-273.
- Singleton, V.L. and Rossi, J.A. Jr. (1965). Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents, *American Journal of Enology and Viticulture*. 16:144-158.
- Solar, A., Colarič, M., Usenik, V. and Štampar, F. 2006. Seasonal variations of selected flavonoids, phenolic acids and quinines in annual shoots of common walnut (*Juglans regia* L.). *Plant Science* 170:453-461.
- Štampar, F., Solar, A., Hudina, M., Veberič, R. and Colarič, M. (2006). Traditional walnut liqueur – cocktail of phenolics. *Food Chemistry* 95:627-631.
- Sun, T. and Ho, C. 2005. Antioxidant activities of buckwheat extracts. *Food Chemistry* 90:743–749.
- Turkmen, N., Sari, F. and Velioglu, Y.S. 2006. Effect of extraction solvents on concentration and antioxidant activity of black and black mate polyphenols determined by ferrous tartrate and Folin-Ciocalteu methods. *Food Chemistry* 99:838-841.
- Wang, H. and Helliwell, K. 2001. Determination of flavonols in green and black tea leaves and green tea infusions by high-performance liquid chromatography. *Food Research International* 34:223–227.
- Yilmaz, Y. and Toledo, R.T. 2006. Oxygen radical absorbance capacities of grape/wine industry byproducts and effect of solvent type on extraction of grape seed polyphenols. *Journal of Food Composition and Analysis* 19:41–48.
- Yu, J., Ahmedna, M. and Goktepe, I. 2005. Effects of processing methods and extraction solvents on concentration and antioxidant activity of peanut skin phenolics. *Food Chemistry* 90:199-206.