

Tight relative t -designs on two shells in hypercubes, and Hahn and Hermite polynomials*

Eiichi Bannai

*Faculty of Mathematics, Kyushu University (emeritus), Japan, and
National Center for Theoretical Sciences, National Taiwan University, Taiwan*

Etsuko Bannai

National Center for Theoretical Sciences, National Taiwan University, Taiwan

Hajime Tanaka[†] 

*Research Center for Pure and Applied Mathematics, Graduate School of Information
Sciences, Tohoku University, Sendai 980-8579, Japan*

Yan Zhu[‡] 

*College of Science, University of Shanghai for Science and Technology,
Shanghai 200093, China*

Received 3 June 2020, accepted 24 May 2021, published online 14 April 2022

Abstract

Relative t -designs in the n -dimensional hypercube \mathcal{Q}_n are equivalent to weighted regular t -wise balanced designs, which generalize combinatorial t - (n, k, λ) designs by allowing multiple block sizes as well as weights. Partly motivated by the recent study on tight Euclidean t -designs on two concentric spheres, in this paper we discuss tight relative t -designs in \mathcal{Q}_n supported on two shells. We show under a mild condition that such a relative t -design induces the structure of a coherent configuration with two fibers. Moreover, from this structure we deduce that a polynomial from the family of the Hahn hypergeometric orthogonal polynomials must have only integral simple zeros. The Terwilliger algebra is the main tool to establish these results. By explicitly evaluating the behavior of the zeros of the Hahn polynomials when they degenerate to the Hermite polynomials under an appropriate limit

*This work was also partially supported by the Research Institute for Mathematical Sciences at Kyoto University.

[†]Corresponding author. Hajime Tanaka was supported by JSPS KAKENHI Grant Numbers JP25400034 and JP17K05156.

[‡]Yan Zhu was supported by NSFC Grant No. 11801353.

process, we prove a theorem which gives a partial evidence that the non-trivial tight relative t -designs in \mathcal{Q}_n supported on two shells are rare for large t .

Keywords: Relative t -design, association scheme, coherent configuration, Terwilliger algebra, Hahn polynomial, Hermite polynomial.

Math. Subj. Class. (2020): 05B30, 05E30, 33C45

1 Introduction

This paper is a contribution to the study of *relative t -designs* in Q -polynomial association schemes. In the Delsarte theory [16], the concept of t -designs is introduced for arbitrary Q -polynomial association schemes. For the Johnson scheme $J(n, k)$, the t -designs in the sense of Delsarte are shown to be the same thing as the combinatorial t - (n, k, λ) designs. There are similar interpretations of t -designs in some other important families of Q -polynomial association schemes [16, 17, 19, 34, 41]. The concept of relative t -designs is also due to Delsarte [18], and is a relaxation of that of t -designs. Relative t -designs can again be interpreted in several cases, including $J(n, k)$. For the n -dimensional hypercube \mathcal{Q}_n (or the binary Hamming scheme $H(n, 2)$) which will be our central focus in this paper, these are equivalent to the weighted regular t -wise balanced designs, which generalize the combinatorial t - (n, k, λ) designs by allowing multiple block sizes as well as weights.

The Delsarte theory has a counterpart for the unit sphere \mathbb{S}^{n-1} in \mathbb{R}^n , established by Delsarte, Goethals, and Seidel [20]. The t -designs in \mathbb{S}^{n-1} are commonly called the *spherical t -designs*, and are essentially the equally-weighted *cubature formulas* of degree t for the spherical integration, a concept studied extensively in numerical analysis. Spherical t -designs were later generalized to *Euclidean t -designs* by Neumaier and Seidel [35] (cf. [21]). Euclidean t -designs are in general supported on multiple concentric spheres in \mathbb{R}^n , and it follows that we may think of them as the natural counterpart of relative t -designs in \mathbb{R}^n . This point of view was discussed in detail by Bannai and Bannai [3]. See also [7, 8]. The success and the depth of the theory of Euclidean t -designs (cf. [38]) has been one driving force for the recent research activity on relative t -designs in Q -polynomial association schemes; see, e.g., [3, 5, 6, 7, 8, 9, 11, 32, 51, 53, 54].

A relative t -design in a Q -polynomial association scheme (X, \mathcal{R}) is often defined as a certain *weighted subset* of the vertex set X , i.e., a pair (Y, ω) of a subset Y of X and a function $\omega: Y \rightarrow (0, \infty)$. We are given in advance a ‘base vertex’ $x \in X$, and (Y, ω) gives a ‘degree- t approximation’ of the shells (or spheres or subconstituents) with respect to x on which Y is supported. See Sections 2 and 3 for formal definitions. Bannai and Bannai [3] proved a Fisher-type lower bound on $|Y|$, and we call (Y, ω) *tight* if it attains this bound. We may remark that t must be even in this case. In this paper, we continue the study (cf. [5, 9, 32, 51, 53]) of tight relative t -designs in the hypercubes \mathcal{Q}_n , which are one of the most important families of Q -polynomial association schemes. The Delsarte theory directly applies to the tight relative t -designs in \mathcal{Q}_n supported on one shell, say, the k^{th} shell, as these are equivalent to the tight combinatorial t - (n, k, λ) designs. (We note that the k^{th} shell induces $J(n, k)$.) Our aim is to extend this structure theory to those supported on two shells. We may view the results of this paper roughly as counterparts to (part of) the

results by Bannai and Bannai [2, 4] on tight Euclidean t -designs on two concentric spheres.

Let $t = 2e$ be even. In Theorem 5.3, which is our first main result, we show under a mild condition that a tight relative $2e$ -design in \mathcal{Q}_n supported on two shells induces the structure of a coherent configuration with two fibers. Moreover, from this structure we deduce that a certain polynomial of degree e , known as a *Hahn polynomial*, must have only integral simple zeros. We note that the case $e = 1$ was handled previously by Bannai, Bannai, and Bannai [5]. The Hahn polynomials are a family of hypergeometric orthogonal polynomials in the Askey scheme [31, Section 1.5], and that their zeros are integral provides quite a strong necessary condition on the existence of such relative $2e$ -designs. The corresponding necessary condition for the tight combinatorial $2e$ - (n, k, λ) designs from the Delsarte theory was used successfully by Bannai [1]; that is to say, he showed that, for each given integer $e \geq 5$, there exist only finitely many non-trivial tight $2e$ - (n, k, λ) designs, where n and k (and thus λ) vary. See also [22, 36, 52]. We extend Bannai's method to prove our second main result, Theorem 7.1, which presents a version of his theorem for our case.

The sections other than Sections 5 and 7 are organized as follows. We collect the necessary background material in Sections 2 and 3. Section 3 also includes a few general results on relative t -designs in Q -polynomial association schemes. As in [6, 44], our main tool in the analysis of relative t -designs is the *Terwilliger algebra* [46, 47, 48], which is a non-commutative semisimple \mathbb{C} -algebra containing the adjacency algebra. Section 4 is devoted to detailed descriptions of the Terwilliger algebra of \mathcal{Q}_n . It is well known (cf. [30, 31]) that the Hahn polynomials $({}_3F_2)$ degenerate to the Hermite polynomials $({}_2F_0)$ by an appropriate limit process, and a key in Bannai's method above was to evaluate precisely the behavior of the zeros of the Hahn polynomials in this process. In Section 6, we revisit this part of the method in a form suited to our purpose. Our account will also be simpler than that in [1]. In Appendix, we provide a proof of a number-theoretic result (Proposition 7.2) which is a variation of a result of Schur [40, Satz I].

2 Coherent configurations and association schemes

We begin by recalling the concept of coherent configurations.

Definition 2.1. The pair (X, \mathcal{R}) of a finite set X and a set \mathcal{R} of non-empty subsets of X^2 is called a *coherent configuration* on X if it satisfies the following (C1) – (C4):

(C1) \mathcal{R} is a partition of X^2 .

(C2) There is a subset \mathcal{R}_0 of \mathcal{R} such that

$$\bigsqcup_{R \in \mathcal{R}_0} R = \{(x, x) : x \in X\}.$$

(C3) \mathcal{R} is invariant under the transposition $\tau : (x, y) \mapsto (y, x)$ ($(x, y) \in X^2$), i.e., $R^\tau \in \mathcal{R}$ for all $R \in \mathcal{R}$.

(C4) For all $R, S, T \in \mathcal{R}$ and $(x, y) \in T$, the number

$$p_{R,S}^T := |\{z \in X : (x, z) \in R, (z, y) \in S\}|$$

is independent of the choice of $(x, y) \in T$.

Moreover, a coherent configuration (X, \mathcal{R}) on X is called *homogeneous* if $|\mathcal{R}_0| = 1$, and an *association scheme* if $R^T = R$ for all $R \in \mathcal{R}$.

Remark 2.2. Suppose that a finite group \mathfrak{G} acts on X , and let \mathcal{R} be the set of the orbitals of \mathfrak{G} , that is to say, the orbits of \mathfrak{G} in its natural action on X^2 . Then (X, \mathcal{R}) is a coherent configuration. Moreover, (X, \mathcal{R}) is homogeneous (resp. an association scheme) if and only if the action of \mathfrak{G} on X is transitive (resp. generously transitive, i.e., for any $x, y \in X$ we have $(x^g, y^g) = (y, x)$ for some $g \in \mathfrak{G}$).

Let (X, \mathcal{R}) be a coherent configuration as above. For every $R \in \mathcal{R}_0$, let Φ_R be the subset of X such that $R = \{(x, x) : x \in \Phi_R\}$. Then we have

$$\bigsqcup_{R \in \mathcal{R}_0} \Phi_R = X.$$

We call the Φ_R ($R \in \mathcal{R}_0$) the *fibers* of (X, \mathcal{R}) . By setting in (C4) either $R \in \mathcal{R}_0$ and $S = T$, or $S \in \mathcal{R}_0$ and $R = T$, it follows that for every $T \in \mathcal{R}$, we have $T \subset \Phi_R \times \Phi_S$ for some $R, S \in \mathcal{R}_0$. In particular, (X, \mathcal{R}) is homogeneous whenever it is an association scheme. Let

$$\gamma_{R,S} = |\{T \in \mathcal{R} : T \subset \Phi_R \times \Phi_S\}| \quad (R, S \in \mathcal{R}_0).$$

The matrix

$$[\gamma_{R,S}]_{R,S \in \mathcal{R}_0},$$

which is symmetric by (C3), is called the *type* of (X, \mathcal{R}) .

Let $M_X(\mathbb{C})$ be the \mathbb{C} -algebra of all complex matrices with rows and columns indexed by X , and let $V = \mathbb{C}^X$ be the \mathbb{C} -vector space of complex column vectors with coordinates indexed by X . We endow V with the Hermitian inner product

$$\langle u, v \rangle = v^\dagger u \quad (u, v \in V),$$

where \dagger denotes adjoint. For every $R \in \mathcal{R}$, let $A_R \in M_X(\mathbb{C})$ be the adjacency matrix of the graph (X, R) (directed, in general), i.e.,

$$(A_R)_{x,y} = \begin{cases} 1 & \text{if } (x, y) \in R, \\ 0 & \text{otherwise,} \end{cases} \quad (x, y \in X).$$

Then (C1) – (C4) above are rephrased as follows:

$$(A1) \quad \sum_{R \in \mathcal{R}} A_R = J \text{ (the all-ones matrix).}$$

$$(A2) \quad \sum_{R \in \mathcal{R}_0} A_R = I \text{ (the identity matrix).}$$

$$(A3) \quad (A_R)^\dagger \in \{A_S : S \in \mathcal{R}\} \quad (R \in \mathcal{R}).$$

$$(A4) \quad A_R A_S = \sum_{T \in \mathcal{R}} p_{R,S}^T A_T \quad (R, S \in \mathcal{R}).$$

Let

$$\mathbf{A} = \text{span}\{A_R : R \in \mathcal{R}\}.$$

Then from (A2) and (A4) it follows that \mathbf{A} is a subalgebra of $M_X(\mathbb{C})$, called the *adjacency algebra* of (X, \mathcal{R}) . We note that \mathbf{A} is semisimple as it is closed under \dagger by virtue of (A3). By (A1), \mathbf{A} is also closed under entrywise (or *Hadamard* or *Schur*) multiplication, which we denote by \circ . The A_R are the (central) primitive idempotents of \mathbf{A} with respect to \circ , i.e.,

$$A_R \circ A_S = \delta_{R,S} A_R, \quad \sum_{R \in \mathcal{R}} A_R = J.$$

Remark 2.3. If (X, \mathcal{R}) arises from a group action as in Remark 2.2, then \mathbf{A} coincides with the centralizer algebra (or Hecke algebra or commutant) for the corresponding permutation representation $g \mapsto P_g$ ($g \in \mathfrak{G}$) on V , i.e.,

$$\mathbf{A} = \{B \in M_X(\mathbb{C}) : BP_g = P_g B \ (g \in \mathfrak{G})\}.$$

A subalgebra of $M_X(\mathbb{C})$ is called a *coherent algebra* if it contains J , and is closed under \circ and \dagger . We note that the coherent algebras are precisely the adjacency algebras of coherent configurations. It is clear that the intersection of coherent algebras in $M_X(\mathbb{C})$ is again a coherent algebra. In particular, for any subset \mathcal{S} of $M_X(\mathbb{C})$, we can speak of the smallest coherent algebra containing \mathcal{S} , which we call the *coherent closure* of \mathcal{S} .

From now on, we assume that (X, \mathcal{R}) is an association scheme. As is the case for many examples of association schemes, we write

$$\mathcal{R} = \{R_0, R_1, \dots, R_n\}, \quad \text{where } \mathcal{R}_0 = \{R_0\},$$

and say that (X, \mathcal{R}) has n classes. We will then abbreviate $p_{i,j}^k = p_{R_i, R_j}^{R_k}$, $A_i = A_{R_i}$, and so on. The adjacency algebra \mathbf{A} is commutative in this case, and hence it has a basis E_0, E_1, \dots, E_n consisting of the (central) primitive idempotents, i.e.,

$$E_i E_j = \delta_{i,j} E_i, \quad \sum_{i=0}^n E_i = I.$$

Put differently, $E_0 V, E_1 V, \dots, E_n V$ are the maximal common eigenspaces (or homogeneous components or isotypic components) of \mathbf{A} , and the E_i are the corresponding orthogonal projections. Since the A_i are real symmetric matrices, so are the E_i . Note that the matrix $|X|^{-1} J \in \mathbf{A}$ is an idempotent with rank one, and thus primitive. We will always set

$$E_0 = \frac{1}{|X|} J.$$

For convenience, we let

$$A_i = E_i := O \text{ (the zero matrix) if } i < 0 \text{ or } i > n.$$

Though our focus in this paper will be on Q -polynomial association schemes, we first recall the P -polynomial property for completeness. We say that the association scheme (X, \mathcal{R}) is P -polynomial (or *metric*) with respect to the ordering A_0, A_1, \dots, A_n if there

are non-negative integers a_i, b_i, c_i ($0 \leq i \leq n$) such that $b_n = c_0 = 0$, $b_{i-1}c_i \neq 0$ ($1 \leq i \leq n$), and

$$A_1 A_i = b_{i-1} A_{i-1} + a_i A_i + c_{i+1} A_{i+1} \quad (0 \leq i \leq n),$$

where b_{-1} and c_{n+1} are indeterminates. In this case, A_1 recursively generates \mathbf{A} , and hence has $n + 1$ distinct eigenvalues $\theta_0, \theta_1, \dots, \theta_n \in \mathbb{R}$, where we write

$$A_1 = \sum_{i=0}^n \theta_i E_i. \tag{2.1}$$

We note that (X, \mathcal{R}) is P -polynomial as above precisely when the graph (X, R_1) is a distance-regular graph and (X, R_i) is the distance- i graph of (X, R_1) ($0 \leq i \leq n$). See, e.g., [10, 12, 27, 15] for more information on distance-regular graphs.

We say that (X, \mathcal{R}) is Q -polynomial (or cometric) with respect to the ordering E_0, E_1, \dots, E_n if there are real scalars a_i^*, b_i^*, c_i^* ($0 \leq i \leq n$) such that $b_n^* = c_0^* = 0$, $b_{i-1}^* c_i^* \neq 0$ ($1 \leq i \leq n$), and

$$E_1 \circ E_i = \frac{1}{|X|} (b_{i-1}^* E_{i-1} + a_i^* E_i + c_{i+1}^* E_{i+1}) \quad (0 \leq i \leq n), \tag{2.2}$$

where b_{-1}^* and c_{n+1}^* are indeterminates. In this case, $|X|E_1$ recursively generates \mathbf{A} with respect to \circ , and hence has $n + 1$ distinct entries $\theta_0^*, \theta_1^*, \dots, \theta_n^* \in \mathbb{R}$, where we write

$$|X|E_1 = \sum_{i=0}^n \theta_i^* A_i. \tag{2.3}$$

We call the θ_i^* the dual eigenvalues of $|X|E_1$. We may remark that $E_1 \circ E_i$, being a principal submatrix of $E_1 \otimes E_i$, is positive semidefinite, so that the scalars a_i^*, b_i^* , and c_i^* are non-negative (the so-called Krein condition). The Q -polynomial association schemes are an important subject in their own right, and we refer the reader to [23, 29] and the references therein for recent activity.

Below we give two fundamental examples of P - and Q -polynomial association schemes, both of which come from transitive group actions. See [10, 12, 16] for the details.

Example 2.4. Let v and k be positive integers with $v > k$, and let X be the set of k -subsets of $\{1, 2, \dots, v\}$. Set $n = \min\{k, v - k\}$. For $x, y \in X$ and $0 \leq i \leq n$, we let $(x, y) \in R_i$ if $|x \cap y| = k - i$. The R_i are the orbitals of the symmetric group \mathfrak{S}_v acting on X . We call (X, \mathcal{R}) a Johnson scheme and denote it by $J(v, k)$. The eigenvalues of A_1 are given in decreasing order by

$$\theta_i = (k - i)(v - k - i) - i \quad (0 \leq i \leq n),$$

and $J(v, k)$ is Q -polynomial with respect to the corresponding ordering of the E_i (cf. (2.1)).

Example 2.5. Let $q \geq 2$ be an integer and let $X = \{0, 1, \dots, q - 1\}^n$. For $x, y \in X$ and $0 \leq i \leq n$, we let $(x, y) \in R_i$ if x and y differ in exactly i coordinate positions. The R_i are the orbitals of the wreath product $\mathfrak{S}_q \wr \mathfrak{S}_n$ of the symmetric groups \mathfrak{S}_q and \mathfrak{S}_n acting

on X . We call (X, \mathcal{R}) a *Hamming scheme* and denote it by $H(n, q)$. The eigenvalues of A_1 are given in decreasing order by

$$\theta_i = n(q - 1) - qi \quad (0 \leq i \leq n),$$

and $H(n, q)$ is Q -polynomial with respect to the corresponding ordering of the E_i (cf. (2.1)). The Hamming scheme $H(n, 2)$ is also known as the n -cube (or n -dimensional hypercube) and is denoted by \mathcal{Q}_n .

Assumption 2.6. For the rest of this section and in Section 3, we assume that (X, \mathcal{R}) is an association scheme and is Q -polynomial with respect to the ordering E_0, E_1, \dots, E_n of the primitive idempotents.

In general, for any positive semidefinite Hermitian matrices $B, C \in M_X(\mathbb{C})$, we have (cf. [45])

$$(B \circ C)V = \text{span}(BV \circ CV),$$

where

$$BV \circ CV = \{u \circ v : u \in BV, v \in CV\}.$$

Hence it follows from (2.2) that

$$\text{span}(E_1V \circ E_iV) = \begin{cases} E_{i-1}V + E_iV + E_{i+1}V & \text{if } a_i^* \neq 0, \\ E_{i-1}V + E_{i+1}V & \text{if } a_i^* = 0, \end{cases} \quad (0 \leq i \leq n), \quad (2.4)$$

from which it follows that

$$\begin{aligned} \sum_{i=0}^h \sum_{j=0}^k \text{span}(E_iV \circ E_jV) &= \sum_{i=0}^h \sum_{j=0}^k \text{span}(\underbrace{E_1V \circ \dots \circ E_1V}_{i \text{ times}} \circ E_jV) \\ &= \sum_{i=0}^{h+k} E_iV \end{aligned} \quad (2.5)$$

for $0 \leq h, k \leq n$. See also [10, Section 2.8].

We now fix a ‘base vertex’ $x \in X$. Let

$$X_i = \{y \in X : (x, y) \in R_i\} \quad (0 \leq i \leq n).$$

We call the X_i the *shells* (or *spheres* or *subconstituents*) of (X, \mathcal{R}) with respect to x . For every i ($0 \leq i \leq n$), define the diagonal matrix $E_i^* = E_i^*(x) \in M_X(\mathbb{C})$ by

$$(E_i^*)_{y,y} = \begin{cases} 1 & \text{if } y \in X_i, \\ 0 & \text{otherwise,} \end{cases} \quad (y \in X).$$

Then we have

$$E_i^* E_j^* = \delta_{i,j} E_i^*, \quad \sum_{i=0}^n E_i^* = I.$$

We call the E_i^* the *dual idempotents* of (X, \mathcal{R}) with respect to x . The subspace

$$\mathbf{A}^* = \mathbf{A}^*(x) = \text{span}\{E_0^*, E_1^*, \dots, E_n^*\}$$

is then a subalgebra of $M_X(\mathbb{C})$, which we call the *dual adjacency algebra* of (X, \mathcal{R}) with respect to x . The *Terwilliger algebra* (or *subconstituent algebra*) of (X, \mathcal{R}) with respect to x is the subalgebra $T = T(x)$ of $M_X(\mathbb{C})$ generated by A and A^* [46, 47, 48]. We note that T is semisimple as it is closed under \dagger .

Remark 2.7. If (X, \mathcal{R}) arises from a group action as in Remark 2.2, which we recall is generously transitive in this case, then T is a subalgebra of the centralizer algebra for the action of the stabilizer \mathfrak{G}_x of x in \mathfrak{G} . The two algebras are known to be equal, e.g., for $J(v, k)$ and $H(n, q)$; see [25, 43].

For every subset Y of X , let $\hat{Y} \in V$ be the characteristic vector of Y , i.e.,

$$(\hat{Y})_y = \begin{cases} 1 & \text{if } y \in Y, \\ 0 & \text{otherwise,} \end{cases} \quad (y \in X).$$

In particular, \hat{X} denotes the all-ones vector in V . We will simply write \hat{x} for the characteristic vector of the singleton $\{x\}$. With this notation established, we have

$$\hat{X}_i = E_i^* \hat{X} = A_i \hat{x} \quad (0 \leq i \leq n),$$

from which it follows that

$$T\hat{x} = \text{span}\{\hat{X}_i : 0 \leq i \leq n\} = \text{span}\{E_i \hat{x} : 0 \leq i \leq n\}. \tag{2.6}$$

The T -module $T\hat{x}$ is easily seen to be irreducible with dimension $n + 1$ (cf. [46, Lemma 3.6]), and is called the *primary T -module*.

We define the *dual adjacency matrix* $A_1^* = A_1^*(x) \in M_X(\mathbb{C})$ by (cf. (2.3))

$$A_1^* = |X| \text{diag } E_1 \hat{x} = \sum_{i=0}^n \theta_i^* E_i^*. \tag{2.7}$$

Since the θ_i^* are mutually distinct, A_1^* generates A^* . Moreover, since

$$A_1^* v = |X|(E_1 \hat{x}) \circ v \quad (v \in V),$$

it follows from (2.4) that

$$E_i A_1^* E_j = O \quad \text{if } |i - j| > 1 \quad (0 \leq i, j \leq n). \tag{2.8}$$

Let W be an irreducible T -module. We define the *dual support* W_s^* , the *dual endpoint* $r^*(W)$, and the *dual diameter* $d^*(W)$ of W by

$$W_s^* = \{i : E_i W \neq 0\}, \quad r^*(W) = \min W_s^*, \quad d^*(W) = |W_s^*| - 1,$$

respectively. We call W *dual thin* if $\dim E_i W \leq 1$ ($0 \leq i \leq n$). We note that the primary T -module $T\hat{x}$ is dual thin, and that it is a unique irreducible T -module up to isomorphism which has dual endpoint zero or dual diameter n . The following lemma is an easy consequence of (2.8):

Lemma 2.8 ([46, Lemma 3.12]). *With reference to Assumption 2.6, write $A_1^* = A_1^*(x)$, $A^* = A^*(x)$, $T = T(x)$. Let W be an irreducible T -module and set $r^* = r^*(W)$, $d^* = d^*(W)$. Then the following hold:*

1. $A_1^* E_i W \subset E_{i-1} W + E_i W + E_{i+1} W$ ($0 \leq i \leq n$).
2. $W_s^* = \{r^*, r^* + 1, \dots, r^* + d^*\}$.
3. $E_i A_1^* E_j W \neq 0$ if $|i - j| = 1$ ($r^* \leq i, j \leq r^* + d^*$).
4. Suppose that W is dual thin. Then

$$\sum_{h=0}^i E_{r^*+h} W = \sum_{h=0}^i (A_1^*)^h E_{r^*} W \quad (0 \leq i \leq d^*).$$

In particular, $W = A^* E_{r^*} W$.

3 Relative t -designs in Q -polynomial association schemes

In this section, we develop some general theory on relative t -designs in Q -polynomial association schemes.

Recall Assumption 2.6. Throughout this section, we fix a base vertex $x \in X$, and write $E_i^* = E_i^*(x)$ ($0 \leq i \leq n$), $A_1^* = A_1^*(x)$, $A^* = A^*(x)$, and $T = T(x)$. In Introduction, we meant by a weighted subset of X a pair (Y, ω) of a subset Y of X and a function $\omega: Y \rightarrow (0, \infty)$. For convenience, however, we extend the domain of ω to X by setting $\omega(y) = 0$ for every $y \in X \setminus Y$. We will also naturally identify V with the set of complex functions on X , so that $\omega \in V$ and $Y = \text{supp } \omega$. In our discussions on relative t -designs, we will often consider the set

$$L = L_Y = \{\ell : Y \cap X_\ell \neq \emptyset\}, \tag{3.1}$$

and say that (Y, ω) is supported on $\bigsqcup_{\ell \in L} X_\ell$.

For comparison, we begin with the algebraic definition of t -designs in (X, \mathcal{R}) due to Delsarte [16, 17].

Definition 3.1. A weighted subset (Y, ω) of X is called a t -design in (X, \mathcal{R}) if $E_i \omega = 0$ for $1 \leq i \leq t$.

Delsarte [18] generalized this concept as follows:

Definition 3.2. A weighted subset (Y, ω) of X is called a relative t -design in (X, \mathcal{R}) (with respect to x) if $E_i \omega \in \text{span}\{E_i \hat{x}\}$ for $1 \leq i \leq t$.

Remark 3.3. Delsarte introduced the concept of t -designs for subsets Y of X in [16], i.e., when $\omega = \hat{Y}$, whereas in [17, 18] he mostly considered general (i.e., not necessarily non-negative) non-zero vectors $\omega \in V$ in the discussions on t -designs and relative t -designs. Some facts/results below, such as Examples 3.4 and 3.5, Proposition 3.6, and Theorem 3.8, are still valid for general $\omega \in V$, but the Fisher-type lower bound on $|Y| = |\text{supp } \omega|$ (cf. Theorem 3.9) makes sense only when ω is non-negative.

For the Johnson and Hamming schemes, Delsarte [16, 17, 18] showed that these algebraic concepts indeed have geometric interpretations:

Example 3.4. Let (X, \mathcal{R}) be the Johnson scheme $J(v, k)$ from Example 2.4. Then (Y, ω) is a t -design if and only if, for every t -subset z of $\{1, 2, \dots, v\}$, the sum λ_z of the values

$\omega(y)$ over those $y \in Y$ such that $z \subset y$, is a constant independent of z . On the other hand, (Y, ω) is a relative t -design if and only if the above λ_z depends only on $|x \cap z|$. We note that (Y, \hat{Y}) is a t -design if and only if Y is a t - (v, k, λ) design (cf. [13, Chapter II.4]) for some λ .

Example 3.5. Let (X, \mathcal{R}) be the Hamming scheme $H(n, q)$ from Example 2.5. Then (Y, ω) is a t -design if and only if, for every t -subset \mathcal{T} of $\{1, 2, \dots, n\}$ and every function $f: \mathcal{T} \rightarrow \{0, 1, \dots, q - 1\}$, the sum $\lambda_{\mathcal{T}, f}$ of the values $\omega(y)$ over those $y = (y_1, y_2, \dots, y_n) \in Y$ such that $y_i = f(i)$ ($i \in \mathcal{T}$), is a constant independent of the pair (\mathcal{T}, f) . On the other hand, (Y, ω) is a relative t -design if and only if the above $\lambda_{\mathcal{T}, f}$ depends only on $|\{i \in \mathcal{T} : x_i = f(i)\}|$, where $x = (x_1, x_2, \dots, x_n)$. We note that (Y, \hat{Y}) is a t -design if and only if the transpose of the $|Y| \times n$ matrix formed by arranging the elements of Y (in any order) is an orthogonal array $\text{OA}(|Y|, n, q, t)$ (cf. [13, Chapter III.6]). For the case $q = 2$, i.e., for \mathcal{Q}_n , if we choose the base vertex as $x = (0, 0, \dots, 0)$, then (Y, \hat{Y}) is a relative t -design if and only if Y is a regular t -wise balanced design of type t - (n, L, λ) (cf. [38, Section 4.4]) for some λ , where L is from (3.1), and where we identify the elements of $X = \{0, 1\}^n$ with their supports.

Similar results hold for some other important families of P - and Q -polynomial association schemes; see, e.g., [17, 18, 19, 34, 41].

Proposition 3.6 (cf. [3, Theorem 4.5]). *With reference to Assumption 2.6, let (Y, ω) be a weighted subset supported on $\bigsqcup_{\ell \in L} X_\ell$. Then we have*

$$\omega|_{\mathbf{T}\hat{x}} = \sum_{\ell \in L} \frac{\langle \omega, \hat{X}_\ell \rangle}{|X_\ell|} \hat{X}_\ell, \tag{3.2}$$

where $\omega|_{\mathbf{T}\hat{x}}$ denotes the orthogonal projection of ω on the primary \mathbf{T} -module $\mathbf{T}\hat{x}$. Moreover, (Y, ω) is a relative t -design if and only if

$$\langle \omega, v \rangle = \langle \omega|_{\mathbf{T}\hat{x}}, v \rangle = \sum_{\ell \in L} \frac{\langle \omega, \hat{X}_\ell \rangle}{|X_\ell|} \langle \hat{X}_\ell, v \rangle$$

for every $v \in \sum_{i=0}^t E_i V$.

Proof. Recall (2.6). The first part follows since the \hat{X}_i form an orthogonal basis of $\mathbf{T}\hat{x}$ with $\|\hat{X}_i\|^2 = |X_i|$. The second part is also immediate from

$$E_i \omega \in \text{span}\{E_i \hat{x}\} \iff E_i \omega \in \mathbf{T}\hat{x} \iff E_i \omega|_{\mathbf{T}\hat{x}} = E_i \omega. \quad \square$$

Remark 3.7. It is clear that (X_ℓ, \hat{X}_ℓ) is a relative n -design for every $0 \leq \ell \leq n$. Hence, if (Y, ω) is a relative t -design such that $X_\ell \subset Y$ for some ℓ , and if ω is constant on X_ℓ , then the weighted subset $(Y \setminus X_\ell, (I - E_\ell^*)\omega)$ obtained by discarding X_ℓ from Y is again a relative t -design. This observation is particularly important when applying Theorem 3.8 below; for example, we can always assume that $0 \notin L$.

The following is a slight generalization of Delsarte’s Assmus–Mattson theorem for Q -polynomial association schemes [18, Theorem 8.4], and can also be viewed as a variation of [9, Theorem 3.3], which in turn generalizes [28, Proposition 1]. See also [11]. The proof is in fact identical to that of [44, Theorem 4.3], but we include it below because of the potential importance of the result.

Theorem 3.8. *With reference to Assumption 2.6, let (Y, ω) be a relative t -design supported on $\bigsqcup_{\ell \in L} X_\ell$. Then $(Y \cap X_\ell, E_\ell^* \omega)$ is a relative $(t - |L| + 1)$ -design for every $\ell \in L$.*

Proof. Let $U = (\mathbf{T}\hat{x})^\perp$ be the orthogonal complement of $\mathbf{T}\hat{x}$ in V , which we recall is the sum of all the non-primary irreducible \mathbf{T} -modules in V . On the one hand, we have

$$\omega|_U \in \sum_{\ell \in L} E_\ell^* U.$$

Since A_1^* generates \mathbf{A}^* and has at most $|L|$ distinct eigenvalues on this subspace (cf. (2.7)), it follows that

$$\mathbf{A}^* \omega|_U = \text{span}\{\omega|_U, A_1^* \omega|_U, \dots, (A_1^*)^{|L|-1} \omega|_U\}. \tag{3.3}$$

On the other hand, since $E_0 U = 0$, that (Y, ω) is a relative t -design is rephrased as

$$\omega|_U \in \sum_{i=t+1}^n E_i U.$$

Hence it follows from (2.8) and (3.3) that

$$\mathbf{A}^* \omega|_U \subset \sum_{k=0}^{|L|-1} (A_1^*)^k \sum_{i=t+1}^n E_i U \subset \sum_{i=t-|L|+2}^n E_i U.$$

In particular, for every $\ell \in L$ we have

$$E_\ell^* \omega|_U \in \sum_{i=t-|L|+2}^n E_i U.$$

In other words, $(Y \cap X_\ell, E_\ell^* \omega)$ is a relative $(t - |L| + 1)$ -design, as desired. □

Bannai and Bannai [3, Theorem 4.8] established the following Fisher-type lower bound on the size of a relative t -design with t even:

Theorem 3.9. *With reference to Assumption 2.6, let (Y, ω) be a relative $2e$ -design ($e \in \mathbb{N}$) supported on $\bigsqcup_{\ell \in L} X_\ell$. Then*

$$|Y| \geq \dim \left(\sum_{\ell \in L} E_\ell^* \right) \binom{e}{\sum_{i=0} E_i V}.$$

Definition 3.10. A relative $2e$ -design (Y, ω) is called *tight* if equality holds above.

Recall from Example 3.5 that the relative t -designs in the hypercubes are equivalent to the weighted regular t -wise balanced designs.

Example 3.11. Let (X, \mathscr{R}) be the n -cube \mathcal{Q}_n from Example 2.5. Xiang [51] showed that if $e \leq \ell \leq n - e$ for every $\ell \in L$, then

$$\dim \left(\sum_{\ell \in L} E_\ell^* \right) \binom{e}{\sum_{i=0} E_i V} = \sum_{i=0}^{\min\{|L|-1, e\}} \binom{n}{e-i}. \tag{3.4}$$

We may remark that (cf. [12, Theorem 9.2.1])

$$\dim E_i V = \binom{n}{i} \quad (0 \leq i \leq n). \tag{3.5}$$

See also [32] and [6, Theorem 2.7, Example 2.9].

Example 3.12. Consider a symmetric $2-(n + 1, k, \lambda)$ design (cf. [13, Chapter II.6]). Observe that removing a point yields a tight relative 2-design in \mathcal{Q}_n with $L = \{k - 1, k\}$. Likewise, taking the complement of every block which contains a given point followed by removing that point gives rise to a tight relative 2-design in \mathcal{Q}_n with $L = \{k, n + 1 - k\}$. The complement of this is yet another example¹ such that $L = \{k - 1, n - k\}$. See [32, Section 3] and [50, Theorem 8]. Note that the weights are constant for these three examples. On the other hand, Bannai, Bannai, and Bannai [5, Theorem 2.2] showed that there is a tight relative 2-design in \mathcal{Q}_n with $L = \{2, n/2\}$ for $n \equiv 6 \pmod{8}$, provided that a Hadamard matrix of order $n/2 + 1$ exists. This construction provides examples in which the weights take two distinct values depending on the shells. See also [53].

Example 3.13. Working with the tight $4-(23, 7, 1)$ and $4-(23, 16, 52)$ designs instead of a symmetric $2-(n + 1, k, \lambda)$ design as in Example 3.12, we obtain four tight relative 4-designs in \mathcal{Q}_{22} with constant weight such that

$$L \in \{\{6, 7\}, \{6, 15\}, \{7, 16\}, \{15, 16\}\}.$$

See [9, Theorem 6.3] and [32, Section 3].

Let (Y, ω) be a tight relative $2e$ -design supported on $\bigsqcup_{\ell \in L} X_\ell$. Bannai, Bannai, and Bannai [5, Theorem 2.1] showed that if the stabilizer of x in the automorphism group of (X, \mathcal{R}) acts transitively on each of the shells X_i then ω is constant on $Y \cap X_\ell$ for every $\ell \in L$. The next theorem generalizes this result by replacing group actions by combinatorial regularity. Observe that the fibers of the coherent closure of \mathbf{T} are in general finer than the shells X_i .

Theorem 3.14. *With reference to Assumption 2.6, let (Y, ω) be a tight relative $2e$ -design ($e \in \mathbb{N}$) supported on $\bigsqcup_{\ell \in L} X_\ell$. For every $\ell \in L$, the weight ω is constant on $Y \cap X_\ell$ provided that X_ℓ remains a fiber of the coherent closure of \mathbf{T} .*

Proof. Let (cf. (3.2))

$$D = \text{diag } \omega, \quad \tilde{D} = \text{diag } \omega|_{\mathbf{T}\hat{x}} = \sum_{\ell \in L} \frac{\langle \omega, \hat{X}_\ell \rangle}{|\hat{X}_\ell|} E_\ell^*.$$

Note that $\tilde{D} \in \mathbf{T}$. Let F be the orthogonal projection onto BV , where

$$B = \sqrt{\tilde{D}} \sum_{i=0}^e E_i \in \mathbf{T}.$$

Observe that

$$BV = (BB^\dagger)V,$$

¹It seems that this construction is missing in [50, Theorem 8].

and that F is written as a polynomial in the Hermitian (in fact, real and symmetric) matrix BB^\dagger . In particular, $F \in \mathcal{T}$.

Since (Y, ω) is tight, we have

$$\dim BV = \dim \sqrt{\tilde{D}} \left(\sum_{\ell \in L} E_\ell^* \right) \left(\sum_{i=0}^e E_i V \right) = |Y|.$$

Let $u_1, u_2, \dots, u_{|Y|}$ be an orthonormal basis of BV , and let

$$G = [u_1 \mid u_2 \mid \cdots \mid u_{|Y|}].$$

Then we have

$$F = GG^\dagger. \tag{3.6}$$

Let

$$D' = D|_{Y \times Y}, \quad \tilde{D}' = \tilde{D}|_{Y \times Y}, \quad F' = F|_{Y \times Y}, \quad G' = G|_{Y \times \{1, 2, \dots, |Y|\}},$$

where $|_{Y \times Y}$ etc. mean taking corresponding submatrices. Note that these are square matrices, and that D' and \tilde{D}' are invertible. Then it follows that

$$(G')^\dagger D' (\tilde{D}')^{-1} G' = I_{|Y|}. \tag{3.7}$$

Indeed, since we may write

$$u_i = \sqrt{\tilde{D}} v_i, \quad \text{where } v_i \in \sum_{r=0}^e E_r V \quad (1 \leq i \leq |Y|),$$

it follows from (2.5) (applied to $h = k = e$) and Proposition 3.6 that the (i, j) -entry of the LHS in (3.7) is equal to

$$(v_i)^\dagger D v_j = \langle \omega, v_i \circ \bar{v}_j \rangle = \langle \omega|_{\mathbf{T}\hat{x}}, v_i \circ \bar{v}_j \rangle = (v_i)^\dagger \tilde{D} v_j = \langle u_j, u_i \rangle = \delta_{i,j},$$

where $\bar{}$ means complex conjugate. By (3.6) and (3.7), we have

$$I_{|Y|} = D' (\tilde{D}')^{-1} G' (G')^\dagger = D' (\tilde{D}')^{-1} F',$$

so that

$$(D')^{-1} = (\tilde{D}')^{-1} F'. \tag{3.8}$$

In particular, F' is a diagonal matrix.

Now, let $\ell \in L$ and suppose that X_ℓ remains a fiber of the coherent closure of \mathcal{T} . Then the (y, y) -entry of $F \in \mathcal{T}$ is constant for $y \in X_\ell$ (cf. (A1) and (A2)), and the same is true for \tilde{D} . Hence it follows from (3.8) that $\omega(y) = D_{y,y}$ must be constant for $y \in Y \cap X_\ell$. This completes the proof. \square

4 The Terwilliger algebra of \mathcal{Q}_n

For the rest of this paper, we will focus on relative t -designs in the n -cube \mathcal{Q}_n from Example 2.5. We will need detailed descriptions of the Terwilliger algebra of \mathcal{Q}_n and its irreducible modules, and we collect these in this section. Thus, we assume that $(X, \mathcal{R}) = \mathcal{Q}_n$, where $X = \{0, 1\}^n$. We again fix a base vertex $x \in X$, and write $E_i^* = E_i^*(x)$ ($0 \leq i \leq n$), $A_1^* = A_1^*(x)$, and $\mathbf{T} = \mathbf{T}(x)$. The Q -polynomial ordering we consider is the one given in Example 2.5.²

Proposition 4.1 (cf. [39, Section I.C]). *We have*

$$\mathbf{T} = \text{span}\{E_i^* A_j E_k^* : 0 \leq i, j, k \leq n\}. \tag{4.1}$$

In particular, \mathbf{T} is a coherent algebra.

Proof. The RHS in (4.1) is a subspace of \mathbf{T} . Recall from Example 2.5 that \mathcal{Q}_n admits the action of $\mathfrak{G} = \mathfrak{S}_2 \wr \mathfrak{S}_n$. The stabilizer \mathfrak{G}_x of x in \mathfrak{G} is isomorphic to \mathfrak{S}_n , and it is immediate to see that every orbital of \mathfrak{G}_x is of the form

$$\{(y, z) \in X \times X : (x, y) \in R_i, (y, z) \in R_j, (z, x) \in R_k\}$$

for some i, j , and k , where the corresponding adjacency matrix is $E_i^* A_j E_k^*$. Hence the RHS in (4.1) agrees with the centralizer algebra for the action of \mathfrak{G}_x on X , which is a coherent algebra; cf. Remark 2.3. Since \mathbf{T} is generated by the A_i and the E_i^* , the result follows. □

Lemma 4.2. *For $0 \leq i, j, k \leq n$, we have $E_i^* A_j E_k^* \neq O$ if and only if*

$$j \in \{|i - k|, |i - k| + 2, |i - k| + 4, \dots, \min\{i + k, 2n - i - k\}\}.$$

Proof. Routine. □

Next we recall basic facts about the irreducible \mathbf{T} -modules. Let W be an irreducible \mathbf{T} -module. We define the *support* W_s , the *endpoint* $r(W)$, and the *diameter* $d(W)$ of W by

$$W_s = \{i : E_i^* W \neq 0\}, \quad r(W) = \min W_s, \quad d(W) = |W_s| - 1,$$

respectively. We call W *thin* if $\dim E_i^* W \leq 1$ ($0 \leq i \leq n$).

Theorem 4.3 (cf. [26]). *Let W be an irreducible \mathbf{T} -module and set $r = r(W)$, $r^* = r^*(W)$, $d = d(W)$, and $d^* = d^*(W)$. Then W is thin, dual thin, and we have*

$$r = r^*, \quad d = d^* = n - 2r, \quad W_s = W_s^* = \{r, r + 1, \dots, n - r\}.$$

Moreover, the isomorphism class of W is determined by r .

Remark 4.4. Recall that the universal enveloping algebra $U(\mathfrak{sl}_2(\mathbb{C}))$ is defined by the generators $\mathfrak{r}, \mathfrak{h}$ and the relations

$$\mathfrak{r}\mathfrak{h} - \mathfrak{h}\mathfrak{r} = \mathfrak{h}, \quad \mathfrak{h}\mathfrak{r} - \mathfrak{r}\mathfrak{h} = 2\mathfrak{r}, \quad \mathfrak{h}\mathfrak{h} - \mathfrak{h}\mathfrak{h} = -2\mathfrak{h}.$$

²If n is even then \mathcal{Q}_n has another Q -polynomial ordering $E_0, E_{n-1}, E_2, E_{n-3}, \dots$ in terms of the natural ordering; cf. [10, p. 305].

There is a surjective homomorphism $U(\mathfrak{sl}_2(\mathbb{C})) \rightarrow \mathbf{T}$ such that (cf. [26, Lemma 7.5])

$$\mathfrak{k} \mapsto \sum_{i=1}^n E_{i-1} A_1^* E_i, \quad \mathfrak{h} \mapsto \sum_{i=0}^{n-1} E_{i+1} A_1^* E_i, \quad \mathfrak{h} \mapsto A_1.$$

Every irreducible \mathbf{T} -module is then irreducible as an $\mathfrak{sl}_2(\mathbb{C})$ -module. We also obtain another surjective homomorphism $U(\mathfrak{sl}_2(\mathbb{C})) \rightarrow \mathbf{T}$ by interchanging A_1 and A_1^* and replacing the E_i by the E_i^* above; cf. [26, Lemma 5.3].

From now on, we fix an orthogonal irreducible decomposition

$$V = \bigoplus_{W \in \Lambda} W \tag{4.2}$$

of the standard module V . In view of Theorem 4.3, let

$$\Lambda_r = \{W \in \Lambda : r(W) = r^*(W) = r\} \quad (0 \leq r \leq \lfloor n/2 \rfloor), \tag{4.3}$$

and fix a unit vector $v_W \in E_r W$ for each $W \in \Lambda_r$. Since

$$\dim E_i V = \sum_{W \in \Lambda} \dim E_i W = \sum_{r=0}^i |\Lambda_r| \quad (0 \leq i \leq \lfloor n/2 \rfloor) \tag{4.4}$$

by Theorem 4.3, it follows from (3.5) that

$$|\Lambda_r| = \binom{n}{r} - \binom{n}{r-1} \quad (0 \leq r \leq \lfloor n/2 \rfloor).$$

It is known (cf. [26, Theorem 9.2]) that if $W \in \Lambda_r$ then the vectors

$$E_r^* v_W, E_{r+1}^* v_W, \dots, E_{n-r}^* v_W \tag{4.5}$$

form an orthogonal basis of W , called a *standard basis* of W . By [26, Lemma 6.6], we also have

$$\|E_i^* v_W\|^2 = \binom{n-2r}{i-r} \|E_r^* v_W\|^2 \quad (r \leq i \leq n-r). \tag{4.6}$$

We note that

$$1 = \|v_W\|^2 = \sum_{i=r}^{n-r} \|E_i^* v_W\|^2 = 2^{n-2r} \|E_r^* v_W\|^2. \tag{4.7}$$

For $W, W' \in \Lambda_r$, we observe that the linear map $W \rightarrow W'$ defined by

$$E_i^* v_W \mapsto E_i^* v_{W'} \quad (r \leq i \leq n-r)$$

is an isometric isomorphism of \mathbf{T} -modules. Let

$$\check{E}_r^{i,j} = \frac{2^{n-2r}}{\sqrt{\binom{n-2r}{i-r} \binom{n-2r}{j-r}}} \sum_{W \in \Lambda_r} (E_i^* v_W)(E_j^* v_W)^\dagger \quad (r \leq i, j \leq n-r). \tag{4.8}$$

Then we have

$$(\check{E}_r^{i,j})^\dagger = \check{E}_r^{j,i} \quad (r \leq i, j \leq n - r), \tag{4.9}$$

and from (4.6) and (4.7) it follows that

$$\check{E}_r^{i,j} \check{E}_{r'}^{i',j'} = \delta_{r,r'} \delta_{j,i'} \check{E}_r^{i,j'}$$

for $0 \leq r, r' \leq \lfloor n/2 \rfloor$, $r \leq i, j \leq n - r$, and $r' \leq i', j' \leq n - r'$. By Theorem 4.3 and Wedderburn’s theorem (cf. [14, Section 3]), \mathbf{T} is isomorphic to the direct sum of full matrix algebras

$$\mathbf{T} \cong \bigoplus_{r=0}^{\lfloor n/2 \rfloor} M_{n-2r+1}(\mathbb{C}),$$

and the $\check{E}_r^{i,j}$ form an orthogonal basis of \mathbf{T} . See also [24, Section 2]. We note that

$$E_i^* \mathbf{T} E_j^* = \text{span}\{\check{E}_r^{i,j} : 0 \leq r \leq \min\{i, j, n - i, n - j\}\} \quad (0 \leq i, j \leq n). \tag{4.10}$$

We now recall the *Hahn polynomials* [31, Section 1.5]

$$Q_r(\xi; \alpha, \beta, N) = {}_3F_2\left(\begin{matrix} -\xi, -r, r + \alpha + \beta + 1 \\ \alpha + 1, -N \end{matrix} \middle| 1\right) \in \mathbb{R}[\xi] \quad (0 \leq r \leq N), \tag{4.11}$$

where

$${}_sF_t\left(\begin{matrix} a_1, \dots, a_s \\ b_1, \dots, b_t \end{matrix} \middle| c\right) = \sum_{i=0}^{\infty} \frac{(a_1)_i \cdots (a_s)_i c^i}{(b_1)_i \cdots (b_t)_i i!},$$

and

$$(a)_i = a(a + 1) \cdots (a + i - 1).$$

For $\alpha, \beta > -1$, or for $\alpha, \beta < -N$, we have

$$\begin{aligned} \sum_{\xi=0}^N \binom{\alpha + \xi}{\xi} \binom{\beta + N - \xi}{N - \xi} Q_r(\xi; \alpha, \beta, N) Q_{r'}(\xi; \alpha, \beta, N) \\ = \delta_{r,r'} \frac{(-1)^r (r + \alpha + \beta + 1)_{N+1} (\beta + 1)_r r!}{(2r + \alpha + \beta + 1) (\alpha + 1)_r (-N)_r N!}. \end{aligned} \tag{4.12}$$

Our aim is to describe the entries of the $\check{E}_r^{i,j}$. In view of (4.9), we will assume for the rest of this section that

$$0 \leq i \leq j \leq n.$$

By Proposition 4.1 and Lemma 4.2, we have

$$E_i^* \mathbf{T} E_j^* = \text{span}\{E_i^* A_{2\xi+j-i} E_j^* : 0 \leq \xi \leq \min\{i, n - j\}\}.$$

Moreover, it follows that (cf. (4.10))

$$\begin{aligned} E_i^* A_{2\xi+j-i} E_j^* \\ = \sum_{r=0}^{\min\{i, n-j\}} {}_3F_2\left(\begin{matrix} -\xi, -r, r - n - 1 \\ j - n, -i \end{matrix} \middle| 1\right) \frac{\binom{j}{i-\xi} \binom{n-j}{\xi} \binom{j-r}{j-i} \sqrt{\binom{n-2r}{j-r}}}{\binom{j}{i} \sqrt{\binom{n-2r}{i-r}}} \check{E}_r^{i,j}. \end{aligned} \tag{4.13}$$

This formula can be found in [33, Section 10]. See also [39, 49] for similar calculations.

If $i \leq n - j$ then

$${}_3F_2 \left(\begin{matrix} -\xi, -r, r - n - 1 \\ j - n, -i \end{matrix} \middle| 1 \right) = Q_r(\xi; j - n - 1, -j - 1, i).$$

Since

$$\binom{j}{i - \xi} \binom{n - j}{\xi} = (-1)^i \binom{j - n - 1 + \xi}{\xi} \binom{-j - 1 + i - \xi}{i - \xi},$$

it follows from (4.12) (applied to $\alpha = j - n - 1, \beta = -j - 1, N = i$) and (4.13) that, for $0 \leq r \leq i$,

$$\begin{aligned} \sum_{\xi=0}^i {}_3F_2 \left(\begin{matrix} -\xi, -r, r - n - 1 \\ j - n, -i \end{matrix} \middle| 1 \right) E_i^* A_{2\xi+j-i} E_j^* \\ = \frac{(-1)^r (r - n - 1)_{i+1} (-j)_r r!}{(2r - n - 1)(j - n)_r (-i)_r i!} \cdot \frac{(-1)^i \binom{j-r}{j-i} \sqrt{\binom{n-2r}{j-r}}}{\binom{j}{i} \sqrt{\binom{n-2r}{i-r}}} \check{E}_r^{i,j} \\ = \frac{\binom{n}{i} \binom{n-i}{r} \sqrt{\binom{n-2r}{j-r}}}{\left(\binom{n}{r} - \binom{n}{r-1} \right) \binom{n-j}{r} \sqrt{\binom{n-2r}{i-r}}} \check{E}_r^{i,j}. \end{aligned}$$

Likewise, if $n - j \leq i$ then

$${}_3F_2 \left(\begin{matrix} -\xi, -r, r - n - 1 \\ j - n, -i \end{matrix} \middle| 1 \right) = Q_r(\xi; -i - 1, i - n - 1, n - j).$$

In this case, since

$$\binom{j}{i - \xi} \binom{n - j}{\xi} \binom{j}{i}^{-1} = (-1)^{n-j} \binom{-i - 1 + \xi}{\xi} \binom{i - 1 - j - \xi}{n - j - \xi} \binom{n - i}{n - j}^{-1},$$

again it follows from (4.12) (applied to $\alpha = -i - 1, \beta = i - n - 1, N = n - j$) and (4.13) that, for $0 \leq r \leq n - j$,

$$\begin{aligned} \sum_{\xi=0}^{n-j} {}_3F_2 \left(\begin{matrix} -\xi, -r, r - n - 1 \\ j - n, -i \end{matrix} \middle| 1 \right) E_i^* A_{2\xi+j-i} E_j^* \\ = \frac{(-1)^r (r - n - 1)_{n-j+1} (i - n)_r r!}{(2r - n - 1)(-i)_r (j - n)_r (n - j)!} \cdot \frac{(-1)^{n-j} \binom{j-r}{j-i} \sqrt{\binom{n-2r}{j-r}}}{\binom{n-i}{n-j} \sqrt{\binom{n-2r}{i-r}}} \check{E}_r^{i,j} \\ = \frac{\binom{n}{i} \binom{n-i}{r} \sqrt{\binom{n-2r}{j-r}}}{\left(\binom{n}{r} - \binom{n}{r-1} \right) \binom{n-j}{r} \sqrt{\binom{n-2r}{i-r}}} \check{E}_r^{i,j}. \end{aligned}$$

In either case, it follows that

$$\check{E}_r^{i,j} = \frac{\left(\binom{n}{r} - \binom{n}{r-1} \right) \binom{n-j}{r} \sqrt{\binom{n-2r}{i-r}}}{\binom{n}{i} \binom{n-i}{r} \sqrt{\binom{n-2r}{j-r}}}$$

$$\times \sum_{\xi=0}^{\min\{i,n-j\}} {}_3F_2\left(-\xi, -r, r-n-1 \mid j-n, -i \mid 1\right) E_i^* A_{2\xi+j-i} E_j^* \tag{4.14}$$

for $0 \leq i \leq j \leq n$ and $0 \leq r \leq \min\{i, n-j\}$.

5 Tight relative $2e$ -designs on two shells in \mathcal{Q}_n

We retain the notation of the previous sections. In this section, we discuss tight relative $2e$ -designs (Y, ω) in \mathcal{Q}_n supported on two shells $X_\ell \sqcup X_m$, i.e., $L = \{\ell, m\}$ (cf. (3.1)). Recall from (3.4) that we have in this case

$$|Y| = \binom{n}{e} + \binom{n}{e-1},$$

but recall also that this is valid under the additional condition that $e \leq \ell, m \leq n - e$. However, both $(Y \cap X_\ell, E_\ell^* \omega)$ and $(Y \cap X_m, E_m^* \omega)$ are relative $(2e - 1)$ -designs by Theorem 3.8, so that if $\ell < 2e$ or $\ell > n - 2e$ for example, then $(Y \cap X_\ell, E_\ell^* \omega)$ must be trivial in view of Example 3.5, i.e., $X_\ell \subset Y$ and ω is constant on X_ℓ , and hence $(Y \cap X_m, E_m^* \omega)$ is by itself a relative $2e$ -design; cf. Remark 3.7. This shows that the above condition is not a restrictive one. We also note that

Lemma 5.1. *Let (Y, ω) be a relative t -design in \mathcal{Q}_n supported on $\bigsqcup_{\ell \in L} X_\ell$. Then $(Y', A_n \omega)$ is a relative t -design supported on $\bigsqcup_{\ell \in L} X_{n-\ell}$, where $Y' = \{y' : y \in Y\}$, and for every $y \in X, y'$ denotes the unique vertex such that $(y, y') \in R_n$.*

Proof. Immediate from $E_i A_n \in \text{span}\{E_i\}$ ($0 \leq i \leq n$). □

In view of the above comments, we now make the following assumption:

Assumption 5.2. *In this section, let (Y, ω) be a tight relative $2e$ -design ($e \in \mathbb{N}$) in \mathcal{Q}_n supported on two shells $X_\ell \sqcup X_m$, where*

$$e \leq \ell < m \leq n - \ell (\leq n - e).$$

Our aim is to show that Y then induces the structure of a coherent configuration with two fibers, and to obtain a necessary condition on the existence of such (Y, ω) akin to Delsarte’s theorem on tight $2e$ -designs. To this end, we first recall the proof of (3.4) given in [6, Theorem 2.7, Example 2.9] under the above assumption.

For convenience, set

$$E_L^* = E_\ell^* + E_m^*.$$

By (4.2) and (4.3), we have

$$E_L^* \left(\sum_{i=0}^e E_i V \right) = \sum_{r=0}^e \sum_{W \in \Lambda_r} E_L^* \left(\sum_{i=r}^e E_i W \right). \tag{5.1}$$

Let $W \in \Lambda_r$, where $0 \leq r \leq e$. Recall Theorem 4.3 and also the standard basis (4.5) of W . If $r = e$ then $E_e W$ is spanned by v_W , and hence we have

$$E_L^* E_e W = \text{span}\{E_L^* v_W\}.$$

Note that $E_L^*v_W$ is non-zero by Assumption 5.2, and hence

$$\dim E_L^*E_eW = 1$$

in this case. Suppose next that $0 \leq r < e$. On the one hand, since

$$E_L^* \left(\sum_{i=r}^e E_iW \right) \subset E_L^*W = E_\ell^*W + E_m^*W,$$

we have

$$\dim E_L^* \left(\sum_{i=r}^e E_iW \right) \leq 2.$$

On the other hand, it follows from (2.8) that

$$v_W, A_1^*v_W \in E_rW + E_{r+1}W \subset \sum_{i=r}^e E_iW, \tag{5.2}$$

and hence

$$E_L^*v_W, E_L^*A_1^*v_W \in E_L^* \left(\sum_{i=r}^e E_iW \right).$$

Moreover, we have (cf. (2.7))

$$E_L^*v_W = E_\ell^*v_W + E_m^*v_W, \quad E_L^*A_1^*v_W = \theta_\ell^*E_\ell^*v_W + \theta_m^*E_m^*v_W,$$

so that these two vectors are non-zero and are linearly independent by Assumption 5.2 and since $\theta_\ell^* \neq \theta_m^*$. It follows that

$$\dim E_L^* \left(\sum_{i=r}^e E_iW \right) = 2.$$

Note that in this case we in fact have

$$E_L^* \left(\sum_{i=r}^e E_iW \right) = \text{span}\{E_\ell^*v_W, E_m^*v_W\},$$

as

$$E_\ell^*v_W = E_L^* \frac{\theta_m^*I - A_1^*}{\theta_m^* - \theta_\ell^*} v_W, \quad E_m^*v_W = E_L^* \frac{\theta_\ell^*I - A_1^*}{\theta_\ell^* - \theta_m^*} v_W. \tag{5.3}$$

Combining these comments, we now obtain (3.4) as follows:

$$\begin{aligned} \dim E_L^* \left(\sum_{i=0}^e E_iV \right) &= \sum_{r=0}^e \sum_{W \in \Lambda_r} \dim E_L^* \left(\sum_{i=r}^e E_iW \right) \\ &= |\Lambda_e| + \sum_{r=0}^{e-1} 2|\Lambda_r| \\ &= \dim E_eV + \dim E_{e-1}V \end{aligned}$$

$$= \binom{n}{e} + \binom{n}{e-1},$$

where we have used (3.5) and (4.4).

By the above discussions, the set of vectors below forms an orthogonal basis of the subspace (5.1):

$$\left(\bigsqcup_{r=0}^{e-1} \bigsqcup_{W \in \Lambda_r} \{E_\ell^* v_W, E_m^* v_W\} \right) \bigsqcup \left(\bigsqcup_{W \in \Lambda_e} \{E_L^* v_W\} \right).$$

As in the proof of Theorem 3.14, let

$$D = \text{diag } \omega.$$

We next apply \sqrt{D} to the above basis vectors and compute their inner products. First, let $W, W' \in \bigsqcup_{r=0}^{e-1} \Lambda_r$. It is clear that

$$\left\langle \sqrt{D}E_\ell^* v_W, \sqrt{D}E_m^* v_{W'} \right\rangle = \left\langle \sqrt{D}E_m^* v_W, \sqrt{D}E_\ell^* v_{W'} \right\rangle = 0. \tag{5.4}$$

By (5.3), we have

$$(E_\ell^* \overline{v_W}) \circ (E_\ell^* v_{W'}) = E_L^* u,$$

where $\bar{}$ means complex conjugate, and

$$u = \left(\frac{\theta_m^* I - A_1^*}{\theta_m^* - \theta_\ell^*} \overline{v_W} \right) \circ \left(\frac{\theta_m^* I - A_1^*}{\theta_m^* - \theta_\ell^*} v_{W'} \right).$$

Observe that u belongs to $\sum_{i=0}^{2e} E_i V$ by (2.5) (applied to $h = k = e$) and (5.2). Hence, by Proposition 3.6 we have

$$\begin{aligned} \left\langle \sqrt{D}E_\ell^* v_W, \sqrt{D}E_\ell^* v_{W'} \right\rangle &= \langle \omega, E_L^* u \rangle \\ &= \langle \omega, u \rangle \\ &= \frac{\langle \omega, \hat{X}_\ell \rangle}{|X_\ell|} \langle \hat{X}_\ell, u \rangle + \frac{\langle \omega, \hat{X}_m \rangle}{|X_m|} \langle \hat{X}_m, u \rangle \\ &= \frac{\langle \omega, \hat{X}_\ell \rangle}{|X_\ell|} \langle \hat{X}_\ell, E_L^* u \rangle \\ &= \frac{\langle \omega, \hat{X}_\ell \rangle}{|X_\ell|} \langle E_\ell^* v_W, E_\ell^* v_{W'} \rangle \\ &= \delta_{W,W'} \frac{\langle \omega, \hat{X}_\ell \rangle}{|X_\ell|} \|E_\ell^* v_W\|^2. \end{aligned} \tag{5.5}$$

Likewise, we have

$$\left\langle \sqrt{D}E_m^* v_W, \sqrt{D}E_m^* v_{W'} \right\rangle = \delta_{W,W'} \frac{\langle \omega, \hat{X}_m \rangle}{|X_m|} \|E_m^* v_W\|^2. \tag{5.6}$$

Next, let $W \in \bigsqcup_{r=0}^{e-1} \Lambda_r$ and $W' \in \Lambda_e$. Then, by the same argument we have

$$\left\langle \sqrt{D}E_\ell^* v_W, \sqrt{D}E_L^* v_{W'} \right\rangle = \left\langle \sqrt{D}E_m^* v_W, \sqrt{D}E_L^* v_{W'} \right\rangle = 0. \tag{5.7}$$

Finally, let $W, W' \in \Lambda_e$. In this case, we have

$$\begin{aligned} & \left\langle \sqrt{D}E_L^*v_W, \sqrt{D}E_L^*v_{W'} \right\rangle \\ &= \delta_{W,W'} \left(\frac{\langle \omega, \hat{X}_\ell \rangle}{|\hat{X}_\ell|} \|E_\ell^*v_W\|^2 + \frac{\langle \omega, \hat{X}_m \rangle}{|\hat{X}_m|} \|E_m^*v_W\|^2 \right). \end{aligned} \quad (5.8)$$

Since (Y, ω) is a tight relative $2e$ -design, it follows from (5.4) – (5.8) that the set of vectors below is an orthogonal basis of the subspace $\sqrt{D}V = \text{span}\{\hat{y} : y \in Y\}$ of dimension $|Y| = \binom{n}{e} + \binom{n}{e-1}$:

$$\left(\bigsqcup_{r=0}^{e-1} \bigsqcup_{W \in \Lambda_r} \left\{ \sqrt{D}E_\ell^*v_W, \sqrt{D}E_m^*v_W \right\} \right) \sqcup \left(\bigsqcup_{W \in \Lambda_e} \left\{ \sqrt{D}E_L^*v_W \right\} \right).$$

For convenience, set

$$Y_\ell = Y \cap X_\ell, \quad Y_m = Y \cap X_m.$$

We will naturally make the following identification by discarding irrelevant entries:

$$\begin{aligned} \sqrt{D}E_\ell^*V &= \text{span}\{\hat{y} : y \in Y_\ell\} \longleftrightarrow \mathbb{C}^{Y_\ell}, \\ \sqrt{D}E_m^*V &= \text{span}\{\hat{y} : y \in Y_m\} \longleftrightarrow \mathbb{C}^{Y_m}. \end{aligned}$$

We write

$$\Lambda_r = \{W_r^1, W_r^2, \dots, W_r^{|\Lambda_r|}\} \quad (0 \leq r \leq e).$$

For $0 \leq r \leq e$, define a $|Y_\ell| \times |\Lambda_r|$ matrix H_r^ℓ and a $|Y_m| \times |\Lambda_r|$ matrix H_r^m by

$$\begin{aligned} H_r^\ell &= \left[\sqrt{D}E_\ell^*v_{W_r^1} \mid \cdots \mid \sqrt{D}E_\ell^*v_{W_r^{|\Lambda_r|}} \right], \\ H_r^m &= \left[\sqrt{D}E_m^*v_{W_r^1} \mid \cdots \mid \sqrt{D}E_m^*v_{W_r^{|\Lambda_r|}} \right]. \end{aligned}$$

We then define a characteristic matrix H of (Y, ω) by

$$H = \left[\begin{array}{c|c|c|c|c|c|c} H_0^\ell & \cdots & H_{e-1}^\ell & O & \cdots & O & H_e^\ell \\ \hline O & \cdots & O & H_0^m & \cdots & H_{e-1}^m & H_e^m \end{array} \right].$$

We note that H is a square matrix of size $|Y| = \binom{n}{e} + \binom{n}{e-1}$. By (4.6), (4.7), and (5.4) – (5.8), and since

$$|X_i| = \binom{n}{i} \quad (0 \leq i \leq n),$$

we have

$$H^\dagger H = \left(\bigoplus_{r=0}^{e-1} \kappa_r^\ell I_{|\Lambda_r|} \right) \oplus \left(\bigoplus_{r=0}^{e-1} \kappa_r^m I_{|\Lambda_r|} \right) \oplus \kappa_e I_{|\Lambda_e|}, \quad (5.9)$$

where

$$\kappa_r^\ell = \frac{\omega_\ell \binom{n-2r}{\ell-r}}{2^{n-2r} \binom{n}{\ell}}, \quad \kappa_r^m = \frac{\omega_m \binom{n-2r}{m-r}}{2^{n-2r} \binom{n}{m}} \quad (0 \leq r < e),$$

$$\kappa_e = \frac{\omega_\ell \binom{n-2e}{\ell-e}}{2^{n-2e} \binom{n}{\ell}} + \frac{\omega_m \binom{n-2e}{m-e}}{2^{n-2e} \binom{n}{m}},$$

and we abbreviate

$$\omega_\ell = \langle \omega, \hat{X}_\ell \rangle, \quad \omega_m = \langle \omega, \hat{X}_m \rangle.$$

Let K denote the diagonal matrix on the RHS in (5.9). Then it follows that

$$\begin{aligned} I_{|Y|} &= HK^{-1}H^\dagger \\ &= \left[\begin{array}{c|c} \sum_{r=0}^e \frac{1}{\kappa_r^\ell} H_r^\ell (H_r^\ell)^\dagger & \frac{1}{\kappa_e} H_e^\ell (H_e^m)^\dagger \\ \hline \frac{1}{\kappa_e} H_e^\ell (H_e^\ell)^\dagger & \sum_{r=0}^e \frac{1}{\kappa_r^m} H_r^m (H_r^m)^\dagger \end{array} \right], \end{aligned} \tag{5.10}$$

where we write

$$\kappa_e^\ell = \kappa_e^m := \kappa_e \tag{5.11}$$

for brevity. In particular, we have

$$\frac{1}{\kappa_e} H_e^\ell (H_e^m)^\dagger = O. \tag{5.12}$$

Moreover, from (5.9) and (5.10) it follows that

$$\left(\frac{1}{\kappa_r^\ell} H_r^\ell (H_r^\ell)^\dagger \right) \left(\frac{1}{\kappa_{r'}^\ell} H_{r'}^\ell (H_{r'}^\ell)^\dagger \right) = \delta_{r,r'} \frac{1}{\kappa_r^\ell} H_r^\ell (H_r^\ell)^\dagger \quad (0 \leq r, r' < e), \tag{5.13}$$

$$\frac{1}{\kappa_e} H_e^\ell (H_e^\ell)^\dagger = I_{|Y_\ell|} - \sum_{r=0}^{e-1} \frac{1}{\kappa_r^\ell} H_r^\ell (H_r^\ell)^\dagger, \tag{5.14}$$

$$\left(\frac{1}{\kappa_r^m} H_r^m (H_r^m)^\dagger \right) \left(\frac{1}{\kappa_{r'}^m} H_{r'}^m (H_{r'}^m)^\dagger \right) = \delta_{r,r'} \frac{1}{\kappa_r^m} H_r^m (H_r^m)^\dagger \quad (0 \leq r, r' < e), \tag{5.15}$$

$$\frac{1}{\kappa_e} H_e^m (H_e^m)^\dagger = I_{|Y_m|} - \sum_{r=0}^{e-1} \frac{1}{\kappa_r^m} H_r^m (H_r^m)^\dagger. \tag{5.16}$$

Note that the matrices $(\kappa_r^\ell)^{-1} H_r^\ell (H_r^\ell)^\dagger, (\kappa_r^m)^{-1} H_r^m (H_r^m)^\dagger$ ($0 \leq r < e$) are non-zero since H_r^ℓ, H_r^m are non-zero. Likewise, by setting

$$\kappa_r = \sqrt{\kappa_r^\ell \kappa_r^m} \quad (0 \leq r < e)$$

for brevity, we have

$$\left(\frac{1}{\kappa_r^\ell} H_r^\ell (H_r^\ell)^\dagger \right) \left(\frac{1}{\kappa_{r'}} H_{r'}^\ell (H_{r'}^m)^\dagger \right) = \delta_{r,r'} \frac{1}{\kappa_r} H_r^\ell (H_r^m)^\dagger \quad (0 \leq r, r' < e), \tag{5.17}$$

$$\left(\frac{1}{\kappa_r} H_r^\ell (H_r^m)^\dagger \right) \left(\frac{1}{\kappa_{r'}} H_{r'}^m (H_{r'}^m)^\dagger \right) = \delta_{r,r'} \frac{1}{\kappa_r} H_r^\ell (H_r^m)^\dagger \quad (0 \leq r, r' < e), \tag{5.18}$$

$$\left(\frac{1}{\kappa_r} H_r^\ell (H_r^m)^\dagger \right) \left(\frac{1}{\kappa_{r'}} H_{r'}^m (H_{r'}^\ell)^\dagger \right) = \delta_{r,r'} \frac{1}{\kappa_r^\ell} H_r^\ell (H_r^\ell)^\dagger \quad (0 \leq r, r' < e), \tag{5.19}$$

$$\left(\frac{1}{\kappa_r} H_r^m (H_r^m)^\dagger \right) \left(\frac{1}{\kappa_{r'}} H_{r'}^\ell (H_{r'}^m)^\dagger \right) = \delta_{r,r'} \frac{1}{\kappa_r^m} H_r^m (H_r^m)^\dagger \quad (0 \leq r, r' < e). \tag{5.20}$$

Since the matrices $(\kappa_r^\ell)^{-1}H_r^\ell(H_r^\ell)^\dagger, (\kappa_r^m)^{-1}H_r^m(H_r^m)^\dagger$ ($0 \leq r < e$) are non-zero, it follows from (5.17) – (5.20) that the matrices $(\kappa_r)^{-1}H_r^\ell(H_r^m)^\dagger$ ($0 \leq r < e$) are non-zero and are linearly independent.

It follows from Theorem 3.14 and Proposition 4.1 that ω is constant on each of Y_ℓ and Y_m , from which it follows that

$$D_{y,y} = \omega(y) = \begin{cases} \frac{\omega_\ell}{|Y_\ell|} & \text{if } y \in Y_\ell, \\ \frac{\omega_m}{|Y_m|} & \text{if } y \in Y_m. \end{cases} \tag{5.21}$$

Hence, by comparing with the formula (4.8) for the matrices $\check{E}_r^{i,j}$, we have

$$\frac{1}{\kappa_r^\ell}H_r^\ell(H_r^\ell)^\dagger = \frac{\binom{n}{\ell}}{|Y_\ell|}\check{E}_r^{\ell,\ell}|_{Y_\ell \times Y_\ell} \quad (0 \leq r < e), \tag{5.22}$$

$$\frac{1}{\kappa_e}H_e^\ell(H_e^\ell)^\dagger = \frac{\omega_\ell \binom{n-2e}{\ell-e}}{2^{n-2e}\kappa_e|Y_\ell|}\check{E}_e^{\ell,\ell}|_{Y_\ell \times Y_\ell}, \tag{5.23}$$

$$\frac{1}{\kappa_r^m}H_r^m(H_r^m)^\dagger = \frac{\binom{n}{m}}{|Y_m|}\check{E}_r^{m,m}|_{Y_m \times Y_m} \quad (0 \leq r < e), \tag{5.24}$$

$$\frac{1}{\kappa_e}H_e^m(H_e^m)^\dagger = \frac{\omega_m \binom{n-2e}{m-e}}{2^{n-2e}\kappa_e|Y_m|}\check{E}_e^{m,m}|_{Y_m \times Y_m}, \tag{5.25}$$

$$\frac{1}{\kappa_r}H_r^\ell(H_r^m)^\dagger = \frac{\sqrt{\binom{n}{\ell}\binom{n}{m}}}{\sqrt{|Y_\ell||Y_m|}}\check{E}_r^{\ell,m}|_{Y_\ell \times Y_m} \quad (0 \leq r < e), \tag{5.26}$$

$$\frac{1}{\kappa_e}H_e^\ell(H_e^m)^\dagger = \frac{\sqrt{\omega_\ell\omega_m \binom{n-2e}{\ell-e}\binom{n-2e}{m-e}}}{2^{n-2e}\kappa_e\sqrt{|Y_\ell||Y_m|}}\check{E}_e^{\ell,m}|_{Y_\ell \times Y_m}, \tag{5.27}$$

where $|_{Y_\ell \times Y_\ell}$ etc. mean taking corresponding submatrices. From (5.23) and (5.25) it follows that the matrices $(\kappa_e)^{-1}H_e^\ell(H_e^\ell)^\dagger, (\kappa_e)^{-1}H_e^m(H_e^m)^\dagger$ are also non-zero, since each of $\check{E}_e^{\ell,\ell}|_{Y_\ell \times Y_\ell}, \check{E}_e^{m,m}|_{Y_m \times Y_m}$ has non-zero constant diagonal entries by (4.14).

Let \mathbf{H}' be the set consisting of the $|Y| \times |Y|$ matrices of the form

$$\left[\begin{array}{c|c} \sum_{r=0}^e a_r^{\ell,\ell} \frac{1}{\kappa_r^\ell} H_r^\ell(H_r^\ell)^\dagger & \sum_{r=0}^{e-1} a_r^{\ell,m} \frac{1}{\kappa_r} H_r^\ell(H_r^m)^\dagger \\ \hline \sum_{r=0}^{e-1} a_r^{m,\ell} \frac{1}{\kappa_r} H_r^m(H_r^\ell)^\dagger & \sum_{r=0}^e a_r^{m,m} \frac{1}{\kappa_r^m} H_r^m(H_r^m)^\dagger \end{array} \right],$$

where $a_r^{\ell,\ell}$ etc. are in \mathbb{C} , and we are again using the notation (5.11). By (5.13) – (5.20) and the above comments, \mathbf{H}' is a \mathbb{C} -algebra with

$$\dim \mathbf{H}' = 4e + 2. \tag{5.28}$$

Define

$$S_{\ell,\ell}(Y) = \{j : R_j \cap (Y_\ell \times Y_\ell) \neq \emptyset\},$$

and define $S_{\ell,m}(Y)(= S_{m,\ell}(Y))$ and $S_{m,m}(Y)$ in the same manner. Let \mathbf{H} be the set consisting of the $|Y| \times |Y|$ matrices of the form

$$\left[\begin{array}{c|c} \sum_{j \in S_{\ell,\ell}(Y)} b_j^{\ell,\ell} A_j|_{Y_\ell \times Y_\ell} & \sum_{j \in S_{\ell,m}(Y)} b_j^{\ell,m} A_j|_{Y_\ell \times Y_m} \\ \hline \sum_{j \in S_{m,\ell}(Y)} b_j^{m,\ell} A_j|_{Y_m \times Y_\ell} & \sum_{j \in S_{m,m}(Y)} b_j^{m,m} A_j|_{Y_m \times Y_m} \end{array} \right], \tag{5.29}$$

where $b_j^{\ell,\ell}$ etc. are in \mathbb{C} . Then \mathbf{H} is a \mathbb{C} -vector space with

$$\dim \mathbf{H} = |S_{\ell,\ell}(Y)| + |S_{\ell,m}(Y)| + |S_{m,\ell}(Y)| + |S_{m,m}(Y)|. \tag{5.30}$$

Note that \mathbf{H} is closed under \circ . By (5.22) – (5.26) and Proposition 4.1 (or (4.14)), \mathbf{H}' is a subspace of \mathbf{H} .

By (4.14), (5.14), (5.22), and (5.23), we have

$$\begin{aligned} I_{|Y_\ell|} &= \sum_{r=0}^{e-1} \frac{\binom{n}{\ell}}{|Y_\ell|} \check{E}_r^{\ell,\ell} |_{Y_\ell \times Y_\ell} + \frac{\omega_\ell \binom{n-2e}{\ell-e}}{2^{n-2e} \kappa_e |Y_\ell|} \check{E}_e^{\ell,\ell} |_{Y_\ell \times Y_\ell} \\ &= \frac{1}{|Y_\ell|} \sum_{\xi=0}^{\min\{\ell, n-\ell\}} \left(\sum_{r=0}^{e-1} \left(\binom{n}{r} - \binom{n}{r-1} \right) {}_3F_2 \left(\begin{matrix} -\xi, -r, r-n-1 \\ \ell-n, -\ell \end{matrix} \middle| 1 \right) \right. \\ &\quad \left. + \frac{\omega_\ell \binom{n-2e}{\ell-e} \left(\binom{n}{e} - \binom{n}{e-1} \right)}{2^{n-2e} \kappa_e \binom{n}{\ell}} {}_3F_2 \left(\begin{matrix} -\xi, -e, e-n-1 \\ \ell-n, -\ell \end{matrix} \middle| 1 \right) \right) A_{2\xi} |_{Y_\ell \times Y_\ell}. \end{aligned} \tag{5.31}$$

Hence it follows that $\{\xi \neq 0 : 2\xi \in S_{\ell,\ell}(Y)\}$ is a set of zeros of the polynomial

$$\begin{aligned} \psi_e^{\ell,\ell}(\xi) &= \sum_{r=0}^{e-1} \left(\binom{n}{r} - \binom{n}{r-1} \right) {}_3F_2 \left(\begin{matrix} -\xi, -r, r-n-1 \\ \ell-n, -\ell \end{matrix} \middle| 1 \right) \\ &\quad + \frac{\omega_\ell \binom{n-2e}{\ell-e} \left(\binom{n}{e} - \binom{n}{e-1} \right)}{2^{n-2e} \kappa_e \binom{n}{\ell}} {}_3F_2 \left(\begin{matrix} -\xi, -e, e-n-1 \\ \ell-n, -\ell \end{matrix} \middle| 1 \right) \in \mathbb{R}[\xi]. \end{aligned} \tag{5.32}$$

Note that $\psi_e^{\ell,\ell}(\xi)$ has degree exactly e , from which it follows that

$$|S_{\ell,\ell}(Y)| \leq e + 1. \tag{5.33}$$

Likewise, we find that $\{\xi \neq 0 : 2\xi \in S_{m,m}(Y)\}$ is a set of zeros of the polynomial

$$\begin{aligned} \psi_e^{m,m}(\xi) &= \sum_{r=0}^{e-1} \left(\binom{n}{r} - \binom{n}{r-1} \right) {}_3F_2 \left(\begin{matrix} -\xi, -r, r-n-1 \\ m-n, -m \end{matrix} \middle| 1 \right) \\ &\quad + \frac{\omega_m \binom{n-2e}{m-e} \left(\binom{n}{e} - \binom{n}{e-1} \right)}{2^{n-2e} \kappa_e \binom{n}{m}} {}_3F_2 \left(\begin{matrix} -\xi, -e, e-n-1 \\ m-n, -m \end{matrix} \middle| 1 \right) \in \mathbb{R}[\xi], \end{aligned} \tag{5.34}$$

and hence that

$$|S_{m,m}(Y)| \leq e + 1. \tag{5.35}$$

Finally, by (4.14), (5.12), and (5.27), we have

$$\begin{aligned} O &= \frac{\sqrt{\omega_\ell \omega_m \binom{n-2e}{\ell-e} \binom{n-2e}{m-e}}}{2^{n-2e} \kappa_e \sqrt{|Y_\ell| |Y_m|}} \check{E}_e^{\ell,m} |_{Y_\ell \times Y_m} \\ &= \frac{\sqrt{\omega_\ell \omega_m} \binom{n-m}{e} \binom{n-2e}{\ell-e} \left(\binom{n}{e} - \binom{n}{e-1} \right)}{2^{n-2e} \kappa_e \sqrt{|Y_\ell| |Y_m|} \binom{n}{\ell} \binom{n-\ell}{e}} \end{aligned}$$

$$\times \sum_{\xi=0}^{\min\{\ell, n-m\}} {}_3F_2 \left(\begin{matrix} -\xi, -e, e-n-1 \\ m-n, -\ell \end{matrix} \middle| 1 \right) A_{2\xi+m-\ell} |Y_\ell \times Y_m|.$$

Hence it follows that $\{\xi : 2\xi + m - \ell \in S_{\ell, m}(Y)\}$ is a set of zeros of the polynomial

$$\psi_e^{\ell, m}(\xi) = {}_3F_2 \left(\begin{matrix} -\xi, -e, e-n-1 \\ m-n, -\ell \end{matrix} \middle| 1 \right) \in \mathbb{R}[\xi], \tag{5.36}$$

and that

$$|S_{\ell, m}(Y)| = |S_{m, \ell}(Y)| \leq e. \tag{5.37}$$

By (5.30), (5.33), (5.35), and (5.37), we have

$$\dim \mathbf{H} \leq 4e + 2.$$

Since \mathbf{H}' is a subspace of \mathbf{H} , it follows from (5.28) that $\mathbf{H} = \mathbf{H}'$. In particular, \mathbf{H} is a \mathbb{C} -algebra. It is also clear that \mathbf{H} is closed under \dagger and contains $J_{|\mathcal{Y}|}$. We now conclude that \mathbf{H} is a coherent algebra. Note also that equality holds in each of (5.33), (5.35), and (5.37).

To summarize:

Theorem 5.3. *Recall Assumption 5.2. With the above notation, the following hold:*

- (i) *The set \mathbf{H} from (5.29) is a coherent algebra of type $\begin{bmatrix} e+1 & e \\ e+1 & e \end{bmatrix}$.*
- (ii) *The sets of zeros of the polynomials $\psi_e^{\ell, \ell}(\xi)$, $\psi_e^{m, m}(\xi)$, and $\psi_e^{\ell, m}(\xi)$ from (5.32), (5.34), and (5.36) are given respectively by*

$$\{\xi \neq 0 : 2\xi \in S_{\ell, \ell}(Y)\}, \{\xi \neq 0 : 2\xi \in S_{m, m}(Y)\}, \text{ and} \\ \{\xi : 2\xi + m - \ell \in S_{\ell, m}(Y)\}.$$

In particular, the zeros of these polynomials are integral.

Concerning the scalars ω_ℓ and ω_m appearing in the polynomials $\psi_e^{\ell, \ell}(\xi)$ and $\psi_e^{m, m}(\xi)$, it follows that

Proposition 5.4. *Recall Assumption 5.2. The scalars ω_ℓ and ω_m satisfies*

$$\frac{\omega_m}{\omega_\ell} = \frac{\binom{n}{m} \binom{n-2e}{\ell-e}}{\binom{n}{\ell} \binom{n-2e}{m-e}} \cdot \frac{|Y_m| - \binom{n}{e-1}}{|Y_\ell| - \binom{n}{e-1}}.$$

In particular, the weight function ω is unique up to a scalar multiple.

Proof. By comparing the diagonal entries of both sides in (5.31), we have

$$1 = \frac{\psi_e^{\ell, \ell}(0)}{|Y_\ell|} = \frac{1}{|Y_\ell|} \left(\binom{n}{e-1} + \frac{\omega_\ell \binom{n-2e}{\ell-e} \left(\binom{n}{e} - \binom{n}{e-1} \right)}{2^{n-2e} \kappa_e \binom{n}{\ell}} \right).$$

Likewise,

$$1 = \frac{\psi_e^{m, m}(0)}{|Y_m|} = \frac{1}{|Y_m|} \left(\binom{n}{e-1} + \frac{\omega_m \binom{n-2e}{m-e} \left(\binom{n}{e} - \binom{n}{e-1} \right)}{2^{n-2e} \kappa_e \binom{n}{m}} \right).$$

By eliminating κ_e , we obtain the formula for $\omega_m(\omega_\ell)^{-1}$. The uniqueness of ω follows from this and (5.21). □

Example 5.5. Suppose that $e = 1$. In this case, Theorem 5.3(i) was previously obtained by Bannai, Bannai, and Bannai [5, Theorem 2.2 (i)]. Moreover, Theorem 5.3(ii) and Proposition 5.4 are together equivalent to [5, Proposition 4.3].

Example 5.6. Suppose that $e = 2$. Then we have

$$\begin{aligned} \psi_e^{\ell,m}(\xi) &= 1 + \frac{(-\xi)(-2)(1-n)}{(m-n)(-\ell)} + \frac{(-\xi)(1-\xi)(-2)(-1)(1-n)(2-n)}{(m-n)(m-n+1)(-\ell)(1-\ell)2} \\ &= 1 - \frac{2(n-1)\xi}{(n-m)\ell} + \frac{(n-1)(n-2)\xi(\xi-1)}{(n-m)(n-m-1)\ell(\ell-1)}. \end{aligned}$$

From Example 3.13 we find two parameter sets satisfying Assumption 5.2:

n	ℓ	m	ξ
22	6	7	3, 5
22	6	15	1, 3

The zeros ξ given in the last column are indeed integers. Note that the other two parameter sets in Example 3.13 correspond to the complements of these two; cf. Lemma 5.1. On the other hand, the existence of tight relative 4-designs with the following feasible parameter sets was left open in [9, Section 6]:

n	ℓ	m	ξ
37	9	16	$\frac{1}{14}(71 \pm \sqrt{337})$
37	9	21	$\frac{1}{14}(55 \pm \sqrt{337})$
41	15	16	$\frac{1}{26}(237 \pm \sqrt{1569})$
41	15	25	$\frac{1}{26}(153 \pm \sqrt{1569})$

Here, we are again taking Lemma 5.1 into account. Observe that the zeros ξ are irrational, thus proving the non-existence.

We end this section with a comment on the expressions of the polynomials $\psi_e^{\ell,\ell}(\xi)$ and $\psi_e^{m,m}(\xi)$. We first invoke the following identity which agrees with the formula of the backward shift operator on the dual Hahn polynomials (cf. [31, Section 1.6]):

$$\begin{aligned} \alpha(N+1)(\alpha+\beta+2r)Q_r(\xi; \alpha-1, \beta, N+1) \\ = (\alpha+r)(\alpha+\beta+r)(N+1-r)Q_r(\xi-1; \alpha, \beta, N) \\ - r(\alpha+\beta+N+1+r)(\beta+r)Q_{r-1}(\xi-1; \alpha, \beta, N). \end{aligned} \tag{5.38}$$

This can be routinely verified by writing the LHS as a linear combination of the polynomials $(1-\xi)_i$ ($0 \leq i \leq r$) using

$$(-\xi)_i = (1-\xi)_i - i(1-\xi)_{i-1},$$

and then comparing the coefficients of both sides. Setting $\alpha = \ell - n$, $\beta = -\ell - 1$, and $N = \ell - 1$ in (5.38), it follows that the first term of the RHS in (5.32) is rewritten as follows:

$$\sum_{r=0}^{e-1} \left(\binom{n}{r} - \binom{n}{r-1} \right) {}_3F_2 \left(\begin{matrix} -\xi, -r, r-n-1 \\ \ell-n, -\ell \end{matrix} \middle| 1 \right)$$

$$\begin{aligned}
 &= \sum_{r=0}^{e-1} \frac{n!(n-2r+1)}{r!(n-r+1)!} Q_r(\xi; \alpha-1, \beta, N+1) \\
 &= \frac{n!}{\ell(n-\ell)} \sum_{r=0}^{e-1} \left(\frac{(\ell-n+r)(r-n-1)(\ell-r)}{r!(n-r+1)!} Q_r(\xi-1; \alpha, \beta, N) \right. \\
 &\quad \left. - \frac{r(r+\ell-n-1)(r-\ell-1)}{r!(n-r+1)!} Q_{r-1}(\xi-1; \alpha, \beta, N) \right) \\
 &= \frac{n!}{\ell(n-\ell)} \cdot (-1) \frac{(\ell-n+e-1)(\ell-e+1)}{(e-1)!(n-e+1)!} Q_{e-1}(\xi-1; \alpha, \beta, N) \\
 &= \binom{n}{e-1} \frac{(n-\ell-e+1)(\ell-e+1)}{\ell(n-\ell)} {}_3F_2 \left(\begin{matrix} 1-\xi, 1-e, e-n-1 \\ \ell-n+1, 1-\ell \end{matrix} \middle| 1 \right).
 \end{aligned}$$

Likewise, the first term of the RHS in (5.34) is given by

$$\begin{aligned}
 &\sum_{r=0}^{e-1} \left(\binom{n}{r} - \binom{n}{r-1} \right) {}_3F_2 \left(\begin{matrix} -\xi, -r, r-n-1 \\ m-n, -m \end{matrix} \middle| 1 \right) \\
 &= \binom{n}{e-1} \frac{(n-m-e+1)(m-e+1)}{m(n-m)} {}_3F_2 \left(\begin{matrix} 1-\xi, 1-e, e-n-1 \\ m-n+1, 1-m \end{matrix} \middle| 1 \right).
 \end{aligned}$$

6 Zeros of the Hahn and Hermite polynomials

Recall the Hahn polynomials $Q_r(\xi; \alpha, \beta, N)$ from (4.11). Recall also that the zeros of orthogonal polynomials are always real and simple; see, e.g., [42, Theorem 3.3.1]. It is well known that we can obtain the Hermite polynomials as limits of the Hahn polynomials; cf. [30, 31]. In this section, we revisit this limit process and describe the limit behavior of the zeros of the $Q_r(\xi; \alpha, \beta, N)$, in a special case which is suited to our purpose.

Assumption 6.1. *Throughout this section, we assume that $\alpha < -N$ and $\beta < -N$, so that the $Q_r(\xi; \alpha, \beta, N)$ satisfy the orthogonality relation (4.12). We consider the following limit:*

$$\epsilon := -\frac{\alpha + \beta}{\sqrt{\alpha\beta N}} \rightarrow +0.$$

We write

$$\alpha = \frac{\alpha_\epsilon}{\epsilon^2}, \quad \beta = \frac{\beta_\epsilon}{\epsilon^2}, \quad N = \frac{N_\epsilon}{\epsilon^2},$$

and assume further that

$$\lim_{\epsilon \rightarrow +0} \frac{N_\epsilon}{\alpha_\epsilon + \beta_\epsilon} = 0, \quad \lim_{\epsilon \rightarrow +0} \frac{\beta_\epsilon}{\alpha_\epsilon + \beta_\epsilon} = \rho \in [0, 1].$$

Remark 6.2. We do not require in Assumption 6.1 that $\alpha_\epsilon, \beta_\epsilon$, and N_ϵ are uniquely determined by ϵ . In other words, these are multi-valued functions of ϵ in general (for admissible values of ϵ), but their limit behaviors are uniformly governed by ϵ .

With reference to Assumption 6.1, observe that

$$\lim_{\epsilon \rightarrow +0} \alpha_\epsilon = \lim_{\epsilon \rightarrow +0} \alpha \epsilon^2 = \lim_{\epsilon \rightarrow +0} \frac{\alpha_\epsilon + \beta_\epsilon}{\beta_\epsilon} \cdot \frac{\alpha_\epsilon + \beta_\epsilon}{N_\epsilon} = -\infty.$$

Likewise, we have

$$\lim_{\epsilon \rightarrow +0} \beta_\epsilon = -\infty, \quad \lim_{\epsilon \rightarrow +0} N_\epsilon = \frac{1}{\rho(1-\rho)} \in [4, \infty].$$

We will work with the *normalized* (or *monic*) Hahn polynomials:

$$q_r(\xi) = q_r(\xi; \epsilon) = \frac{(\alpha + 1)_r (-N)_r}{(r + \alpha + \beta + 1)_r} Q_r(\xi; \alpha, \beta, N). \tag{6.1}$$

Their recurrence relation is given by (cf. [31, Section 1.5])

$$\xi q_r(\xi) = q_{r+1}(\xi) + (a_r + b_r)q_r(\xi) + a_{r-1}b_r q_{r-1}(\xi), \tag{6.2}$$

where $q_{-1}(\xi) := 0$, and

$$a_r = \frac{(r + \alpha + \beta + 1)(r + \alpha + 1)(N - r)}{(2r + \alpha + \beta + 1)(2r + \alpha + \beta + 2)},$$

$$b_r = \frac{r(r + \alpha + \beta + N + 1)(r + \beta)}{(2r + \alpha + \beta)(2r + \alpha + \beta + 1)}.$$

For convenience, let

$$\lambda_\epsilon = \sqrt{\frac{2(\alpha_\epsilon + \beta_\epsilon + N_\epsilon)}{\alpha_\epsilon + \beta_\epsilon}}.$$

Note that

$$\lim_{\epsilon \rightarrow +0} \lambda_\epsilon = \sqrt{2}. \tag{6.3}$$

Consider the polynomial $\tilde{q}_r(\eta; \epsilon)$ in the new indeterminate η defined by

$$\tilde{q}_r(\eta) = \tilde{q}_r(\eta; \epsilon) = q_r\left(\frac{\lambda_\epsilon \eta}{\epsilon} + \frac{\alpha_\epsilon N_\epsilon}{(\alpha_\epsilon + \beta_\epsilon)\epsilon^2}\right) \cdot \frac{\epsilon^r}{(\lambda_\epsilon)^r} \in \mathbb{R}[\eta].$$

Note that $\tilde{q}_r(\eta)$ is also monic with degree r in η . Then (6.2) becomes

$$\eta \tilde{q}_r(\eta) = \tilde{q}_{r+1}(\eta) + \frac{1}{\lambda_\epsilon} \left((a_r + b_r)\epsilon - \frac{\alpha_\epsilon N_\epsilon}{(\alpha_\epsilon + \beta_\epsilon)\epsilon} \right) \tilde{q}_r(\eta) + \frac{a_{r-1}b_r \epsilon^2}{(\lambda_\epsilon)^2} \tilde{q}_{r-1}(\eta). \tag{6.4}$$

It is a straightforward matter to show that

$$\frac{1}{\lambda_\epsilon} \left((a_r + b_r)\epsilon - \frac{\alpha_\epsilon N_\epsilon}{(\alpha_\epsilon + \beta_\epsilon)\epsilon} \right) = -(\mu_\epsilon + r\sigma_\epsilon)\epsilon + O(\epsilon^3), \tag{6.5}$$

$$\frac{a_{r-1}b_r \epsilon^2}{(\lambda_\epsilon)^2} = \frac{r}{2} + O(\epsilon^2), \tag{6.6}$$

where

$$\mu_\epsilon := \frac{(\alpha_\epsilon - \beta_\epsilon)N_\epsilon}{\lambda_\epsilon(\alpha_\epsilon + \beta_\epsilon)^2}, \quad \sigma_\epsilon := \frac{(\alpha_\epsilon - \beta_\epsilon)(\alpha_\epsilon + \beta_\epsilon + 2N_\epsilon)}{\lambda_\epsilon(\alpha_\epsilon + \beta_\epsilon)^2}$$

are convergent:

$$\lim_{\epsilon \rightarrow +0} \mu_\epsilon = 0, \quad \lim_{\epsilon \rightarrow +0} \sigma_\epsilon = \frac{1 - 2\rho}{\sqrt{2}}. \tag{6.7}$$

Recall the *Hermite polynomials* [31, Section 1.13]

$$H_r(\eta) = (2\eta)^r {}_2F_0 \left(\begin{matrix} -r/2, -(r-1)/2 \\ - \end{matrix} \middle| -\frac{1}{\eta^2} \right) \in \mathbb{R}[\eta] \quad (r = 0, 1, 2, \dots).$$

Their normalized recurrence relation is given by

$$\eta h_r(\eta) = h_{r+1}(\eta) + \frac{r}{2} h_{r-1}(\eta), \tag{6.8}$$

where

$$h_r(\eta) = \frac{H_r(\eta)}{2^r}, \tag{6.9}$$

and $h_{-1}(\eta) := 0$. We also note that

$$\frac{dh_r}{d\eta}(\eta) = r h_{r-1}(\eta), \tag{6.10}$$

and that

$$h_r(-\eta) = (-1)^r h_r(\eta). \tag{6.11}$$

Since $\tilde{q}_0(\eta) = h_0(\eta) = 1$, it follows from (6.4) – (6.8) that

$$\lim_{\epsilon \rightarrow +0} \tilde{q}_r(\eta; \epsilon) = h_r(\eta) \tag{6.12}$$

in the sense of coefficient-wise convergence.

We now set

$$\tilde{q}_r(\eta; 0) = h_r(\eta),$$

and discuss partial derivatives of $\tilde{q}_r(\eta; \epsilon)$ as a bivariate function of η and ϵ . First, it follows from (6.10) and (6.12) that

$$\lim_{\epsilon \rightarrow +0} \frac{\partial \tilde{q}_r}{\partial \eta}(\eta; \epsilon) = \frac{dh_r}{d\eta}(\eta) = r h_{r-1}(\eta). \tag{6.13}$$

Concerning the partial differentiability of $\tilde{q}_r(\eta; \epsilon)$ with respect to ϵ , it follows that

Lemma 6.3. *The function $\tilde{q}_r(\eta; \epsilon)$ is partially right differentiable with respect to ϵ at $(\eta, 0)$, and we have*

$$\frac{\partial \tilde{q}_r}{\partial \epsilon}(\eta; 0) = \frac{r(1-2\rho)}{3\sqrt{2}} ((r-1+\eta^2)h_{r-1}(\eta) - \eta h_r(\eta)).$$

Proof. Throughout the proof, we fix $\eta \in \mathbb{R}$ and set

$$\Delta_r(\epsilon) = \Delta_r(\eta; \epsilon) = \frac{\tilde{q}_r(\eta; \epsilon) - h_r(\eta)}{\epsilon}.$$

It follows from (6.4) – (6.8) and (6.12) that

$$\begin{aligned} \eta \Delta_r(\epsilon) &= \Delta_{r+1}(\epsilon) - (\mu_\epsilon + r\sigma_\epsilon) \tilde{q}_r(\eta; \epsilon) + \frac{r}{2} \Delta_{r-1}(\epsilon) + O(\epsilon) \\ &= \Delta_{r+1}(\epsilon) - r\sigma_0 h_r(\eta) + \frac{r}{2} \Delta_{r-1}(\epsilon) + o(1), \end{aligned} \tag{6.14}$$

where we set

$$\sigma_0 := \lim_{\epsilon \rightarrow +0} \sigma_\epsilon = \frac{1 - 2\rho}{\sqrt{2}}$$

for brevity. Since $\tilde{q}_0(\eta; \epsilon) = 1$, we have $\Delta_0(\epsilon) = 0$. Solving the recurrence (6.14) using this initial condition and (6.8), we routinely obtain

$$\Delta_r(\epsilon) = \frac{r(r-1)}{2} \sigma_0 h_{r-1}(\eta) + \frac{r(r-1)(r-2)}{12} \sigma_0 h_{r-3}(\eta) + o(1),$$

where $h_{-1}(\eta) = h_{-2}(\eta) = h_{-3}(\eta) := 0$. It follows that $\tilde{q}_r(\eta; \epsilon)$ is partially right differentiable with respect to ϵ at $(\eta, 0)$:

$$\begin{aligned} \frac{\partial \tilde{q}_r}{\partial \epsilon}(\eta; 0) &= \lim_{\epsilon \rightarrow +0} \Delta_r(\epsilon) \\ &= \frac{r(r-1)}{2} \sigma_0 h_{r-1}(\eta) + \frac{r(r-1)(r-2)}{12} \sigma_0 h_{r-3}(\eta). \end{aligned}$$

Finally, from (6.8) it follows that

$$\begin{aligned} \frac{\partial \tilde{q}_r}{\partial \epsilon}(\eta; 0) &= \frac{r(r-1)}{2} \sigma_0 h_{r-1}(\eta) + \frac{r(r-1)}{6} \sigma_0 (\eta h_{r-2}(\eta) - h_{r-1}(\eta)) \\ &= \frac{r(r-1)}{3} \sigma_0 h_{r-1}(\eta) + \frac{r}{3} \sigma_0 \eta (\eta h_{r-1}(\eta) - h_r(\eta)) \\ &= \frac{r\sigma_0}{3} ((r-1 + \eta^2) h_{r-1}(\eta) - \eta h_r(\eta)), \end{aligned}$$

as desired. □

Proposition 6.4. Recall Assumption 6.1. Fix a positive integer e , and let

$$\begin{aligned} \xi_{-\lfloor e/2 \rfloor} &< \dots < \xi_{-1} < (\xi_0) < \xi_1 < \dots < \xi_{\lfloor e/2 \rfloor}, \\ \eta_{-\lfloor e/2 \rfloor} &< \dots < \eta_{-1} < (\eta_0) < \eta_1 < \dots < \eta_{\lfloor e/2 \rfloor} \end{aligned}$$

be the zeros of $q_e(\xi; \epsilon)$ and $h_e(\eta)$ from (6.1) and (6.9), respectively, where ξ_0 and η_0 appear only when e is odd. Then ξ_i satisfies

$$\lim_{\epsilon \rightarrow +0} \left(\xi_i - \frac{\lambda_\epsilon \eta_i}{\epsilon} - \frac{\alpha_\epsilon N_\epsilon}{(\alpha_\epsilon + \beta_\epsilon) \epsilon^2} \right) = \frac{2\rho - 1}{3} (e - 1 + (\eta_i)^2)$$

as a function of ϵ , for $i = -\lfloor e/2 \rfloor, \dots, -1, (0), 1, \dots, \lfloor e/2 \rfloor$.

Proof. Define τ_i by

$$\xi_i = \frac{\lambda_\epsilon (\eta_i + \tau_i)}{\epsilon} + \frac{\alpha_\epsilon N_\epsilon}{(\alpha_\epsilon + \beta_\epsilon) \epsilon^2},$$

so that $\eta_i + \tau_i$ is a zero of $\tilde{q}_e(\eta; \epsilon)$. Then, from (6.12) it follows that

$$\lim_{\epsilon \rightarrow +0} \tau_i = 0. \tag{6.15}$$

For the moment, fix i . Then we have

$$0 = \tilde{q}_e(\eta_i + \tau_i; \epsilon) = \tilde{q}_e(\eta_i; \epsilon) + \frac{\partial \tilde{q}_e}{\partial \eta}(\eta_i + \theta \tau_i; \epsilon) \tau_i$$

for some $\theta \in (0, 1)$ depending on ϵ . Hence, from (6.13), (6.15), Lemma 6.3, and since

$$\tilde{q}_e(\eta_i; 0) = h_e(\eta_i) = 0,$$

it follows that

$$\begin{aligned} \lim_{\epsilon \rightarrow +0} \frac{\tau_i}{\epsilon} &= -\frac{1}{eh_{e-1}(\eta_i)} \lim_{\epsilon \rightarrow +0} \frac{\tilde{q}_e(\eta_i; \epsilon)}{\epsilon} \\ &= -\frac{1}{eh_{e-1}(\eta_i)} \frac{\partial \tilde{q}_e}{\partial \epsilon}(\eta_i; 0) \\ &= \frac{2\rho - 1}{3\sqrt{2}}(e - 1 + (\eta_i)^2), \end{aligned}$$

where we note that $h_e(\eta)$ and $h_{e-1}(\eta)$ have no common zero by the general theory of orthogonal polynomials; see, e.g., [42, Theorem 3.3.2]. By (6.3), we have

$$\lim_{\epsilon \rightarrow +0} \left(\xi_i - \frac{\lambda_\epsilon \eta_i}{\epsilon} - \frac{\alpha_\epsilon N_\epsilon}{(\alpha_\epsilon + \beta_\epsilon) \epsilon^2} \right) = \lim_{\epsilon \rightarrow +0} \frac{\lambda_\epsilon \tau_i}{\epsilon} = \frac{2\rho - 1}{3}(e - 1 + (\eta_i)^2).$$

This completes the proof. □

The following is part of the estimates on the zeros of $h_e(\eta)$ used in [1].³

Proposition 6.5 ([1, Proposition 13]). *Fix a positive integer e , and let the η_i be as in Proposition 6.4. Then $\eta_{-i} = -\eta_i$ for all i . Moreover, the following hold:*

1. *If e is odd and $e \geq 5$, then $\eta_0 = 0$ and $(\eta_1)^2 < 3/2$.*
2. *If e is even and $e \geq 8$, then $(\eta_2)^2 - (\eta_1)^2 < 3/2$.*

Proof. That $\eta_{-i} = -\eta_i$ is immediate from (6.11). We now write $\eta_i = \eta_i^e$ to compare these zeros for different values of e . Then, as an application of Sturm’s method, it follows that

$$\sqrt{2e + 1} \eta_i^e < \sqrt{2e' + 1} \eta_i^{e'} \quad (i = 1, 2, \dots, \lfloor e'/2 \rfloor),$$

whenever $e' < e$ and $e' \equiv e \pmod{2}$; see the comments preceding (6.31.19) in [42]. Since

$$h_3(\eta) = \eta^3 - \frac{3}{2}\eta, \quad h_4(\eta) = \eta^4 - 3\eta^2 + \frac{3}{4},$$

we have

$$\eta_1^3 = \sqrt{\frac{3}{2}}, \quad \eta_2^4 = \sqrt{\frac{3 + \sqrt{6}}{2}}.$$

Hence, for odd $e \geq 5$ we have

$$(\eta_1^e)^2 < \frac{7}{2e + 1} (\eta_1^3)^2 = \frac{21}{4e + 2} < \frac{3}{2},$$

and for even $e \geq 8$ we have

$$(\eta_2^e)^2 - (\eta_1^e)^2 < (\eta_2^4)^2 < \frac{9}{2e + 1} (\eta_2^4)^2 = \frac{27 + 9\sqrt{6}}{4e + 2} < \frac{3}{2},$$

as desired. □

³Bannai [1] worked with the polynomial $\sqrt{2^e} h_e(\eta/\sqrt{2})$. We may remark that the upper bounds $\sqrt{3}$ mentioned in Proposition 13 (i) and (ii) in [1] should both be 3. See also [22, Proposition 2.4].

7 A finiteness result for tight relative $2e$ -designs on two shells in \mathcal{Q}_n

In this section, we prove that

Theorem 7.1. *For any $\delta \in (0, 1/2)$, there exists $e_0 = e_0(\delta) > 0$ with the property that, for every given integer $e \geq e_0$ and each constant $c > 0$, there are only finitely many tight relative $2e$ -designs (Y, ω) (up to scalar multiples of ω) supported on two shells $X_\ell \sqcup X_m$ in \mathcal{Q}_n satisfying Assumption 5.2 such that*

$$\ell < c \cdot n^\delta. \tag{7.1}$$

Our proof is an application of Bannai’s method from [1]. We will use the following result, which is a variation of [40, Satz I]:

Proposition 7.2. *For any $\vartheta > 0$ and $\delta \in (0, 1/\vartheta)$, there exists $\mathfrak{k}_0 = \mathfrak{k}_0(\vartheta, \delta) > 0$ such that the following holds for every given integer $\mathfrak{k} \geq \mathfrak{k}_0$ and each constant $c > 0$: for all but finitely many pairs $(\mathfrak{a}, \mathfrak{b})$ of positive integers with*

$$\mathfrak{b} < c \cdot \mathfrak{a}^\delta,$$

the product of \mathfrak{k} consecutive odd integers

$$(2\mathfrak{a} + 1)(2\mathfrak{a} + 3) \cdots (2\mathfrak{a} + 2\mathfrak{k} - 1)$$

has a prime factor which is greater than $2\mathfrak{k} + 1$ and whose exponent in this product is greater than that in

$$(\mathfrak{b} + 1)(\mathfrak{b} + 2) \cdots (\mathfrak{b} + \lfloor \vartheta \mathfrak{k} \rfloor).$$

The proof of Proposition 7.2 will be deferred to the appendix.

We will establish Theorem 7.1 by contradiction:

Assumption 7.3. *We fix $\delta \in (0, 1/2)$. Let $\mathfrak{k}_0 = \mathfrak{k}_0(2, \delta) > 0$ be as in Proposition 7.2 (applied to $\vartheta = 2$), and set*

$$e_0 = e_0(\delta) = \max\{2\mathfrak{k}_0, 8\}.$$

We also fix a positive integer $e \geq e_0$ and a constant $c > 0$. Throughout the proof, we assume that there exist infinitely many tight relative $2e$ -designs (Y, ω) in question.

Let Θ denote the set of triples $(\ell, m, n) \in \mathbb{N}^3$ taken by those (Y, ω) in Assumption 7.3. Recall from Proposition 5.4 that ω is uniquely determined by Y up to a scalar multiple. Moreover, for each $(\ell, m, n) \in \Theta$ there are only finitely many choices for Y . Hence we have

$$|\Theta| = \infty. \tag{7.2}$$

For the moment, we fix $(\ell, m, n) \in \Theta$ and consider the polynomial $\psi_e^{\ell, m}(\xi)$ (which also depends on n) from (5.36). We recall that

$$\psi_e^{\ell, m}(\xi) = Q_e(\xi; \alpha, \beta, N),$$

where

$$\alpha = m - n - 1, \quad \beta = -m - 1, \quad N = \ell. \tag{7.3}$$

We note that $\alpha, \beta < -N$ in view of Assumption 5.2. By Theorem 5.3(ii), if we let

$$\xi_{-\lfloor e/2 \rfloor} < \cdots < \xi_{-1} < (\xi_0) < \xi_1 < \cdots < \xi_{\lfloor e/2 \rfloor} \tag{7.4}$$

denote the zeros of $\psi_e^{\ell, m}(\xi)$ (cf. Proposition 6.4), then we have

$$\xi_i \in \{0, 1, \dots, \ell\} \quad \text{for all } i. \tag{7.5}$$

We also rewrite $\psi_e^{\ell, m}(\xi)$ as follows:

$$\psi_e^{\ell, m}(\xi) = \sum_{i=0}^e s_{e-i} (-1)^i (-\xi)_i,$$

where

$$s_{e-i} = \binom{e}{i} \frac{(e-n-1)_i}{(m-n)_i (-\ell)_i} \quad (0 \leq i \leq e).$$

From (7.5) it follows that the polynomial $\psi_e^{\ell, m}(\xi)/s_0$ is monic and integral:

$$\frac{\psi_e^{\ell, m}(\xi)}{s_0} = \sum_{i=0}^e \frac{s_{e-i}}{s_0} (-1)^i (-\xi)_i = (\xi - \xi_{-\lfloor e/2 \rfloor}) \cdots (\xi - \xi_{\lfloor e/2 \rfloor}) \in \mathbb{Z}[\xi], \tag{7.6}$$

where the factor $(\xi - \xi_0)$ appears only when e is odd. Since $(-1)^i (-\xi)_i$ is also monic and integral, and has degree i for $0 \leq i \leq e$, it follows that

$$\frac{s_i}{s_0} = (-1)^i \binom{e}{i} \frac{(n-m-e+1)_i (\ell-e+1)_i}{(n-2e+2)_i} \in \mathbb{Z} \setminus \{0\} \quad (0 \leq i \leq e), \tag{7.7}$$

where that these coefficients are non-zero follows from Assumption 5.2.

We now consider the map $f : \Theta \rightarrow [0, 1]^2$ defined by

$$f(\ell, m, n) = \left(\frac{\ell}{n}, \frac{m}{n} \right) \in [0, 1]^2 \quad ((\ell, m, n) \in \Theta).$$

Recall (7.2). Moreover, from (7.1) it follows that

$$|f^{-1}(a, b)| < \infty \quad ((a, b) \in [0, 1]^2). \tag{7.8}$$

Hence it follows that

$$|f(\Theta)| = \infty,$$

so that $f(\Theta)$ has at least one accumulation point in $[0, 1]^2$. Again by (7.1), such an accumulation point must be of the form

$$(0, \rho) \in [0, 1]^2.$$

We next show that the parameters α, β , and N from (7.3) satisfy Assumption 6.1 when $f(\ell, m, n) \rightarrow (0, \rho)$.

Claim 7.4. $\ell, m, n - m \rightarrow \infty$ as $f(\ell, m, n) \rightarrow (0, \rho)$.

Proof. Since $m, n - m \geq \ell$ by Assumption 5.2, it suffices to show that $\ell \rightarrow \infty$. Suppose the contrary, i.e., that there is a sequence (ℓ_k, m_k, n_k) ($k \in \mathbb{N}$) of distinct elements of Θ such that

$$\lim_{k \rightarrow \infty} f(\ell_k, m_k, n_k) = (0, \rho), \quad \sup_k \ell_k < \infty.$$

Since the ℓ_k are bounded, it follows from (7.5) and (7.6) that there are only finitely many choices for $\psi_e^{\ell, m}(\xi)/s_0$ when (ℓ, m, n) ranges over this sequence. In particular, there are only finitely many choices for each of the coefficients s_1/s_0 and s_2/s_0 , and hence the same is true (cf. (7.7)) for each of

$$\frac{n - m - e + 1}{n - 2e + 2}, \quad \frac{n - m - e + 2}{n - 2e + 3}.$$

However, it is immediate to see that these distinct scalars in turn determine n and m uniquely, from which it follows that the n_k are bounded, a contradiction. \square

Claim 7.5. $\ell m(n - m)/n^2 \rightarrow \infty$ as $f(\ell, m, n) \rightarrow (0, \rho)$.

Proof. If $0 < \rho < 1$ then the result follows from Claim 7.4 and since

$$\frac{m(n - m)}{n^2} \rightarrow \rho(1 - \rho) > 0.$$

Suppose next that $\rho = 1$. Suppose moreover that there is a sequence (ℓ_k, m_k, n_k) ($k \in \mathbb{N}$) of distinct elements of Θ such that

$$\lim_{k \rightarrow \infty} f(\ell_k, m_k, n_k) = (0, 1), \quad \sup_k \frac{\ell_k m_k (n_k - m_k)}{(n_k)^2} < \infty.$$

Since $m_k/n_k \rightarrow 1$, we then have

$$\sup_k \frac{\ell_k (n_k - m_k)}{n_k} < \infty.$$

Let

$$r_k = \frac{(n_k - m_k - e + 1)(\ell_k - e + 1)}{n_k - 2e + 2}, \quad t_k = \frac{(n_k - m_k - e + 2)(\ell_k - e + 2)}{n_k - 2e + 3}.$$

Then the r_k and the t_k are bounded since

$$r_k \approx t_k \approx \frac{\ell_k (n_k - m_k)}{n_k}$$

by Claim 7.4. From (7.7) it follows that s_1/s_0 and s_2/s_0 are bounded as well, and hence take only finitely many non-zero integral values when (ℓ, m, n) ranges over this sequence. It follows that the r_k and the t_k can assume only finitely many values, and then since $r_k \approx t_k$ we must have $r_k = t_k$ for sufficiently large k . However, it is again immediate to see that $r_k \neq t_k$ for every $k \in \mathbb{N}$, and hence this is absurd. It follows that the result holds when $\rho = 1$.

Finally, suppose that $\rho = 0$. For every $(\ell, m, n) \in \Theta$ we have

$$e \frac{(m - e + 1)(\ell - e + 1)}{n - 2e + 2} = \frac{s_1}{s_0} + e(\ell - e + 1),$$

$$\binom{e}{2} \frac{(m-e+1)_2(\ell-e+1)_2}{(n-2e+2)_2} = \frac{s_2}{s_0} + (e-1)(\ell-e+2) \frac{s_1}{s_0} + \binom{e}{2} (\ell-e+1)_2.$$

From (7.7) and Assumption 5.2 it follows that these scalars are non-zero integers. By the same argument as above, but working with these two scalars instead of s_1/s_0 and s_2/s_0 , we conclude that the result holds in this case as well. \square

By Claims 7.4 and 7.5, it follows that the parameters α, β , and N from (7.3) satisfy Assumption 6.1 when $f(\ell, m, n) \rightarrow (0, \rho)$, since

$$-\frac{\alpha + \beta}{\sqrt{\alpha\beta N}} \approx \frac{n}{\sqrt{\ell m(n-m)}}, \quad \frac{N}{\alpha + \beta} \approx -\frac{\ell}{n}, \quad \frac{\beta}{\alpha + \beta} \approx \frac{m}{n}.$$

Note that the scalar ρ in Assumption 6.1 agrees with the one used here in this case. Hence we are now in the position to apply the results of the previous section to $\psi_e^{\ell, m}(\xi)$, which is the Hahn polynomial having these parameters.

Claim 7.6. *We have $\rho = 1/2$. In particular, $(0, 1/2)$ is a unique accumulation point of $f(\Theta)$. Moreover, we have $n = 2m$ for all but finitely many $(\ell, m, n) \in \Theta$.*

Proof. Let the ξ_i be as in (7.4). Then from Propositions 6.4 and 6.5 it follows that

$$\xi_i + \xi_{-i} - \xi_j - \xi_{-j} \rightarrow \frac{4\rho - 2}{3} ((\eta_i)^2 - (\eta_j)^2) \quad \text{for all } i, j, \tag{7.9}$$

as $f(\ell, m, n) \rightarrow (0, \rho)$, where the η_i are the zeros of the monic Hermite polynomial $h_e(\eta)$ from (6.9) as in Proposition 6.4. Recall that $e \geq 8$ by Assumption 7.3. Set $(i, j) = (1, 0)$ in (7.9) if e is odd, and $(i, j) = (2, 1)$ if e is even. Then, since

$$\left| \frac{4\rho - 2}{3} \right| \leq \frac{2}{3},$$

it follows from Proposition 6.5 that the RHS in (7.9) lies in the open interval $(-1, 1)$. However, the LHS in (7.9) is always an integer by (7.5), so that this is possible only when the RHS equals zero, i.e., $\rho = 1/2$. In particular, we have shown that $(0, 1/2)$ is a unique accumulation point of $f(\Theta)$.

Again by (7.5) and (7.9), we then have

$$\xi_i + \xi_{-i} = \xi_j + \xi_{-j} \quad \text{for all } i, j,$$

provided that $f(\ell, m, n)$ is sufficiently close to $(0, 1/2)$. By the uniqueness of the accumulation point and (7.8), this last condition on $f(\ell, m, n)$ can be rephrased as “for all but finitely many $(\ell, m, n) \in \Theta$.” Now, let $\tilde{\xi}$ be the average of the zeros ξ_i of $\psi_e^{\ell, m}(\xi)$. Then the above identity means that the ξ_i are symmetric with respect to $\tilde{\xi}$. Hence, if we write

$$\frac{\psi_e^{\ell, m}(\xi)}{s_0} = \sum_{i=0}^e w_{e-i} (\xi - \tilde{\xi})^i,$$

then we have

$$w_{2i-1} = 0 \quad (1 \leq i \leq \lceil e/2 \rceil)$$

for all but finitely many $(\ell, m, n) \in \Theta$. On the other hand, using (7.6) and (7.7), we routinely obtain

$$w_3 = \binom{e}{3}(n - 2\ell)(n - 2m) \times \frac{(\ell - e + 1)(m - e + 1)(n - \ell - e + 1)(n - m - e + 1)}{(n - 2e + 2)^3(n - 2e + 3)(n - 2e + 4)}.$$

Hence, by Assumption 5.2, that $w_3 = 0$ forces $n = 2m$. The claim is proved. □

By virtue of Claim 7.6, we may now assume without loss of generality that

$$n = 2m \quad ((\ell, m, n) \in \Theta),$$

by discarding a finite number of exceptions. Set

$$\mathfrak{k} = \left\lfloor \frac{e}{2} \right\rfloor,$$

and let c' be a constant such that $c' > 2^\delta c$. Note that

$$\mathfrak{k} \geq \mathfrak{k}_0 = \mathfrak{k}_0(2, \delta)$$

by Assumption 7.3. Let $(\ell, m, 2m) \in \Theta$. We have

$$c \cdot (2m)^\delta < c' \cdot (m - e + 1)^\delta$$

provided that m is large. Hence it follows from Proposition 7.2 (applied to $\vartheta = 2$) and (7.1) that if m is sufficiently large then there is a prime $p > 2\mathfrak{k} + 1$ such that

$$\nu_p((2m - 2e + 3)(2m - 2e + 5) \cdots (2m - 2e + 2\mathfrak{k} + 1)) > \nu_p((\ell - e + 1)_{2\mathfrak{k}}),$$

where $\nu_p(n)$ denotes the exponent of p in n . Assuming that this is the case, let i ($1 \leq i \leq \mathfrak{k}$) be such that

$$\nu_p(2m - 2e + 2i + 1) > 0.$$

Observe that i is unique since $p > 2\mathfrak{k} + 1$, so that we have

$$\nu_p(2m - 2e + 2i + 1) > \nu_p((\ell - e + 1)_{2\mathfrak{k}}).$$

Moreover, we have

$$\gcd(2m - 2e + 2i + 1, m - e + i + j) = \gcd(2j - 1, m - e + i + j) < p$$

for $1 \leq j \leq i$, from which it follows that

$$\nu_p((m - e + i + 1)_i) = 0.$$

By these comments and since

$$2i \leq e < p,$$

it follows from (7.7) (with $n = 2m$) that


$$\nu_p\left(\frac{s_{2i}}{s_0}\right) = \nu_p\left(\frac{(m - e + 1)_{2i}(\ell - e + 1)_{2i}}{(2m - 2e + 2)_{2i}}\right)$$


$$= \nu_p \left(\frac{(m-e+i+1)_i (\ell-e+1)_{2i}}{2^i (2m-2e+3)(2m-2e+5) \cdots (2m-2e+2i+1)} \right) < 0.$$

However, this contradicts the fact that s_{2i}/s_0 is a non-zero integer. Hence we now conclude that Θ must be finite.

The proof of Theorem 7.1 is complete.

ORCID iDs

Hajime Tanaka  <https://orcid.org/0000-0002-5958-0375>

Yan Zhu  <https://orcid.org/0000-0001-7164-9198>

References

- [1] E. Bannai, On tight designs, *Q. J. Math. Oxford Ser. (2)* **28** (1977), 433–448, doi:10.1093/qmath/28.4.433.
- [2] E. Bannai and E. Bannai, Euclidean designs and coherent configurations, in: *Combinatorics and Graphs*, Amer. Math. Soc., Providence, RI, volume 531 of *Contemp. Math.*, pp. 59–93, 2010, doi:10.1090/conm/531/10457.
- [3] E. Bannai and E. Bannai, Remarks on the concepts of t -designs, *J. Appl. Math. Comput.* **40** (2012), 195–207, doi:10.1007/s12190-012-0544-1.
- [4] E. Bannai and E. Bannai, Tight t -designs on two concentric spheres, *Mosc. J. Comb. Number Theory* **4** (2014), 52–77, <http://mjcnt.phystech.edu/en/article.php?id=78>.
- [5] E. Bannai, E. Bannai and H. Bannai, On the existence of tight relative 2-designs on binary Hamming association schemes, *Discrete Math.* **314** (2014), 17–37, doi:10.1016/j.disc.2013.09.013.
- [6] E. Bannai, E. Bannai, S. Suda and H. Tanaka, On relative t -designs in polynomial association schemes, *Electron. J. Comb.* **22** (2015), Paper 4.47, 17, doi:10.37236/4889.
- [7] E. Bannai, E. Bannai, H. Tanaka and Y. Zhu, Design theory from the viewpoint of algebraic combinatorics, *Graphs Comb.* **33** (2017), 1–41, doi:10.1007/s00373-016-1739-2.
- [8] E. Bannai, E. Bannai and Y. Zhu, A survey on tight Euclidean t -designs and tight relative t -designs in certain association schemes, *Proc. Steklov Inst. Math.* **288** (2015), 189–202, doi:10.1134/s0081543815010149.
- [9] E. Bannai, E. Bannai and Y. Zhu, Relative t -designs in binary Hamming association scheme $H(n, 2)$, *Des. Codes Cryptogr.* **84** (2017), 23–53, doi:10.1007/s10623-016-0200-0.
- [10] E. Bannai and T. Ito, *Algebraic Combinatorics. I: Association Schemes*, The Benjamin/Cummings Publishing Co., Inc., Menlo Park, CA, 1984.
- [11] E. Bannai and Y. Zhu, Tight t -designs on one shell of Johnson association schemes, *Eur. J. Comb.* **80** (2019), 23–36, doi:10.1016/j.ejc.2018.02.024.
- [12] A. E. Brouwer, A. M. Cohen and A. Neumaier, *Distance-Regular Graphs*, volume 18 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]*, Springer-Verlag, Berlin, 1989, doi:10.1007/978-3-642-74341-2.
- [13] C. J. Colbourn and J. H. Dinitz (eds.), *Handbook of Combinatorial Designs*, Discrete Mathematics and its Applications (Boca Raton), Chapman & Hall/CRC, Boca Raton, FL, 2nd edition, 2007, doi:10.1201/9781420010541.

- [14] C. W. Curtis and I. Reiner, *Methods of Representation Theory. Vol. I*, Wiley Classics Library, John Wiley & Sons, Inc., New York, 1990, with applications to finite groups and orders, Reprint of the 1981 original, A Wiley-Interscience Publication, https://books.google.si/books?id=HWm_QgAACAAJ.
- [15] E. R. van Dam, J. H. Koolen and H. Tanaka, Distance-regular graphs, *Electron. J. Comb.* **DS22** (2016), Paper No. DS22, 156, doi:10.37236/4925.
- [16] P. Delsarte, An algebraic approach to the association schemes of coding theory, *Philips Res. Rep. Suppl.* (1973), vi+97, <https://users.wpi.edu/~martin/RESEARCH/>.
- [17] P. Delsarte, Association schemes and t -designs in regular semilattices, *J. Comb. Theory Ser. A* **20** (1976), 230–243, doi:10.1016/0097-3165(76)90017-0.
- [18] P. Delsarte, Pairs of vectors in the space of an association scheme, *Philips Res. Rep.* **32** (1977), 373–411.
- [19] P. Delsarte, Hahn polynomials, discrete harmonics, and t -designs, *SIAM J. Appl. Math.* **34** (1978), 157–166, doi:10.1137/0134012.
- [20] P. Delsarte, J. M. Goethals and J. J. Seidel, Spherical codes and designs, *Geom. Dedicata* **6** (1977), 363–388, doi:10.1007/bf03187604.
- [21] P. Delsarte and J. J. Seidel, Fisher type inequalities for Euclidean t -designs, *Linear Algebra Its Appl.* **114/115** (1989), 213–230, doi:10.1016/0024-3795(89)90462-x.
- [22] P. Dukes and J. Short-Gershman, Nonexistence results for tight block designs, *J. Algebraic Comb.* **38** (2013), 103–119, doi:10.1007/s10801-012-0395-8.
- [23] A. L. Gavrilyuk, J. Vidali and J. S. Williford, On few-class Q -polynomial association schemes: feasible parameters and nonexistence results, *Ars Math. Contemp.* **20** (2021), 103–127, doi:10.26493/1855-3974.2101.b76.
- [24] D. Gijswijt, Matrix algebras and semidefinite programming techniques for codes, 2005, arXiv:1007.0906 [math.CO].
- [25] D. Gijswijt, A. Schrijver and H. Tanaka, New upper bounds for nonbinary codes based on the Terwilliger algebra and semidefinite programming, *J. Comb. Theory Ser. A* **113** (2006), 1719–1731, doi:10.1016/j.jcta.2006.03.010.
- [26] J. T. Go, The Terwilliger algebra of the hypercube, *Eur. J. Comb.* **23** (2002), 399–429, doi:10.1006/eujc.2000.0514.
- [27] C. D. Godsil, *Algebraic Combinatorics*, Chapman and Hall Mathematics Series, Chapman & Hall, New York, 1993, <https://books.google.si/books?id=eADtLNckkIMC>.
- [28] S. Kageyama, A property of T -wise balanced designs, *Ars Comb.* **31** (1991), 237–238.
- [29] B. G. Kodalen, Cometric association schemes, 2019, arXiv:1905.06959 [math.CO].
- [30] R. Koekoek, P. A. Lesky and R. F. Swarttouw, *Hypergeometric Orthogonal Polynomials and Their q -Analogues*, Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2010, doi:10.1007/978-3-642-05014-5.
- [31] R. Koekoek and R. F. Swarttouw, *The Askey scheme of hypergeometric orthogonal polynomials and its q -analog*, Delft University of Technology, 1998, {<http://aw.twi.tudelft.nl/~koekoek/askey.html>}.
- [32] Z. Li, E. Bannai and E. Bannai, Tight relative 2- and 4-designs on binary Hamming association schemes, *Graphs Comb.* **30** (2014), 203–227, doi:10.1007/s00373-012-1252-1.
- [33] W. J. Martin and H. Tanaka, Commutative association schemes, *Eur. J. Comb.* **30** (2009), 1497–1525, doi:10.1016/j.ejc.2008.11.001.

- [34] A. Munemasa, An analogue of t -designs in the association schemes of alternating bilinear forms, *Graphs Comb.* **2** (1986), 259–267, doi:10.1007/bf01788100.
- [35] A. Neumaier and J. J. Seidel, Discrete measures for spherical designs, eutactic stars and lattices, *Nederl. Akad. Wetensch. Indag. Math.* **50** (1988), 321–334, doi:10.1016/s1385-7258(88)80011-8.
- [36] C. Peterson, On tight 6-designs, *Osaka Math. J.* **14** (1977), 417–435, doi:10.18910/3990.
- [37] J. B. Rosser and L. Schoenfeld, Approximate formulas for some functions of prime numbers, *Illinois J. Math.* **6** (1962), 64–94, doi:10.1215/ijm/1255631807.
- [38] M. Sawa, M. Hirao and S. Kageyama, *Euclidean Design Theory*, SpringerBriefs in Statistics, Springer, Singapore, 2019, doi:10.1007/978-981-13-8075-4.
- [39] A. Schrijver, New code upper bounds from the Terwilliger algebra and semidefinite programming, *IEEE Trans. Inform. Theory* **51** (2005), 2859–2866, doi:10.1109/tit.2005.851748.
- [40] I. Schur, *Einige Sätze über Primzahlen: mit Anwendungen auf Irreduzibilitätsfragen*, Sitzungsberichte der Preussischen Akademie der Wissenschaften. Physikalisch-mathematische Klasse, Akademie der Wissenschaften in Kommission bei Walter de Gruyter, 1929, <https://books.google.si/books?id=wE3kAAAAAMAAJ>.
- [41] D. Stanton, t -designs in classical association schemes, *Graphs Comb.* **2** (1986), 283–286, doi:10.1007/bf01788103.
- [42] G. Szegő, *Orthogonal Polynomials*, American Mathematical Society Colloquium Publications, Vol. XXIII, American Mathematical Society, Providence, R.I., 4th edition, 1975, <https://people.math.osu.edu/nevai.1/SZEGO/>.
- [43] Y.-Y. Tan, Y.-Z. Fan, T. Ito and X. Liang, The Terwilliger algebra of the Johnson scheme $J(N, D)$ revisited from the viewpoint of group representations, *Eur. J. Comb.* **80** (2019), 157–171, doi:10.1016/j.ejc.2018.02.029.
- [44] H. Tanaka, New proofs of the Assmus-Mattson theorem based on the Terwilliger algebra, *Eur. J. Comb.* **30** (2009), 736–746, doi:10.1016/j.ejc.2008.07.018.
- [45] H. Tanaka, A note on the span of hadamard products of vectors, *Linear Algebra Its Appl.* **430** (2009), 865–867, doi:10.1016/j.laa.2008.09.010.
- [46] P. Terwilliger, The subconstituent algebra of an association scheme. I, *J. Algebr. Comb.* **1** (1992), 363–388, doi:10.1023/a:1022494701663.
- [47] P. Terwilliger, The subconstituent algebra of an association scheme. II, *J. Algebr. Comb.* **2** (1993), 73–103, doi:10.1023/a:1022480715311.
- [48] P. Terwilliger, The subconstituent algebra of an association scheme. III, *J. Algebr. Comb.* **2** (1993), 177–210, doi:10.1023/a:1022415825656.
- [49] F. Vallentin, Symmetry in semidefinite programs, *Linear Algebra Appl.* **430** (2009), 360–369, doi:10.1016/j.laa.2008.07.025.
- [50] D. R. Woodall, Square λ -linked designs, *Proc. London Math. Soc. (3)* **20** (1970), 669–687, doi:10.1112/plms/s3-20.4.669.
- [51] Z. Xiang, A Fisher type inequality for weighted regular t -wise balanced designs, *J. Comb. Theory Ser. A* **119** (2012), 1523–1527, doi:10.1016/j.jcta.2012.04.008.
- [52] Z. Xiang, Nonexistence of nontrivial tight 8-designs, *J. Algebr. Comb.* **47** (2018), 301–318, doi:10.1007/s10801-017-0776-0.
- [53] H. Yue, B. Hou and S. Gao, Note on the tight relative 2-designs on $H(n, 2)$, *Discrete Math.* **338** (2015), 196–208, doi:10.1016/j.disc.2014.09.002.

- [54] Y. Zhu, E. Bannai and E. Bannai, Tight relative 2-designs on two shells in Johnson association schemes, *Discrete Math.* **339** (2016), 957–973, doi:10.1016/j.disc.2015.10.024.

Appendix A Proof of Proposition 7.2

Our proof of Proposition 7.2 is a slight modification of (the first part of) that of [40, Satz I].

For a positive integer n , let

$$\chi_n = 1 \cdot 3 \cdot 5 \cdots (2n - 1) = \frac{(2n)!}{2^n n!}.$$

Observe that the exponent $\nu_p(\chi_n)$ of an odd prime p in χ_n is given by

$$\nu_p(\chi_n) = \sum_{i=1}^{\lfloor \log_p(2n) \rfloor} \left(\left\lfloor \frac{2n}{p^i} \right\rfloor - \left\lfloor \frac{n}{p^i} \right\rfloor \right) = \sum_{i=1}^{\lfloor \log_p(2n) \rfloor} \left\lfloor \frac{n}{p^i} + \frac{1}{2} \right\rfloor, \tag{A.1}$$

where we have used

$$\lfloor \xi \rfloor + \left\lfloor \xi + \frac{1}{2} \right\rfloor = \lfloor 2\xi \rfloor \quad (\xi \in \mathbb{R}).$$

Now, let (a, b) be a pair of positive integers with

$$b < c \cdot a^\delta, \tag{A.2}$$

which does not satisfy the desired property about a prime factor; in other words,

$$\nu_p(\chi_{a+\ell}) - \nu_p(\chi_a) \leq \nu_p((b + \lfloor \vartheta \ell \rfloor)!) - \nu_p(b!) \quad \text{if } p > 2\ell + 1. \tag{A.3}$$

Our aim is to show that a is bounded in terms of ϑ, δ, c , and ℓ , and hence so is b by (A.2), from which it follows that there are only finitely many such pairs. (We will specify $\ell_0 = \ell_0(\vartheta, \delta)$ at the end of the proof.) To this end, we may assume for example that

$$a > \ell, \quad c \cdot a^\delta > \ell + 1. \tag{A.4}$$

Without loss of generality, we may also assume that

$$b > \ell, \tag{A.5}$$

for otherwise the pair $(a, \ell + 1)$ would also satisfy (A.2) and (A.3).

Let

$$s = \frac{\chi_{a+\ell}}{\chi_\ell \chi_a} \cdot \frac{b!}{(b + \lfloor \vartheta \ell \rfloor)!}.$$

Then from (A.3) it follows that

$$s \leq \prod_{3 \leq p \leq 2\ell+1} p^{\nu_p(s)}, \tag{A.6}$$

where the product in the RHS is over the odd primes $p \leq 2\ell + 1$, and where

$$\nu_p(s) = \nu_p(\chi_{a+\ell}) - \nu_p(\chi_\ell) - \nu_p(\chi_a) + \nu_p(b!) - \nu_p((b + \lfloor \vartheta \ell \rfloor)!).$$

By (A.1), for every odd prime p we have

$$\nu_p(s) \leq \nu_p(\chi_{a+\ell}) - \nu_p(\chi_\ell) - \nu_p(\chi_a)$$

$$= \sum_{i=1}^{\lfloor \log_p(2\alpha+2\ell) \rfloor} \left(\left\lfloor \frac{\alpha + \ell}{p^i} + \frac{1}{2} \right\rfloor - \left\lfloor \frac{\ell}{p^i} + \frac{1}{2} \right\rfloor - \left\lfloor \frac{\alpha}{p^i} + \frac{1}{2} \right\rfloor \right).$$

Note that

$$\left\lfloor \xi + \eta + \frac{1}{2} \right\rfloor - \left\lfloor \xi + \frac{1}{2} \right\rfloor - \left\lfloor \eta + \frac{1}{2} \right\rfloor \in \{-1, 0, 1\} \quad (\xi, \eta \in \mathbb{R}).$$

Hence it follows that

$$\nu_p(\mathfrak{s}) \leq \log_p(2\alpha + 2\ell) = \frac{\ln(2\alpha + 2\ell)}{\ln p} \tag{A.7}$$

for every odd prime p . From (A.6) and (A.7) it follows that

$$\ln \mathfrak{s} \leq (\pi(2\ell + 1) - 1) \ln(2\alpha + 2\ell), \tag{A.8}$$

where $\pi(n)$ denotes the number of primes at most n .

On the other hand, we have

$$\mathfrak{s} = \frac{(2\alpha + 2\ell)! \ell! \alpha!}{(\alpha + \ell)! (2\ell)! (2\alpha)!} \cdot \frac{\mathfrak{b}!}{(\mathfrak{b} + \lfloor \vartheta \ell \rfloor)!}.$$

Using Stirling’s formula

$$\ln(n!) = \left(n + \frac{1}{2} \right) \ln n - n + \frac{\ln 2\pi}{2} + r_n,$$

where

$$0 < r_n < \frac{1}{12n},$$

we obtain

$$\begin{aligned} \ln \mathfrak{s} &> (\alpha + \ell) \ln(\alpha + \ell) - \ell \ln \ell - \alpha \ln \alpha + \vartheta \ell - 2 \\ &\quad + \left(\mathfrak{b} + \frac{1}{2} \right) \ln \mathfrak{b} - \left(\mathfrak{b} + \vartheta \ell + \frac{1}{2} \right) \ln(\mathfrak{b} + \vartheta \ell). \end{aligned} \tag{A.9}$$

Let

$$\tilde{\alpha} = \frac{\alpha}{\ell}, \quad \tilde{\mathfrak{b}} = \frac{\mathfrak{b}}{\ell}.$$

Note that

$$\tilde{\alpha}, \tilde{\mathfrak{b}} > 1, \tag{A.10}$$

in view of (A.4) and (A.5). With this notation, we have

$$\begin{aligned} \ln \mathfrak{s} &> \ell \ln \tilde{\alpha} + (\tilde{\alpha} + 1) \ell \ln \left(1 + \frac{1}{\tilde{\alpha}} \right) - \vartheta \ell \ln \ell + \vartheta \ell - 2 \\ &\quad - \vartheta \ell \ln \tilde{\mathfrak{b}} - \left((\tilde{\mathfrak{b}} + \vartheta) \ell + \frac{1}{2} \right) \ln \left(1 + \frac{\vartheta}{\tilde{\mathfrak{b}}} \right) \\ &> \ell \ln \tilde{\alpha} - \vartheta \ell \ln \ell - 2 - \vartheta \ell \ln \tilde{\mathfrak{b}} - \left(\vartheta \ell + \frac{1}{2} \right) \ln \left(1 + \frac{\vartheta}{\tilde{\mathfrak{b}}} \right) \end{aligned}$$

$$> (1 - \vartheta\delta)\mathfrak{k} \ln \tilde{\mathfrak{a}} - \vartheta\mathfrak{k} \ln c - \vartheta\delta\mathfrak{k} \ln \mathfrak{k} - 2 - \left(\vartheta\mathfrak{k} + \frac{1}{2}\right) \ln(1 + \vartheta), \tag{A.11}$$

where the first inequality is a restatement of (A.9), the second follows from

$$0 < \ln(1 + \xi) < \xi \quad (\xi > 0),$$

and the last one follows from (A.2) and (A.10).

Concerning the prime-counting function $\pi(\mathfrak{n})$, it is known that [37, (3.6)]

$$\pi(\mathfrak{n}) < 1.25506 \frac{\mathfrak{n}}{\ln \mathfrak{n}} \quad (\mathfrak{n} > 1).$$

By this, (A.8), and (A.10), we have

$$\begin{aligned} \ln \mathfrak{s} &\leq (\pi(2\mathfrak{k} + 1) - 1) \left(\ln \tilde{\mathfrak{a}} + \ln 2\mathfrak{k} \left(1 + \frac{1}{\tilde{\mathfrak{a}}} \right) \right) \\ &< \left(1.25506 \frac{2\mathfrak{k} + 1}{\ln(2\mathfrak{k} + 1)} - 1 \right) (\ln \tilde{\mathfrak{a}} + \ln 4\mathfrak{k}). \end{aligned} \tag{A.12}$$

Combining (A.11) and (A.12), it follows that

$$\begin{aligned} &\left((1 - \vartheta\delta)\mathfrak{k} - 1.25506 \frac{2\mathfrak{k} + 1}{\ln(2\mathfrak{k} + 1)} + 1 \right) \ln \tilde{\mathfrak{a}} \\ &< \vartheta\mathfrak{k} \ln c + \vartheta\delta\mathfrak{k} \ln \mathfrak{k} + 2 + \left(\vartheta\mathfrak{k} + \frac{1}{2}\right) \ln(1 + \vartheta) \\ &\quad + \left(1.25506 \frac{2\mathfrak{k} + 1}{\ln(2\mathfrak{k} + 1)} - 1 \right) \ln 4\mathfrak{k}. \end{aligned} \tag{A.13}$$

If we set

$$\mathfrak{k}_0 = \mathfrak{k}_0(\vartheta, \delta) = \frac{1}{2} \left(\exp\left(\frac{2.51012}{1 - \vartheta\delta}\right) - 1 \right) > 0$$

for example, then we have

$$(1 - \vartheta\delta)\mathfrak{k} - 1.25506 \frac{2\mathfrak{k} + 1}{\ln(2\mathfrak{k} + 1)} + 1 \geq \frac{1 + \vartheta\delta}{2} > 0 \quad (\mathfrak{k} \geq \mathfrak{k}_0).$$

Hence, whenever $\mathfrak{k} \geq \mathfrak{k}_0$, it follows from (A.13) that $\ln \mathfrak{a} = \ln \tilde{\mathfrak{a}} + \ln \mathfrak{k}$ is bounded in terms of ϑ, δ, c , and \mathfrak{k} , from which and (A.2) it follows that there are only finitely many choices for the pairs $(\mathfrak{a}, \mathfrak{b})$.

This completes the proof of Proposition 7.2.