



Revision of *Xanthopsis bodracus* Makarenko, 1956 (Crustacea: Decapoda) from the Palaeogene of Ukraine

Revizija *Xanthopsis bodracus* Makarenko, 1956 (Crustacea: Decapoda) iz paleogena Ukrajine

Vitaly DERNOV

Institute of Geological Sciences, National Academy of Sciences of Ukraine, 55 b, Oles Honchar Str., Kyiv, 01601, Ukraine; National Museum of Natural History, National Academy of Sciences of Ukraine, 15, Bohdan Khmelnytskyi Str., Kyiv, 01054, Ukraine, e-mail: vitalydernov@gmail.com

Prejeto / Received 23. 1. 2025; Sprejeto / Accepted 4. 4. 2025; Objavljeno na spletu / Published online 30. 7. 2025

Key words: decapod crustaceans, *Harpactoxanthopsis*, *Xanthopsis*, Lutetian, taxonomy
Ključne besede: deseteronožni raki, *Harpactoxanthopsis*, *Xanthopsis*, lutecij, taksonomija

Abstract

The present article constitutes a revision of the decapod crustacean species *Xanthopsis bodracus* Makarenko, 1956 from the early Lutetian (Eocene, Palaeogene) of the Crimean Peninsula, southern Ukraine. The distinctive morphological characteristics of *Xanthopsis bodracus* enable its classification within the genus *Harpactoxanthopsis* Via Boada, 1959, as the type series of *Xanthopsis bodracus* precisely aligns with the diagnostic criteria of this genus. Of the five species in the genus *Harpactoxanthopsis*, *Xanthopsis bodracus* is morphologically closest to *Harpactoxanthopsis quadrilobatus* (Desmarest, 1822). However, there are currently insufficient grounds to confidently synonymise *Xanthopsis bodracus* with *Harpactoxanthopsis quadrilobatus*, as the poorly preserved type material, of which only a part has been preserved, does not allow this. In view of this, *Xanthopsis bodracus* has been revised as *Harpactoxanthopsis* sp. The Palaeogene decapod assemblages of Crimea consist of five genera including *Coeloma* A. Milne-Edwards, 1865, *Protocallianassa* Beurlen, 1930, *Arcticocarcinus* Schweitzer et al., 2016, *Xanthopsis* M'Coy, 1849, and *Harpactoxanthopsis* Via Boada, 1959.

Izvilleček

Pričujoči članek predstavlja revizijo vrste deseteronožcev *Xanthopsis bodracus* Makarenko, 1956 iz zgodnjega lutecija (eocen, paleogen) Krimskega polotoka v južni Ukrajini. Izrazite morfološke značilnosti vrste *Xanthopsis bodracus* omogočajo njeno uvrstitev v rod *Harpactoxanthopsis* Via Boada, 1959, saj se tipska serija vrste *Xanthopsis bodracus* natančno ujema z diagnostičnimi znaki tega rodu. Od petih vrst v rodu *Harpactoxanthopsis* je *Xanthopsis bodracus* morfološko najbližje *Harpactoxanthopsis quadrilobatus* (Desmarest, 1822). Vendar trenutno ni dovolj razlogov za zanesljivo sinonimizacijo *Xanthopsis bodracus* s *Harpactoxanthopsis quadrilobatus*, saj slabo ohranjen tipski material, od katerega se je ohranil le del, tega ne omogoča. Glede na to je bil *Xanthopsis bodracus* revidiran kot *Harpactoxanthopsis* sp. Paleogensko dekapodno združbo Krima sestavlja pet rodov. Med njimi so *Coeloma* A. Milne-Edwards, 1865, *Protocallianassa* Beurlen, 1930, *Arcticocarcinus* Schweitzer et al., 2016, *Xanthopsis* M'Coy, 1849, in *Harpactoxanthopsis* Via Boada, 1959.

Introduction

In Ukraine, Palaeogene decapod crustaceans (order Decapoda Latreille, 1802) occur in the Kanivian (Ypresian) and Kyivian (late Lutetian–Bartonian) regional stages of the Dnipro River basin (Radkevich, 1900; Chernyshev, 1949), the Kyivian regional stage in the Donets Basin (Likharev, 1917; Chernyshev, 1949), the Danian, Thanetian, Ypresian, Lutetian, and Rupelian of the Crimean

Peninsula (Makarenko, 1956; Birshtein, 1960; Levitsky, 1974; Korobkov, 1975; Ilyin, 2005; Dervnov & Udovychenko, 2023), the Oligocene of the Ukrainian Carpathians (Gorbach, 1956; Hyžný et al., 2022), and the Nikopol Basin (Griaznov, 1956; Selin, 1964). However, recently this group has not attracted much attention of Ukrainian palaeontologists, despite the fact that its study is of great palaeogeographic importance.

The most diverse assemblages of Ukrainian Palaeogene decapods were recorded from Crimea and consists of at least five genera, *Coeloma* A. Milne-Edwards, 1865, *Protocallianassa* Beurlen, 1930, *Arcticocarcinus* Schweitzer et al., 2016, *Xanthopsis* M'Coy, 1849, and *Harpactoxanthopsis* Via Boada, 1959 (Levitsky, 1974; Ilyin, 2005; Dervov & Udovychenko, 2023).

In 1956, Ukrainian palaeontologist Dr. Dmytro Yelyseyovych Makarenko (1925–2008) described a new crab species, *Xanthopsis bodracus*, from the “crab horizon”, a local marker stratigraphic level containing decapod remains, which occurs in the early Lutetian limestone succession (Vassilenko, 1952; Muratov & Nemkov, 1960; Zernetsky, 1962; Ilyin, 2005; Zernetsky et al., 2015). Unfortunately, Makarenko (1956) did not define the holotype of this species and the place of storage of the examined material. This work is devoted to the revision of the surviving part of the type material

of *Xanthopsis bodracus* Makarenko, 1956, which was found by the present author in the collections of late Dr. Makarenko.

Material and methods

The study is based on a small collection (NMNH-G 8607) of moderately preserved remains of *Xanthopsis bodracus* Makarenko, 1956, consisting of six carapace fragments and their inner moulds preserved in limestone. Of the four specimens of *Xanthopsis bodracus* carapaces figured by Makarenko (1956: figs 1–4; see also Fig. 1), only one specimen (NMNH-G 8607/11) was found in the collection NMNH-G 8607. The carapace mould figured by Makarenko (1956: Fig. 1), the carapace mould with a right claw (Fig. 2 in Makarenko, 1956), and an isolated claw (Fig. 3 in Makarenko, 1956) are absent (probably lost). The images provided by Makarenko (1956) are the only feature that allows us to identify the type series of




Fig. 1. Specimens of the type series of *Xanthopsis bodracus* Makarenko, 1956, as figured in the protologue (Makarenko, 1956: Figs. 1–4). A – “The mould of the carapace of an adult male (dorsal view)” (Makarenko, 1956: Fig. 1); hereinafter translated from Ukrainian by the author. B – “The mould of the carapace with the right claw and abdomen pressed to the ventral surface; ventral view” (Makarenko, 1956: Fig. 2). C – “The cephalothorax mould, which well showing the ornamentation; female individual in dorsal view” (Makarenko, 1956: Fig. 4). D – “The damaged right claw of a large crab specimen; ventral view” (Makarenko, 1956: Fig. 3). Remark: Makarenko (1956) did not indicate the specimen numbers and did not provide data that would allow to determine their size.




Fig. 2. Geographical location (A, B) and stratigraphic position (C) of the type locality of *Xanthopsis bodracus* Makarenko, 1956 (marked with a purple asterisk). Lithological column of the Skelyaste section (Fig. 2C) modified after Zernetsky (1962: p. 196).

Xanthopsis bodracus, since the total amount of the studied material and a holotype are not specified in the protologue.

The material examined was collected by the Dr. Makarenko from the lower Lutetian (Eocene) limestones exposed as a steep cliff on the right bank of the Bodrak River near the village of Skelyaste, Autonomous Republic of Crimea, Ukraine (Fig. 2) during an excursion of a meeting on the Palaeogene stratigraphy of the southern part of the former USSR in 1955.

In 1956, these decapod remains, represented by at least seven specimens, were described as a new species *Xanthopsis bodracus* Makarenko, 1956. The type material of *Xanthopsis bodracus* was considered lost until it was found in November 2023 in the Makarenko's collection stored in the Department of Stratigraphy and Palaeontology of Cenozoic Sediments, Institute of Geological Sciences, National Academy of Sciences of Ukraine, Kyiv. The research was also greatly aided by Dr. Makarenko's notes taken during the 1955 excursion. Now, the studied collection (NMNH-G

8607) is stored in the Department of Geology, National Museum of Natural History, National Academy of Sciences of Ukraine, Kyiv.

The collection NMNH-G 8607 consists of 15 specimens (NMNH-G 8607/01 to NMNH-G 8607/15). The fossil-bearing rock is a white limestone with small shell debris. In addition to decapod remains, the collection NMNH-G 8607 contains tests of unidentified nummulitids, steinkerns of the bivalves *Thracia* sp. and *Chlamys* sp., burrows *?Palaeophycus* isp., and a fragment of the steinkern of the nautiloid *Aturia* sp. bearing a poorly preserved external mould of a serpulid tube and burrows.

The specimen NMNH-G 8607/07 bears the trace fossils assigned to the ichnogenus *Arachnosstega* Bertling, 1992, preserved on the surface of the crab carapace inner mould (see Fig. 4A, C). This ichnogenus is usually interpreted as a domichnia or feeding structure in a consolidated soft- to firmground substrate, produced by detritus- or deposit-feeding polychaetes (Bertling, 1992; Fatka et al., 2011; Zatoń, 2020; Dernov, 2023).

This paper uses the decapod taxonomy proposed by Schweitzer et al. (2010) and morphological terminology and methods of describing fossil decapods summarized by Schweitzer et al. (2024).

Geological setting

The Skelyaste section is represented by a sequence of predominantly white and grey nummulitic non-laminated limestones of the Bakhchisarai and Simferopol formations ascribed to the Ypresian and lower Lutetian (Zernetsky et al., 2015) (see Fig. 2C). Based on foraminifer studies, the Ypresian/Lutetian boundary in this section is located at the base of Unit 7 (personal communication of Dr. Tamara S. Ryabokon, Institute of Geological Sciences of the NAS of Ukraine, February 2024).

Numerous foraminifers, such as *Nummulites distans* Deshayes, 1838, *N. atacicus* Leymerie, 1846, *Discocyclus archiaci* (Schlumberger, 1903), *D. sella* (d'Archiac, 1850), *Assilina spira* (Roissy, 1805), *A. exponens* (Sowerby, 1840), *Operculina gigantea* Mayer, 1876, as well as bivalves *Pycnodonta rarilamella* (Deshayes, 1864), *Spondylus* cf. *tenuispina* Sandberger, 1863, *Chlamys* cf. *opia* Vassilenko, 1952, *Cardium* cf. *gigas* Defrance, 1817, *Chama calcarata* Lamarck, 1806, gastropods *Strombus* sp., *Cerithium* sp., *Rostellaria ampla* Rutot, 1876, and echinoids *Conoclypeus conoides* Agassiz, 1839 occur in the Ypresian part of the limestone succession (Units 1 to 6 – see Fig. 2C) (Zernetsky, 1962; Vyalov, 1975).

The foraminifers *Nummulites polygyratus* Deshayes, 1838, *N. distans* Deshayes, 1838, *Assilina exponens* (Sowerby in Sykes, 1840), *A. irregularis* Carter, 1853, bivalves *Spondylus* cf. *eichwaldi* Fuchs, 1870, *Pseudamussium solea* (Deshayes, 1830), and echinoids *Conoclypeus conoides* Agassiz, 1839 were found in Unit 7, i.e. in the Lutetian part of the limestone succession (Zernetsky, 1962; Vyalov, 1975).

According to Makarenko (1956), the specimens of *Xanthopsis bodracus* was found in a limestone interlayer, about 0.4-m-thick, which lies in the lower part of Unit 7 (Fig. 2C). Numerous foraminifers, shells of bivalves, and moulds of the gastropods, up to 0.45 m long, as well as trace fossils co-occurring with crabs (Makarenko, 1956).

The lower part of the Lutetian in some sections of the Crimea contains the so-called “crab horizon”, which is a local correlation marker (Vassilenko, 1952; Muratov & Nemkov, 1960). In addition to *Xanthopsis bodracus*, *Harpactoxanthopsis lutugini* (Likharev, 1917) and *H. cf. lutugini* (Likharev, 1917) also occur in the “crab horizon”

(Ilyin, 2005). It should be noted, however, that in the Ypresian–Lutetian interval of Crimea, crabs are known both in the highest part of the Ypresian and in the lowest part of the Lutetian (Vassilenko, 1952; Zernetsky, 1962; Vyalov, 1975), but the specimens described by Makarenko (1956: p. 74) come from the basal part of the Lutetian.

The Ypresian-Lutetian nummulitic limestone sequence in the Crimea probably accumulated in the distal part of the inner ramp above the base of wind waves at fast sedimentation rates and high hydrodynamics (Lygina et al., 2010; Zernetsky et al., 2015). Similar environmental conditions have been recorded for the genus *Harpactoxanthopsis* Via Boada, 1959, to which *Xanthopsis bodracus* belongs (see the section “Systematics”), in the Eocene of Croatia and Slovenia (Mikuz, 2002; Gašparič et al., 2015; Križnar & Gašparič, 2019).

Systematics

Makarenko (1956) did not provide a formal diagnosis of *Xanthopsis bodracus*, but briefly described the morphology of the type specimens in detail. Makarenko (1956: p. 75–76; translated from Ukrainian by the author) notes, that “The posterior part of the carapace resembles a trap-ezoid, the lateral sides of which converge at an angle of 80–83° outside the small base. The posterolateral margins are almost straight, rounded. The abdomen is small, flattened, consists of four segments, bent under the cephalothorax and is located in the longitudinal depression of the ventral surface. The dorsal side of the finger is wedge-shaped. The cutting edge of the pollex bears small teeth. The surface of the carpus is rounded and smooth below; above, on the keel projection, there are 5–6 small pointed tubercles. Other four pairs of pereopods are not preserved. The surface of the carapace is evenly convex and no separate regions are distinguished. The carapace is evenly covered with dense, small, round pits. The same pits are present on the claws. Symmetrically arranged diamond-shaped depressions with branches extending posteriorly are present in the most convex part of the carapace. An adult specimen is 41 mm long, 50 mm wide, and has a maximum height of 18 mm”.

Only the material figured by Makarenko (1956: Figs. 1–4) allows us to get an idea of the morphology of *Xanthopsis bodracus*, since the unfigured specimens from the collection NMNH-G 8607 (Fig. 4), although referred to *Xanthopsis bodracus* here, are not part of the type series.

Specimen NMNH-G 8607/11 (Fig. 3A–C) is a moderately-preserved carapace inner mould in




Fig. 3. Specimens of *Xanthopsis bodracus* Makarenko, 1956 of the type series. A–C – specimen NMNH-G 8607/11 (A – posterior view, B – lateral view, C – dorsal view). D – specimen NMNH-G 8607/12 (counterpart of NMNH-G 8607/11) in dorsal view. Scale bars = 5 mm.




Fig. 4. Decapod specimens from the collection NMNH-G 8607. A, C – carapace fragment of *Xanthopsis bodracus* bearing trace fossils *Arachnostega* (specimen NMNH-G 8607/07: A – general view, C – enlarged part of the carapace with *Arachnostega* burrows). B – carapace fragment of *Xanthopsis bodracus* in dorsal view (specimen NMNH-G 8607/08). D – carapace fragment of *Xanthopsis bodracus* in dorsal view (specimen NMNH-G 8607/09). E – carapace fragment of *Xanthopsis bodracus* in ventral view (specimen NMNH-G 8607/10). Scale bars = 5 mm

the dorsal aspect. Specimen NMNH-G 8607/12 is a counterpart impression of the dorsal surface of NMNH-G 8607/11 with a preserved cuticle. The carapace is rounded-hexagonal almost ovate in outline (Fig. 3C), narrowing considerably posteriorly with poorly defined regions, strongly convex longitudinally (Fig. 3B) and moderately convex transversely (Fig. 3A), 34 mm wide and 27 mm long, and almost as wide as long (length/width = 0.79). The greatest width of the carapace is approximately in the middle of its length. The front is slightly convex, almost straight, approximately 12 mm wide (about 33 % of the total carapace width); fronto-orbital width is 23 mm. Almost circular orbits, 4 mm wide and 2.5 mm deep, are poorly preserved (see Figs. 3B, C). The anterolateral margins are not preserved. The posterolateral margins are very slightly convex, almost straight; the posterior margin is narrow, straight, 10 mm wide. The base of a low spine is preserved on the right posterolateral margin in the specimen NMNH-G 8607/11.

On the dorsal surface of the carapace mould, there are weak branchiocardial shallow grooves, which limit the slightly convex cardiac region in the transverse aspect. The ventral surface and pereopods in NMNH-G 8607/11 and NMNH-G 8607/12 are not preserved.

The dorsal surface of the carapace NMNH-G 8607/11 is poorly preserved; only in front, near the frontal margin, small pits occur. Numerous rounded pits are present on the carapace counterpart with a partially preserved cuticle (NMNH-G 8607/12).

Discussion and concluding remarks

The morphological description of *Xanthopsis bodracus* given above does not contradict its description presented by Makarenko (1956). However, he noted some morphological details that cannot be observed on the specimen NMNH-G 8607/11, although their reliability is not in doubt, as they are confirmed by the images (Makarenko,

1956: figs. 1–4). For example, Makarenko (1956: p. 75) reports that the carapace front is divided by four spines, of which the two outer ones form the inner orbital spines; three spines are present on each of the anterolateral margins (see Fig. 1).

The characteristic morphological features of *Xanthopsis bodracus* allow to place it to the genus *Harpactoxanthopsis* Via Boada, 1959 (type species: *Cancer quadrilobata* Desmarest, 1822). The material described above, as well as the images of the type series of *Xanthopsis bodracus* given by Makarenko (1956), fully correspond to the diagnosis of this genus given by Schweitzer (2003: p. 1119), namely: “Carapace about 80 percent as long as wide, ovate; regions poorly defined; branchiocardiac groove usually well-defined along lateral margins of urogastric region. Front with four blunt spines including inner orbital spine; anterolateral margin convex, with five spines excluding outer orbital spine”.

According to Schweitzer et al. (2010), the genus *Harpactoxanthopsis* includes five species, *H. bittneri* (Lörenthey, 1898), *H. lutugini* (Likharev, 1917), *H. quadrilobatus* (Desmarest, 1822), *H. souverbei* (A. Milne-Edwards, 1862), and *H. villaltae* Via Boada, 1959. Of all these species, *H. lutugini* and *H. quadrilobatus* are the most morphologically similar to *Xanthopsis bodracus*.

No morphological differences were found between the crabs described by Makarenko (1956) and *H. quadrilobatus* described from other localities (e.g., Desmarest, 1822; A. Milne Edwards, 1862; Bittner, 1875; Lörenthey and Beurlen, 1929; Vía, 1969; Beschin et al., 1994; Hyžný, 2014; Gašparič et al., 2015). In Ukraine, *Harpactoxanthopsis quadrilobatus* previously was recorded by Chernyshev (1949) from the Kyiv Formation (late Lutetian–Bartonian) of the vicinity of the city of Kyiv. However, there are currently insufficient grounds to confidently synonymise *Xanthopsis bodracus* with *Harpactoxanthopsis quadrilobatus*, as the poorly preserved type material, of which only a part has been preserved, does not

Table 1. Palaeogene decapods of Crimea (modified after Dervov and Udovychenko, 2023, table 1).

Taxa	Age	References
<i>Coeloma vigil</i> Milne-Edwards, 1865	Rupelian	Ilyin, 2005
<i>Harpactoxanthopsis</i> sp.	Early Lutetian	This study
<i>Harpactoxanthopsis</i> cf. <i>lutugini</i> (Likharev, 1917)	Late Lutetian	Ilyin, 2005
	Ypresian	
<i>Harpactoxanthopsis lutugini</i> (Likharev, 1917)	Late Ypresian or early Lutetian	Levitsky, 1974
<i>Xanthopsis nodosa</i> McCoy, 1849	Early Ypresian	Ilyin, 2005
<i>Protocallianassa</i> sp.	Danian	Levitsky, 1974
<i>Arcticocarcinus</i> cf. <i>insignis</i> (Segeber, 1900)	Early Danian	Dervov & Udovychenko, 2023

allow this. In view of this, *Xanthopsis bodracus* has been revised as *Harpactoxanthopsis* sp. Thus, taking into account the data obtained, the Palaeogene decapod assemblages of Crimea consist of five genera (Table 1). These records often have an unclear stratigraphic position and therefore it is difficult to analyse the stratigraphic distribution of taxa.

Acknowledgements

The author is grateful to Dr. Tamara S. Ryabokon for her advice on the Skelyaste section and assistance with the collection of the late Dr. Dmytro Makarenko. The author is sincerely grateful to anonymous reviewers, whose comments and suggestions improved the quality of the final version of the manuscript. The research was carried out within the framework of the scientific theme “Late Precambrian and Phanerozoic biota of Ukraine: biodiversity, revision of systematic composition and phylogeny of leading groups” (No. 0122U001609).

References

- Agassiz, L. 1839: Description des Echinodermes fossiles de la Suisse. 1. *Spatangoides* et *Clypeasteroides*. Nouveau Mémoire de la Société Helvétique des Sciences naturelles, 3: 1–101.
- d’Archiac, E.J.A.D. 1850: Description des fossiles du groupe nummulitique recueillis par M. S.-P. Pratt et M.J. Delbos aux environs de Bayonne et de Dax. Mémoires de la Société géologique de France, deuxième série, 3: 397–456.
- Bertling, M. 1992: *Arachnostega* n. ichnog. – burrowing traces in internal moulds of boring bivalves (late Jurassic, Northern Germany). Paläontologische Zeitschrift, 66: 177–185. <https://doi.org/10.1007/BF02989487>
- Beschin, C., Busulini, A., de Angeli, A. & Tessier, G. 1994: I Crostacei Eocenici della cava “Boschetto” di Nogarole Vicentino (Vicenza-Italia settentrionale). Lavori Società Veneziana di Scienze Naturali, 19: 159–215.
- Beurlen, K. 1930: Vergleichende Stammesgeschichte, Grundlagen, Methoden und Probleme unter besonderer Berücksichtigung der höheren Krebsen. Fortschritte der Geologie und Palaeontologie, 8: 317–586.
- Birshtein, J.A. 1960: Order Decapoda. In: Orlov, Yu.A. (ed.): Fundamentals of Palaeontology, 8. Arthropods: trilobites and crustaceans. State Scientific and Technical Publishing House of Literature on Geology and Protection of Subsoil, Moscow: 439–456.
- Bittner, A. 1875: Die Brachyuren des vicentinischen Tertiärgebirges. Denkschriften der Kaiserlichen Akademie der Wissenschaften. Mathematisch-Naturwissenschaftliche Classe, 34: 63–106.
- Carter, H.J. 1853: Descriptions of some of the larger forms of fossilised Foraminifera in Scinde; with observations on their internal structure. Annals and Magazine of Natural History, 11/63: 161–177. <https://doi.org/10.1080/03745485609496239>
- Chernyshev, B.I. 1949: Subclass Malacostraca. Order Decapoda. In: Ryabinin, A.N. & Korobkov, I.A. (eds.): Atlas of the reference forms of the fossil fauna of the USSR. Vol. 12. Palaeogene. State Publishing House of Geological Literature, Moscow: 289–294.
- Defrance, J.L.M., 1817: Cardites (foss.). In: Cuvier, F. (ed.): Dictionnaire des Sciences Naturelles. Vol. 7. Levrault, Strasbourg & Le Normant, Paris: 87–91.
- Derenyuk, N.Ye. 1984: Geological Map of the Crimean Mountains. Scale 1:200 000. Krymgeologiya, Sevastopol.
- Dernov, V.S. 2023: *Ichtyocoprus affectatus* Makarenko, 1993 is a bivalve boring rock cast, not a coprolite. Collection of scientific works of the Institute of Geological Sciences of the NAS of Ukraine, 16/1: 70–80. <https://doi.org/10.30836/igs.2522-9753./12023.295187>
- Dernov, V. & Udovychenko, M. 2023: The first record of the genus *Arcticocarcinus* Schweitzer et al., 2016 (Decapoda, Brachyura) in the Danian of Crimea (Ukraine) and its palaeobiogeographical significance. Bulletin of Taras Shevchenko National University of Kyiv. Geology, 101/2: 5–14. <https://doi.org/10.17721/1728-2713.101.01>
- Deshayes, G.P. 1824–1837: Description des coquilles fossiles des environs de Paris. Published by the author, Paris: 393 p. <https://doi.org/10.5962/bhl.title.52303>
- Deshayes, G.P. 1838: Description des coquilles fossiles recueillies en Crimée par M. de Verneuil, et observations générales à leur sujet. Mémoires de la Société géologique de France. Série 1, 3: 37–69.
- Deshayes, G.P. 1864–1865: Description des animaux sans vertèbres découverts dans le bassin de Paris pour servir de supplément à la Description des coquilles fossiles des environs de Paris comprenant une revue générale de toutes les espèces actuellement connues. Tome troisième. Mollusques céphalés, deuxième partie. Mollusques céphalopodes. Baillière, Paris: 667 p.

- Desmarest, A.G. 1822: Les crustacés proprement dits. In: Brongniart, A. & Desmarest, A.G. (eds.): Histoire naturelle des crustacés fossiles, sous les rapports zoologiques et géologiques. F.-G. Levrault, Paris: 67–142.
- Fatka, O., Mikuláš, R., Szabad, M., Micka, V. & Valent, M. 2011: *Arachnostega* Bertling, 1992 in the Drumian (Cambrian) sediments of the Teplá-Barrandian region (Czech Republic). *Acta Geologica Polonica*, 61/4: 367–381.
- Fuchs, T. 1870: III. Beiträge zur Kenntniss fossiler Binnenfaunen. III. Die Fauna der Congerien-schichten von Radmanest im Banate. *Jahrbuch der Kaiserlich Königlichen Geologischen Reichsanstalt*, 20: 343–364.
- Gašparič, R., Fraaije, R.H.B., Van Bakel, B.W.M., Jagt, J.W.M. & Skupien, P. 2015: Mesozoic–Cenozoic crustaceans preserved within echinoid tests and bivalve shells. *Bulletin of Geosciences*, 90/3: 601–611. <https://doi.org/10.3140/bull.geosci.1551>
- Gorbach, L.P. 1956: On the crabs from the Menilite shales of the Eastern Carpathians. *Transactions of the Lviv Geological Society*, 2/3: 30–42.
- Griaznov, V.I. 1956: On the confinement of finds of the crabs *Coeloma vigil* M.-Edw. to Oligocene manganese-ore facies. *Reports of the Academy of Sciences of the USSR*, 106: 717–719.
- Hyžný, M. 2014: *Harpactoxanthopsis quadrilobatus* (Desmarest, 1822) from the Eocene of Slovakia and Italy: the phenomenon of inverted images of fossil heterochelous crabs. *Bulletin of the Mizunami Fossil Museum*, 40: 23–27. <https://doi.org/10.5474/geologija.2002.008>
- Hyžný, M., Kovalchuk, O., Świdnicka, E., Barkaszi, Z., Berezovsky, A., Dumitriu, S., Grădianu, I. & Stefaniak, K. 2022: Revisiting brachyuran crabs (Malacostraca: Decapoda) from Oligocene and Miocene fish beds of Europe. *Geologica Carpathica*, 73/6: 579–597. <https://doi.org/10.31577/geolcarp.73.6.3>
- Ilyin, I.V. 2005: Cretaceous and Palaeogene decapod crustaceans (Crustaceomorpha, Decapoda) of the western part of Northern Eurasia. Moscow University Press, Moscow: 295 p.
- Korobkov, I.A. 1975: Decapod crustaceans. In: Grossheim, V.A. & Korobkov, I.A. (eds.): Stratigraphy of the USSR. Palaeogene System. Nedra, Moscow: 433–435.
- Križnar, M. & Gašparič, R. 2019: Eocene of Slovenian Istria and Lutetian “Transitional beds with Crabs”. Book of abstracts, 7th Symposium on Mesozoic and Cenozoic decapod crustaceans (Ljubljana, 17–21 June 2019). *Tiskarna Oman, Kranj*: 116–123.
- Lamarck, J.P. 1806: Memoire sur les fossiles des environs de Paris. *Annales du Museum d’Histoire naturelle de Paris*, 8: 156–166.
- Latreille, P.A. 1802: Histoire Naturelle, Générale et Particulière des Crustacés et des Insectes. Ouvrage Faisant suite à l’Histoire Naturelle Générale et Particulière, Composée par LeClerc de Buffon, et Rédigée par C.S. Sonnini, Membre de Plusieurs Sociétés Savantes. Vol. 3. Dufart, Paris: 468 p. <https://doi.org/10.5962/bhl.title.60678>
- Levitsky, E.S. 1974: Fossil decapod crustaceans of the city of Bakhchisarai area. *Bulletin of the Moscow Society of Naturalists. Geology*, 49: 101–119.
- Leymerie, A. 1846: Mémoire sur le terrain à *Nummulites* (épicrotécé) des Corbières et de la Montagne Noire. *Mémoires de la Société Géologique de France, serie 2*, 1: 337–372.
- Likharev, B.K. 1917: Crab remains from the lower Tertiary of the Pridonetskii Krai. *Yearbook of the Russian Palaeontological Society*, 1: 13–24.
- Lőrenthey, E.: 1898. Beiträge zur Decapodenfauna des ungarischen Tertiärs. *Természetráji Füzetek*, 21: 1–133. <https://doi.org/10.5962/bhl.title.10253>
- Lőrenthey, E. & Beurlen, K. 1929: Die fossilen der Länder der Ungarischen Krone. *Geologica Hungarica, Series Palaeontologica*, 3: 1–420.
- Lygina, Ye.A., Kopaevich, L.F., Nikishin, A.M., Shalimov, I.V. & Yakovishina, Ye.V. 2010: Lower and middle Eocene deposits of the Crimean Peninsula: facies features and depositional environments. *Bulletin of the Moscow University. Series 4. Geology*, 6: 11–22.
- Makarenko, D.Ye. 1956: Crab remains from the Palaeogene deposits of the Crimea. *Geological Journal (Ukraine)*, 16: 74–76.
- Mayer, C. 1876: Description de Coquilles fossiles des terrains tertiaires supérieurs (suite). *Journal de Conchyliologie*, 24: 168–180.
- M’Coy, F. 1849: On the classification of some British fossil Crustacea, with notices of new forms in the university collection at Cambridge. *Annals and Magazine of Natural History*, 4/23: 161–179, 330–335, 392–414. <https://doi.org/10.1080/03745486009494843>
- Mikuž, V. 2002: Nova najdba rakovice *Harpactoxanthopsis quadrilobata* (Desmarest) v eocenskem flišu pri Gračišču blizu Pazina v Istri (Hrvaška). *Geologija*, 45/1: 97–102. <https://doi.org/10.5474/geologija.2002.008>
- Milne-Edwards, A. 1865: Description de quelques Crustacés Nouveaux appartenant à la tribu des Maiens. *Annales De La Société Ento-*

- mologique De France, 4: 133–147. <https://doi.org/10.5962/bhl.title.10083>
- Muratov, M.V. & Nemkov, G.I. 1960: Palaeogene deposits of the Bakhchisarai area and their significance for Palaeogene stratigraphy of the southern part of the USSR. In: Yanshin, A.L., Vyalov, O.S., Dolgoplov, N.N. & Menner, V.V. (eds.): Palaeogene deposits of the southern part of the European part of the USSR. Publishing of the Academy of Sciences of the USSR, Moscow: 15–23.
- Quayle, W.J. & Collins, J.S.H. 1981: New Eocene crabs from the Hampshire Basin. *Palaeontology*, 24: 733–758.
- Radkevich, G.O. 1900: On the lower Tertiary deposits in the vicinity of Kaniv. *Transactions of the Kyiv Society of Naturalists*, 16: 1–45.
- de Roissy, F. 1805: Histoire naturelle générale et particulière des Mollusques; livre faisant suite aux œuvres de Buffon, et partie du cours complet d'Histoire naturelle rédigé par C. S. Sonnini. Tome 5. Dufart, Paris: 590 p.
- Rutot, A. 1876: Note sur quelques fossiles recueillis dans le diluvium des environs de Tongres. *Annales de la Société Malacologique de Belgique*, 10: 7–20.
- Sandberger, C.L.F. 1858–1863: Die Conchylien des Mainzer Tertiärbeckens. Kriedel und Niedner, Wiesbaden: 458 p. <https://doi.org/10.5962/bhl.title.13953>
- Schlumberger, C. 1903: Troisième note sur les Orbitoides. *Bulletin de la Société Géologique de France. Série 4*, 3: 273–289.
- Schweitzer, C.E. 2003: Utility of proxy characters for classification of fossils: an example from the fossil Xanthoidea (Crustacea: Decapoda: Brachyura). *Journal of Paleontology*, 77/6: 1107–1128. [https://doi.org/10.1666/0022-3360\(2003\)077<1107:uopfcf>2.0.co;2](https://doi.org/10.1666/0022-3360(2003)077<1107:uopfcf>2.0.co;2)
- Schweitzer, C.E., Feldmann, R.M., Audo, D., Charbonnier, S., Fraaije, R.H.B., Frantescu, O.D., Hyžný, M., Karasawa, H., Klompmaker, A.A., Luque, J., Robins, C.M., Schram, F.R., Schweigert, G. & Tshudy, D. 2024: Treatise online. Part R, Revised, Volume 1: Generalized external adult Decapoda morphology. The University of Kansas Press, Lawrence: 125 p.
- Schweitzer, C.E., Feldmann, R.M., Garassino, A., Karasawa, H. & Schweigert, G. 2010: Systematic list of fossil decapod crustacean species. *Crustaceana monographs*, 10. Brill, Leiden and Boston: 222 p. <https://doi.org/10.1163/193724012x626575>
- Schweitzer, C.E., Karasawa, H., Luque, J. & Feldmann, R.M. 2016: Phylogeny and classification of Necrocarinoidea Förster, 1968 (Brachyura: Raninoidea) with description of two new genera. *Journal of Crustacean Biology*, 36/3: 338–372. <https://doi.org/10.1163/1937240X-00002432>
- Selin, Y.I. 1964: Oligocene stratigraphy and molluscs of the Bolshe-Tokmaysky manganese-ore district. Nedra, Moscow: 240 p.
- Sowerby, J. de C. 1834–1843: The mineral conchology of Great Britain; or Coloured figures and descriptions of those remains of testaceous animals or shells, which have been preserved at various times and depths in the earth. Vol. 7. Privately published by author, London: 80 p.
- Sykes, W.H. 1840: A notice respecting some fossils collected in Cutch, by Capt. Walter Smee, of the Bombay Army. *Transactions of the Geological Society of London*, 2/5: 715–719. <https://doi.org/10.1144/transgslb.5.3.715>
- Vassilenko, V.K. 1952: Eocene stratigraphy and mollusk fauna of Crimea. *Proceedings of the All-Union Research Geological Exploration Institute*, 59: 1–128.
- Vía, L. 1969: Crustáceos Decápodos del Eoceno español. *Pirineos*, 91–94: 1–479.
- Via Boada, L. 1959: Decápodos fósiles del Eoceno español. *Boletín del Instituto Geológico y Minero de España*, 70: 331–402.
- Vyalov, O.S. 1975: The Bakhchisarai Palaeogene section. 2. Main outcrops of Eocene and Oligocene deposits. *Geology and geochemistry of fossil fuels*, 47: 80–88.
- Zatoń, M. 2020: No evidence for fungal infection of Upper Jurassic horseshoe crabs: A comment on Błażejowski et al. (2019). *Palaeogeography, Palaeoclimatology, Palaeoecology*, 554: article 109142. <https://doi.org/10.1016/j.palaeo.2019.03.043>
- Zernetsky, B.F. 1962: Crimea. Eocene. In: Bondarchuk, V.G. (ed.): *Stratigraphy of the Ukrainian SSR. Palaeogene*. Publishing House of the Academy of Sciences of the Ukrainian SSR, Kyiv: 183–202.
- Zernetsky, B.F., Ryabokon, T.S. & Lulyeva, S.A. 2015: Issue of studying of the Eocene sedimentary succession of the Crimean and Kerch peninsulas. *Collection of scientific works of the IGS of the NAS of Ukraine*, 8: 33–62. <https://doi.org/10.30836/igs.2522-9753.2015.145320>