

# Classification of convex polyhedra by their rotational orbit Euler characteristic

Jurij Kovič \*

*IMFM, Jadranska 19, 1000 Ljubljana, Slovenia;  
UP FAMNIT, Glagoljaška 8, 6000 Koper, Slovenia*

Received 2 February 2015, accepted 10 October 2015, published online 22 August 2016

---

## Abstract

Let  $\mathcal{P}$  be a polyhedron whose boundary consists of flat polygonal faces on some compact surface  $S(\mathcal{P})$  (not necessarily homeomorphic to the sphere  $S^2$ ). Let  $vo_R(\mathcal{P}), eo_R(\mathcal{P}), fo_R(\mathcal{P})$  be the numbers of rotational orbits of vertices, edges and faces, respectively, determined by the group  $G = G_R(\mathcal{P})$  of all the rotations of the Euclidean space  $E^3$  preserving  $\mathcal{P}$ . We define the *rotational orbit Euler characteristic* of  $\mathcal{P}$  as the number  $Eo_R(\mathcal{P}) = vo_R(\mathcal{P}) - eo_R(\mathcal{P}) + fo_R(\mathcal{P})$ .

Using the Burnside lemma we obtain the lower and the upper bound for  $Eo_R(\mathcal{P})$  in terms of the genus of the surface  $S(\mathcal{P})$ . We prove that  $Eo_R \in \{2, 1, 0, -1\}$  for any convex polyhedron  $\mathcal{P}$ . In the non-convex case  $Eo_R$  may be arbitrarily large or small.

*Keywords:* Polyhedron, rotational orbit, Euler characteristic.

*Math. Subj. Class.:* 52B05, 52B10

---

## 1 Introduction

CONTEXT: Euler (1752) discovered the famous relation  $v - e + f = 2$  between the numbers of vertices  $v$ , edges  $e$  and faces  $f$  of any convex polyhedron. This *Euler polyhedron formula* was implicitly stated in the formulas of Descartes (1630)  $p = 2f + 2v - 4$ ,  $p = 2e$ , where  $p$  is the number of "plane angles" – corners of faces determined by pairs of adjacent edges ([5], p.469).

The number  $\chi = v - e + f$  can be defined for any *map* (a graph cellularly embedded into a compact surface  $S$ ) and is called its *Euler characteristic*. It is related to the *genus*  $g$  of the surface  $S$  as follows:  $\chi = 2 - 2g$  ([5], p. 473.) and it may be used for the

---

\*This work is supported in part by the Slovenian Research Agency (research program P1-0294 and research project N1-0032).

*E-mail address:* jurij.kovic@siol.net (Jurij Kovič)

classification of surfaces by two parameters: one is  $\chi$  and the other is orientability (or non-orientability) of the surface. In [4] we introduced the concept of *Euler orbit characteristic*  $Eo = vo - eo + fo$ , where  $vo, eo, fo$  denote the number of orbits of vertices, edges and faces, respectively, determined by the group of all rotations and reflections of the Euclidean space  $E^3$ , preserving the polyhedron  $\mathcal{P}$ , and we used it for the classification of the 92 Johnson solids ([4], p.258). In this paper we introduce a similar concept, called *rotational orbit Euler characteristic*, and we use it for the classification of convex polyhedra.

**Definition 1.1.** The rotational Euler orbit characteristic of the polyhedron  $\mathcal{P}$  is defined as the number  $Eo_R = vo_R - eo_R + fo_R$  where  $vo_R, eo_R, fo_R$  are the numbers of rotational orbits of the vertices, edges and faces, respectively, of  $\mathcal{P}$  (these orbits are determined by the group  $G_R(\mathcal{P})$  of all the rotations of the Euclidean space  $E^3$  preserving  $\mathcal{P}$ ).

**Proposition 1.2.**  $Eo_R = 1$  for all the Platonic solids and all the  $n$ -prisms and  $n$ -antiprisms, while  $1 \leq Eo_R \leq 2$  for all the Archimedean solids.

*Proof.* In the Table 1 the number of rotational orbits of Platonic and Archimedean solids are given. These values can be easily found for each solid directly or deduced from the symmetry-type graphs of Platonic and Archimedean solids [3]. The 5 Platonic solids have just one rotational orbit of vertices, edges and faces. The 13 Archimedean solids have at most two rotational orbits of vertices and at most three rotational orbits of edges and faces. The  $n$ -prisms and the  $n$ -antiprisms have just one rotational orbit of vertices and two rotational orbits of edges and faces. □

class	solid $\mathcal{P}$	vertex pattern	$vo_R$	$eo_R$	$fo_R$	$Eo_R$
I.	tetrahedron	(3.3.3)	1	1	1	1
I.	octahedron	(3.3.3.3)	1	1	1	1
I.	cube	(4.4.4)	1	1	1	1
I.	icosahedron	(3.3.3.3.3)	1	1	1	1
I.	dodecahedron	(5.5.5)	1	1	1	1
II.	truncated tetrahedron	(3.4.3.4)	1	1	2	2
II.	truncated octahedron	(3.5.3.5)	1	1	2	2
III.	truncated cube	(3.6.6)	1	1	2	2
III.	truncated octahedron	(3.8.8)	1	1	2	2
III.	truncated dodecahedron	(4.6.6)	1	1	2	2
III.	truncated icosahedron	(3.10.10)	1	1	2	2
III.	truncated dodecahedron	(5.6.6)	1	1	2	2
IV.	rhombicuboctahedron	(3.4.4.4)	1	2	3	2
IV.	rhombicosidodecahedron	(3.4.5.4)	1	2	3	2
V.	truncated cuboctahedron	(4.6.8)	2	3	3	2
V.	truncated icosidodecahedron	(4.6.10)	2	3	3	2
VI.	snub cube	(3.3.3.3.4)	1	3	3	1
VI.	snub dodecahedron	(3.3.3.3.5)	1	3	3	1
VII.	$n$ -prism	(4.4. $n$ )	1	2	2	1
VIII.	$n$ -antiprism	(3.3. $n$ )	1	2	2	1

Table 1: Values of  $vo_R, eo_R, fo_R$  for Platonic and Archimedean solids and for the infinite families of  $n$ -prisms and  $n$ -antiprisms.

MOTIVATION: Similar bounds on  $Eo_R$  exist for the Johnson solids (i.e. convex polyhedra with regular polygonal faces and at least two orbits of vertices [2]). The direct motivation

for writing this paper came from the empirical observation that the values of  $Eo_R$  for the 92 Johnson solids are in a small range between  $-1$  and  $2$ . This was discovered during the process of constructing a table of 16 parameters of the Johnson solids presented in [4], while the range for  $Eo$  for the same solids turned out to be bigger:  $0 \leq Eo \leq 5$ .

COMPARISON OF  $Eo$  AND  $Eo_R$ : The two characteristics behave very differently on the set of all convex polyhedra: the main result of the paper (Theorem 2.1) states that the relation  $-1 \leq Eo_R \leq 2$  holds for all convex polyhedra, while for  $Eo$  there is no fixed upper bound, we can obtain only the following estimate:  $Eo = vo - eo + fo \leq vo + fo \leq vo_R + fo_R = \leq (vo_R - eo_R + fo_R) + eo_R = Eo_R + eo_R \leq 2 + eo_R$ .

**Definition 1.3.** Let  $G_R(\mathcal{P}) = \{R_1, R_2, \dots, R_{n-1}, R_n = Id\}$  be the group of rotational symmetries of the polyhedron  $\mathcal{P}$ . The poles of the rotation  $R_i$  are the points in which the axis of the rotation  $R_i$  intersects the surface  $S(\mathcal{P})$ . Let  $v_p(R_i)$ ,  $e_p(R_i)$ ,  $f_p(R_i)$  denote the numbers of poles of  $R_i$  in the vertices, edge centers and face centers of  $\mathcal{P}$ , respectively. The number  $E_p(R_i) = v_p(R_i) - e_p(R_i) + f_p(R_i)$  is called the Euler polar characteristics of the rotation  $R_i$ .

**Lemma 1.4.** *Let  $n_i$  denote the number of poles of any non-trivial rotation  $R_i$  of the polyhedral map  $\mathcal{P}$  on the surface  $S$  of genus  $g$ . Then*

$$\begin{aligned} n_i &\leq 2(g + 1), \\ n_i &\in \{0, 2, 4, \dots, 2(g + 1)\}, \\ v_p(R_i) + e_p(R_i) + f_p(R_i) &= n_i, \\ 0 &\leq e_p(R_i) \leq n_i, \\ E_p(R_i) &= n_i - 2e_p(R_i), \\ -2(g + 1) &\leq -n_i \leq E_p(R_i) \leq n_i \leq 2(g + 1). \end{aligned}$$

*If the order of the rotation  $R_i$  is greater than 2 (i.e.  $R_i^n = id$  and  $n > 2$ ), then  $e_p(R_i) = 0$  and  $E_p(R_i) = n_i$ .*

*Proof.* Any line intersecting  $\mathcal{P}$  has at most  $2g$  intersecting points with  $S$ . If  $P$  is a convex polyhedron then any nontrivial rotation  $R_i$  has exactly two poles, hence  $n_i = 2$ . If  $n_i > 2$  then each segment of the rotational axis  $r_i$  not lying in the interior of  $P$  contributes two poles (hence  $n_i$  is an even number!) and at least one new handle. Thus it "increases" the genus of  $S$  for 1 (since it is well known that the genus counts the numbers of "handles" of a surface), therefore it must be  $n_i \leq 2(g + 1)$ . The poles can be only in vertices, edge centers or face centers, hence  $v_p(R_i) + e_p(R_i) + f_p(R_i) = n_i$  and  $0 \leq e_p(R_i) \leq n_i$ . Obviously  $E_p(R_i) = v_p(R_i) - e_p(R_i) + f_p(R_i) = n_i - 2e_p(R_i)$ . Therefore the upper bound for  $E_p(R_i)$  is  $n_i$  and the lower bound is  $-n_i$ .  $\square$

**Corollary 1.5.** *If  $\mathcal{P}$  is a convex polyhedron, then  $E_p(R_i) = 2 - 2e_p(R_i) \in \{2, 0, -2\}$ . If  $e_p = 0$  then  $E_p = 2$ , if  $e_p = 1$  then  $E_p = 1$ , and if  $e_p = 2$  then  $E_p = -2$ . If the order of the rotation  $R_i$  is greater than 2, then  $E_p(R_i) = 2$ .*

*Proof.* Every convex polyhedron is homeomorphic (by a radial projection from any point of its interior) to a sphere, which has genus  $g = 0$ . Now  $n_i = 2$  and the formulas follow from the Lemma 1.4.  $\square$

The next tool we need (in order to prove the main result, Theorem 2.1) is the Burnside lemma, a standard tool for calculating the number of orbits.

**Lemma 1.6.** (Burnside lemma) *Let a group  $G$  act on some set  $Q$ . Let  $|G| = n$  denote the number of elements of  $G$  and let  $|Fix(g)|$  denote the number of elements  $a$  of the set  $Q$ , preserved by the given element  $g$  of the group:  $g(a) = a$ . Then the number of orbits  $Qo$  of the set  $Q$  is given by the formula*

$$Qo = \frac{1}{|G|} \sum_{g \in G} |Fix(g)|.$$

If the group of rotational symmetries of a convex polyhedron is the cyclical group  $C_n$  then the exact value for the Euler orbit characteristic  $Eo_R(\mathcal{P})$  can be obtained by a straightforward application of the Burnside lemma. This is a generalization of the similar result for the spherical polyhedra ([4], p.253).

**Proposition 1.7.** *Let  $\mathcal{P}$  be a convex polyhedron. If  $G_R(\mathcal{P}) = C_n$  where  $C_n$  is generated by a rotation  $R_1$  (thus  $R_1^n = I$ ), then  $Eo_R(\mathcal{P}) \in \{0, 1, 2\}$ .*

*Proof.* The identity transformation fixes all vertices, edges and faces while the other  $n - 1$  rotations fix only the poles. Hence we get by the Burnside lemma and using the Euler formula  $v - e + f = 2$  (valid for any convex polyhedron) the following formulas for the numbers of rotational orbits:

$$\begin{aligned} v o_R &= \frac{1}{n}(v + (n - 1)v_p), \\ e o_R &= \frac{1}{n}(e + (n - 1)e_p), \\ f o_R &= \frac{1}{n}(f + (n - 1)f_p), \\ Eo_R &= \frac{1}{n}(2 + (n - 1)E_p(R_1)), \end{aligned}$$

and using  $E_p(R_1) = 2 - 2e_p(R_1)$  we see: if  $e_p(R_1) = 0$  then  $E_p(R_1) = 2$  and  $Eo_R = 2$ ; if  $e_p(R_1) = 1$  then  $n = 2$ ,  $E_p(R_1) = 0$  and  $Eo_R = 1$ ; if  $e_p(R_1) = 2$  then  $n = 2$ ,  $E_p(R_1) = -2$  and  $Eo_R = 0$ . Thus, if  $n > 2$  then  $Eo_R = 2$ . □

## 2 The main result

**Theorem 2.1.** *Let  $\mathcal{P}$  be a polyhedron with faces on the surface  $S$  of genus  $g$ . Then*

$$Eo_R(\mathcal{P}) = \frac{1}{n}(\chi(\mathcal{P}) + \sum_{i=1}^{n-1} E_p(R_i)),$$

and we get the following bounds on  $Eo_R(\mathcal{P})$ :

$$\frac{1}{n}(\chi(\mathcal{P}) - \sum_{i=1}^{n-1} 2(g + 1)) \leq Eo_R(\mathcal{P}) \leq \frac{1}{n}(\chi(\mathcal{P}) + \sum_{i=1}^{n-1} 2(g + 1)).$$

If  $\mathcal{P}$  is a convex polyhedron, then

$$-1 \leq Eo_R(\mathcal{P}) \leq 2.$$

*Proof.* Let  $n$  be the number of elements in the group  $G_R(P)$ .

The identity transformation fixes each vertex, edge or face.

Every rotation  $R_i$  fixes  $v_p(R_i)$  vertices,  $e_p(R_i)$  edges and  $f_p(R_i)$  faces.

Therefore, by the Burnside lemma:

$$v_{o_R} = \frac{1}{n}(v + v_p(R_1) + \cdots + v_p(R_{n-1})),$$

$$e_{o_R} = \frac{1}{n}(e + e_p(R_1) + \cdots + e_p(R_{n-1})),$$

$$f_{o_R} = \frac{1}{n}(f + f_p(R_1) + \cdots + f_p(R_{n-1})),$$

$$E_{o_R} = \frac{1}{n}(\chi + E_p(R_1) + \cdots + E_p(R_{n-1})).$$

Using  $-2(g+1) \leq E_p(R_i) \leq 2(g+1)$  (proved in Lemma 1.4) we get

$$\frac{1}{n}(\chi - \sum_{i=1}^{n-1} 2(g+1)) \leq E_{o_R} \leq \frac{1}{n}(\chi + \sum_{i=1}^{n-1} 2(g+1)).$$

If  $\mathcal{P}$  is a convex polyhedron, then  $g = 0$ ,  $\chi = 2$ , hence

$$E_{o_R} \leq \frac{1}{n}(2 + (n-1)2) = 2,$$

$$E_{o_R} \geq \frac{1}{n}(2 + (n-1)(-2)) = \frac{1}{n}(4 + n(-2)) \geq -2 + \frac{4}{n} \geq -1,$$

because  $\frac{4}{n} > 0$  and  $E_{o_R}$  must be an integer. □

Thus there are 4 classes  $C_2, C_1, C_0, C_{-1}$  of convex polyhedra, whose  $E_{o_R}$  are 2, 1, 0,  $-1$ , respectively.

Is the lower bound  $E_{o_R} = -1$  actually obtained, and (if it is so) for which convex polyhedra? And is there any simple description of these four classes?

**Proposition 2.2.** *Let  $a, b, c$  be the numbers of rotations  $R_i$  in the group  $G_R(\mathcal{P})$  of a convex polyhedron for which  $E_p(R_i)$  equals 2, 0 and  $-2$ , respectively, and let  $n$  be the number of elements in  $G_R(\mathcal{P})$ . Then*

$$E_{o_R}(\mathcal{P}) = \frac{1}{n}(2 + a \cdot 2 + c(-2)) = \frac{2}{n}(1 + a - c).$$

*Thus the number  $1 + a - c$  is an integer multiple of  $\frac{n}{2}$ . The numbers  $a$  and  $c$  can be (for each of the 4 possible values of  $E_{o_R}$ ) expressed by  $b, n$  and  $E_{o_R}$ .*

*Proof.* This formula follows immediately from the Burnside lemma. Also, it is clear that

$$a + b + c + 1 = n,$$

hence

$$a + c = n - b - 1.$$

The equation  $\frac{2}{n}(1 + a - c) = Eo_R$  implies  $2(1 + a - c) = Eo_R \cdot n$  and

$$c - a = 1 - \frac{Eo_R \cdot n}{2}.$$

Then  $(a + c) + (c - a) = 2c = n - b - 1 + 1 - \frac{Eo_R \cdot n}{2} = \frac{(2 - Eo_R) \cdot n}{2} - b$  and

$$c = \frac{(2 - Eo_R)n - 2b}{4}.$$

Similarly,  $2a = n - b - 1 - 1 + \frac{Eo_R \cdot n}{2}$ , hence

$$a = \frac{n \cdot (2 + Eo_R) - (2b + 4)}{4}.$$

For example, if  $Eo_R = -1$  and  $b = 0$  then  $a = \frac{n}{4} - 1$  and  $c = \frac{3n}{4}$ . In that case  $n$  must be divisible by 4. □

**Example 2.3.** To find such a solid with 4 symmetries we have to look for one having three rotations with poles in edge centers! The lower bound  $Eo_R = -1$  is really obtained for the Johnson solid J84 (Snub Disphenoid, see Figure 1), where  $vo_R = 2, eo_R = 6, fo_R = 3$ , hence  $Eo_R = 2 - 6 + 3 = -1$ . Here the number of symmetries is 4 (the identity transformation and 3 rotations of order two with axes going through edge centers),  $b = 0, a = 0$  and  $c = 3$ . Thus this lower bound  $-1$  is sharp.




Figure 1: The Johnson solid J84, also known as the Snub Disphenoid.

**Remark 2.4.** A rotational axis of a non-convex polyhedron may have more than 2 "poles". As a consequence, there is no upper or lower bound for  $Eo_R$  in the non-convex case.

### 3 Classification of convex polyhedra

As an immediate consequence of the formulas in Proposition 3 we get:

**Corollary 3.1.** *The four classes  $C_2, C_1, C_0, C_{-1}$  of convex polyhedra (whose  $Eo_R$  are  $2, 1, 0, -1$ , respectively) can be characterized as follows:*

$$C_2: a - c = n - 1$$

$$C_1: a - c = -1 + \frac{n}{2}$$

$$C_0: a - c = -1$$

$$C_{-1}: a - c = -1 - \frac{n}{2}, \text{ all poles in edge centers, } a = 0.$$

**Corollary 3.2.** *If  $a \geq n/2$  then  $Eo_R \in \{1, 2\}$ .*

*Proof.* The relation  $a \geq n/2$  is a sufficient condition for  $a - c > 0$  (since there is also the identity transformation in the group  $G_R(\mathcal{P})$ ) that holds only for polyhedra from  $C_2$  and  $C_1$ . □

**Corollary 3.3.** *Let  $q$  be the number of all rotations  $R_i \in G_R(\mathcal{P})$  with the property that  $R_i$  has the same rotational axis as some  $k$ -fold rotation for any  $k > 2$ . If  $q \geq n/2$ , where  $n$  is the order of the group  $G_R$ , then  $Eo_R(\mathcal{P}) \in \{1, 2\}$ .*

*Proof.* No rotation with such an axis can have any of its two poles in an edge center, hence  $a \geq q \geq n/2$ , therefore  $Eo_R(\mathcal{P}) \in \{1, 2\}$ . □

Now we can classify convex polyhedra with respect to their rotational symmetry groups and their rotational orbit Euler characteristic.

The only possible rotational groups of the Euclidean space  $E^3$  are the rotational groups of 1) the  $n$ -gonal pyramid, 2) the  $n$ -gonal dipyrmaid or prism, 3) the regular tetrahedron, 4) the cube or the regular octahedron, 5) the regular dodecahedron or the regular icosahedron ([1], p.34).

**Theorem 3.4.** *Convex polyhedra with at least one rotational symmetry can be classified by their  $G_R$  and by their  $Eo_R$  into 13 classes (in Table 2 the impossible cases are marked with  $\emptyset$ ):*

	$C_2$	$C_1$	$C_0$	$C_{-1}$
$C_n$ cyclical group				$\emptyset$
$D_n$ dihedral group				
$T$ tetrahedron group			$\emptyset$	$\emptyset$
$O$ octahedron group			$\emptyset$	$\emptyset$
$D$ dodecahedron group			$\emptyset$	$\emptyset$

Table 2: Classification of convex polyhedra by  $G_R$  and  $Eo_R$ .

*Proof.* If  $G_R = C_n$  then  $Eo_R \neq -1$  (by Proposition 2). If  $G_R \in \{T, O, D\}$  then  $q \geq n/2$  (Table 3) hence  $\mathcal{P}$  cannot be in  $C_0$  or  $C_{-1}$  (by Corollary 3.3). □

	2	3	4	5	$n$	$q$
$T$ tetrahedron	3	4			$3 + 4 \cdot 2 + 1 = 12$	9
$O$ cube	6	4	3		$6 + 4 \cdot 2 + 3 \cdot 3 + 1 = 24$	17
$D$ dodecahedron	15	10		6	$15 + 10 \cdot 2 + 12 \cdot 4 + 1 = 60$	44

Table 3: Numbers of the 2-,3-,4-,5-fold axes in the solids  $\mathcal{P}$  with the rotational groups  $T$ ,  $O$  or  $D$ , orders  $n$  of the groups  $T$ ,  $O$ ,  $D$  and the numbers  $q$  of all rotations  $R_i \in G_R(\mathcal{P})$  with the property that  $R_i$  has the same rotational axis as some  $k$ -fold rotation for any  $k > 2$ .

## References

- [1] H. S. M. Coxeter, W.O.J. Moser, *Generators and Relations for Discrete Groups*, Springer Verlag, Berlin, Göttingen, Heidelberg, 1957.
- [2] N. W. Johnson, Convex polyhedra with regular faces, *Canad. J. Math.* **18** (1966), 169–200.
- [3] J. Kovič, Symmetry-type graphs of Platonic and Archimedean solids, *Math. Commun.* **16** (No 2) (2011), 491–507.
- [4] J. Kovič, Characterization of convex polyhedra with regular polygonal faces by minimal number of parameters, *J. Combin. Math. Combin. Comput.* **89** (2014), 249–263.
- [5] J. Stillwell, *Mathematics and Its History*, Springer, New York, 2010.