

**Symbiotic efficiency, biosorption and the growth of rhizobia  
on Horse gram plants under aluminium stress**

Učinkovitost simbioze, biosorpcije in rasti rizobijev  
pri vrsti *Macrotyloma uniflorum* zaradi aluminijevega stresa

Prabhavati Edulamudi<sup>a</sup>, Anthony Johnson Anthony Masilamani<sup>c</sup>, Venkata Ramana Sai Gopal Divi<sup>b</sup>,  
Vishnuvardhan Zakkula<sup>a</sup>, Umamaheswara Rao Vanga<sup>a</sup>, Veera Mallaiiah Konada<sup>a</sup>

<sup>a</sup>Department of Botany and Microbiology, Acharya Nagarjuna University, Guntur,  
Andhra Pradesh, India.

<sup>b</sup>Department of Virology, Sri Venkateswara University, Tirupati, Andhra Pradesh, India.

<sup>c</sup>Department of Botany, Sri Krishnadevaraya University, Ananthapur, Andhra Pradesh, India.

\*correspondence: prabha\_anumicro@rediffmail.com

**Abstract:** The aim of the present study was to evaluate the tolerance potential of Horse gram rhizobia to aluminium (Al) toxicity, the enhancement in pod formation, symbiotic efficiency and biosorption potential in the rhizobia inoculated Horse gram (*Macrotyloma uniflorum* (Lam.) Verdc.) plants. Initially, 32 isolates of Horse gram rhizobia were screened for their tolerance of Al in growth media. Among the 32 strains, HGR 4, 6, 13 and 25 that were more tolerant were inoculated individually to Horse gram plants and the plants were then screened for the ability of pod formation, symbiotic efficiency and biosorption potential. Among them, maximum pod formation was observed in Horse gram upon inoculation with HGR-6 and grown at 400  $\mu\text{g g}^{-1}$  of Al. Maximum nodulation was observed in Horse gram upon inoculation with HGR-6 and HGR-13 grown at 200  $\mu\text{g g}^{-1}$  Al. Leghaemoglobin content was maximum on inoculation with HGR-13 at 400  $\mu\text{g g}^{-1}$  of Al. The strain HGR-13 has shown biosorption potential in soil and as well as in root nodules even at 300  $\mu\text{g g}^{-1}$  of Al though it was maximum at 100  $\mu\text{g g}^{-1}$ . This study demonstrated that the Horse gram plants inoculated with *Rhizobium* strains HGR - 4, 6, 13 and 25, besides having nitrogen fixing ability also have the ability to grow in Al contaminated soils. Hence, Horse gram plants associated with these strains of rhizobia could be used in phytoremediation of metal (Al) contaminated soils.

**Key words:** Aluminium, biosorption, metal tolerance, phytoremediation, *Rhizobium*, symbiotic efficiency.

**Izveček:** Namen študije je bil ovrednostiti tolerančni potencial rizobijev za aluminij pri vrsti *Macrotyloma uniflorum* (Lam.) Verdc. Preučevali smo tvorbo strokov, učinkovitost simbioze in biosorpcijski potencial 32 izolatov rizobijev. Med njimi so bili najbolj tolerantni sevi HGR 4, 6, 13 in 25, ki smo jih v nadaljevanju inokulirali individualno na rastline in spremljali 3 parametre. Največjo tvorbo strokov smo izmerili pri sevu HGR-6 in rasti pri 400  $\mu\text{g g}^{-1}$  Al. Največ nodulov je bilo pri inokulaciji s HGR-6 in HGR-13 pri 200  $\mu\text{g g}^{-1}$  Al. Vsebnost leghemoglobina je bila najvišja pri inokulaciji s HGR-13 pri 400  $\mu\text{g g}^{-1}$  Al. Sev HGR-13 je imel biosorpcijski potencial tako v tleh kot v koreninskih nodulih celo pri 300  $\mu\text{g g}^{-1}$  Al, največji pa je bil pri

100  $\mu\text{g g}^{-1}$  Al. Raziskava kaže, da imajo rastline vrste *Macrotyloma uniflorum*, ki so inokulirane s sevi *Rhizobium* HGR - 4, 6, 13 in 25 poleg sposobnosti vezave dušika tudi sposobnost rasti v tleh, onesnaženih z Al. Zato so lahko take rastline primerne za fitoremediacijske postopke s kovinami (Al) onesnaženih tal.

**Ključne besede:** aluminij, biosorpcija, toleranca za kovine, fitoremediacija, *Rhizobium*, učinkovitost simbioze.

## Introduction

Contamination of soils by metals is wide spread due to human, agricultural and industrial activities (Beladi et al. 2011). These activities result in the accumulation of traces of metals in agricultural soils which pose a threat for food safety and public health (Dary et al. 2010). This accumulation of metals leads to loss of soil fertility, since the composition of microbial flora and microbial activities are affected severely (Krujatz et al. 2011). Some metals, though essential in micro-quantities for organisms, are harmful in excess. Aluminium (Al), the third most abundant element in the earth's crust after oxygen and silicon, comprises approximately 7% of its mass (Foy et al. 1978). Al toxicity limits world's agricultural productivity, as Al becomes more soluble in acidic conditions and is the major toxic element in acidic soils and water (Sledge et al. 2005). Excess of Al resists the crop as a result of direct inhibition of nutrient uptake or impairment in root cell function (Kochian 1995, Matsumoto 2000) and productivity of plants (Kochian et al. 2004).

Focus on use of legume plants associated with microorganisms for bioremediation of metals is growing everyday (Carrasco et al. 2005). The *Rhizobium*-legume association has an advantage in which both the microorganisms and plants may influence metal solubility, bioavailability. Added to this is an advantage of enhanced nitrogen content of soils when used for bioremediation or phytoremediation (Pajuelo et al. 2008, Dary et al. 2010). Therefore, isolation of rhizobia, capable to tolerate these metal stress is essential for efficient metal remediation and also the nitrogen fixation under metal pollution (Woldeyohannes et al. 2007). The significance in choice of Horse gram for our study is its adaptability to poor adverse climatic conditions, which are unsuitable for other pulse crops. Horse gram is cultivated as a grain legume and

fodder crop in the states of Tamil Nadu, Karnataka, Andhra Pradesh and Orissa of South India during Kharif and Rabi (the two agricultural monsoon seasons). Our study targets to analyze the effect of Al tolerant *Rhizobium* strains on pod formation, symbiotic efficiency content and Al biosorption potential of Horse gram plants upon inoculation with the chosen rhizobial strains.

## Materials and methods

### *Isolation and analysis of rhizobial strains*

Soil samples were collected from various regions in united Andhra Pradesh, India for sowing the plants. Root nodules were isolated from these plants. Root nodules were surface sterilized and the rhizobial strains were further isolated on Yeast Extract Mannitol (YEM) agar medium (Vincent 1970) with 0.0025% Congo red dye. All these isolates were subjected to biochemical and 16S rRNA sequence analysis. YEM agar medium was prepared with varying concentrations of  $\text{Al}_2(\text{SO}_4)_2$  i.e. 50, 100, 200, 300, 500, 750 and 1000  $\mu\text{g g}^{-1}$ . After solidification, all the isolates were inoculated and incubated at room temperature for 72 hrs. After incubation, the colony diameter was monitored. Replicates were maintained for each metal concentration. Initially 32 isolates of Horse gram rhizobia (HGR's) were screened for the tolerance of aluminium.

### *Inoculation with rhizobial strains grown under different concentrations of $\text{Al}_2(\text{SO}_4)_2$*

Seeds of Horse gram used during the study were obtained from local fields of Andhra Pradesh, India. The pots of the study were filled with soil sterilized in an autoclave at 121°C for 3 hrs each on three alternative days. Horse gram seeds were

surface sterilized with 70% ethanol for 3 min, followed by sodium hypochlorite treatment for 3 min and then rinsed six times with sterilized water, dried. The rhizobial suspension of isolates used in the study were grown in YEM broth in flasks shaken at 120 rpm at  $28 \pm 2^\circ\text{C}$  for 3 days to obtain a cell density of  $6 \times 10^9$  cells  $\text{ml}^{-1}$ . Horse gram plants were inoculated with the selected strains HGR-4 (GQ483457), HGR-6 (GQ483458), HGR-13 (GQ483459) and HGR-25 (GQ483460) which performed well during the initial screening on Al tolerance assay. To perform the inoculations, sterilized seeds were coated with the rhizobial strain by soaking the seeds in liquid culture medium for 2 hrs using 10% (wt/vol) gum Arabic as adhesive to deliver approximately  $10^9$  cells  $\text{seed}^{-1}$ . Respective controls were maintained with seeds treated in sterilized distilled water. The inoculated seeds (20 seeds  $\text{pot}^{-1}$ ) were sown in clay pots using 2 kg sterilized soil. In order to evaluate the effect of Al metal on the Horse gram plants inoculated with 4 HGR strains, plants were maintained with  $\text{Al}_2(\text{SO}_4)_3$  supplements of 50, 100, 200, 300 and 400  $\mu\text{g g}^{-1} \text{kg}^{-1}$ . Negative controls were also maintained using the Horse gram plants natively available with  $\text{Al}_2(\text{SO}_4)_3$  supplements of 50, 100, 200, 300 and 400  $\mu\text{g g}^{-1} \text{kg}^{-1}$ . For the purpose of comparison controls were also maintained without adding Al. Three replicates were maintained for each treatment. The pots were watered regularly and were maintained in an open field conditions and allowed to grow.

#### *Analysis of plants for nodulation, symbiotic efficiency and biosorption potential*

The number of pods formed were counted post 40 days of sowing. The plants were observed for nodulation regularly after the seedlings emerged. Five plants in each treatment were picked up randomly and nodulation characteristics were evaluated 40 days after sowing, as it was previously observed during the study that highest nodulation of Horse gram occurred on 40<sup>th</sup> day. For biochemical analysis, plants raised in different concentrations of Al were collected, the amount of leghaemoglobin was estimated (Tu et al. 1970) post 40 days of sowing. Soil pH, organic matter and total nitrogen (N) (Jackson 1973) and total phosphorus (P) (Olsen et al. 1954) were also es-

timated. The amount of sand, silt and clay present in the soil were also analyzed (Black 1965).

For elemental analysis, root nodules were collected and washed under tap water to remove sediments and soil. Then they were washed in 0.02% detergent (tween-20) and once again in tap water. They were again washed with 0.1 N HCl. Finally, the nodules were washed twice with distilled water. The nodules were dried at  $80^\circ\text{C}$  for 48 hrs in hot air oven and they were ground to a very fine powder. From this, 0.5 grams of powdered tissue was added to 5 ml of conc.  $\text{HNO}_3$  for cold digestion at room temperature. Then 5 ml of conc.  $\text{HNO}_3$  and  $\text{H}_2\text{O}_2$  were added to the digested sample in 10:4 ratio, the samples were heated to a volume of 2 ml. The clear solution obtained was made up to 25 ml with deionized water (millipore) and used for elemental analysis. Soil samples were also subjected to acid digestion with slight modifications and were used in elemental analysis. Al concentration present in the sample was determined (APHA 22<sup>nd</sup> Edition, 3111 B) by Atomic Absorption Spectroscopy (AAS) (THERMO AAS Model No: ICE 3000). The system was operated using the Thermo scientific SOLAAR data station V 11.02 software. Argon was used as inert gas during operation. The instrument's operating conditions included Furnace instrumental mode, Lamp current at 15 mA, Wavelength of 232 nm, 0.2  $\mu\text{g/l}$  Gas flow, 0.2 nm band width and 72 sec of Furnace programme total time.

#### *Statistical analysis*

Statistical analysis was done in three replicates for each treatment. The mean and standard error (SE) were calculated using Microsoft Office Excel 2007. To test the statistical significance, all the values were analyzed by ANOVA, using SPSS Statistics, Version 20 (Armonk, 2011). Bars indicate means  $\pm$  SE and were significant at 5% level of significance ( $P$  value  $< 0.05$ ).


## **Results**

Most of the 32 rhizobial isolates were able to grow on YEM agar plates containing 1000  $\mu\text{g g}^{-1}$  concentration of Al, but the diameter of the colonies varied with the isolate i.e. from 2 mm (HGR-16)

to 14 mm (HGR-13). At this concentration the colonies are round, white, translucent, raised and convex with entire margins. Similar results were obtained in YEM broth also. Hence four isolates of the study HGR - 4, 6, 13 and 25 which performed better were further used for evaluation of other parameters.


Al supported good growth of Horse gram plants up to  $300 \mu\text{g g}^{-1}$ , when the plants were inoculated with the four rhizobial strains. Later, the growth of the plants decreased with increase of metal concentration. The plants inoculated with the strains HGR-4, 6, 13 and 25 have shown their maximum pod formation at 100 to  $400 \mu\text{g g}^{-1}$  of Al. Horse gram rhizobial inoculation increased pod formation when compared to control. Among the four selected strains the strain HGR-4 inoculated plants have shown maximum pod formation at  $200 \mu\text{g g}^{-1}$ , HGR-6 at  $400 \mu\text{g g}^{-1}$ , HGR-13 at  $300 \mu\text{g g}^{-1}$  and HGR-25 has shown at  $100 \mu\text{g g}^{-1}$  of Al only (Fig. 1). They were significant at 5% level ( $P$  value  $< 0.05$ ).

In the present study, nodules appeared post 13 days of sowing on tap root and as well as on lateral roots at all the Al concentrations experimented. The total number of nodules formed per plant ranged from 8 to 22. The plants inoculated with the strain HGR-6 and 13 have shown highest nodulation at  $200 \mu\text{g g}^{-1}$  of Al. The number of nodules formed were more when rhizobia inoculated to Horse gram plants when compared to control. They were significant at 5% level ( $P$  value  $< 0.05$ ). However, the increased concentration of Al reduced the number of nodules. HGR-4 and 25 inoculated plants had maximum number of nodules at  $100 \mu\text{g g}^{-1}$  of Al only (Fig. 2). The leghaemoglobin content was maximum in the Horse gram plants inoculated with the strains HGR-6 and HGR-13 at  $400 \mu\text{g g}^{-1}$  of Al (Fig. 3). The plants inoculated with HGR-4 and HGR-25 have shown their maximum at  $50 \mu\text{g g}^{-1}$  only. But, these values were more than in control. They were significant at 5% level ( $P$  value  $< 0.05$ ).




**Figure 1:** Formation of pods on Horse gram plants inoculated with the four *Rhizobium* strains in response to varying concentrations of aluminium. Data show mean value  $\pm$  SE.

**Slika 1:** Tvorba strokov pri rastlinah *Macrotyloma uniflorum*, inokuliranih s štirimi sevi rizobija pri različnih koncentracijah aluminija. Prikazane so povprečne vrednosti  $\pm$  SN.



**Figure 2:** Formation of root nodules on Horse gram plants inoculated with the four *Rhizobium* strains in response to varying concentrations of aluminium. Data show mean value  $\pm$  SE.

**Slika 2:** Tvorba koreninskih nodulov pri rastlinah *Macrotyloma uniflorum*, inokuliranih s štirimi sevi rizobija pri različnih koncentracijah aluminija. Prikazane so povprečne vrednosti  $\pm$  SN.




**Figure 3:** Leghaemoglobin content in root nodules of Horse gram plants inoculated with the four *Rhizobium* strains in response to varying concentrations of aluminium. Data show mean value  $\pm$  SE.

**Slika 3:** Vsebnost leghemoglobina v koreninskih nodulih pri rastlinah *Macrotyloma uniflorum*, inokuliranih s štirimi sevi rizobija pri različnih koncentracijah aluminija. Prikazane so povprečne vrednosti  $\pm$  SN.

The amount of total nitrogen (%) and phosphorus (%) present in the soil is 0.85 and 1.24 respectively. The total content of organic matter in soil is 1.20, sand 18, silt 16, clay 42 and the pH of the soil is 6.44. Among these four strains HGR-13

has shown more number of nodules and maximum leghaemoglobin content at high concentrations of Al. So, we have selected HGR-13 to estimate biosorption potential of the Horse gram rhizobia. Biosorption potential of the strain HGR-13 was



**Figure 4:** Biosorption potential of the strain HGR-13 in soil after deplantation of Horse gram plants at different concentrations of aluminium. Data show mean value  $\pm$  SE.

**Slika 4:** Biosorpcijski potencial seva HGR-13 po presaditvi rastlin *Macrotyloma uniflorum*, v tla z različno koncentracijo aluminija. Prikazane so povprečne vrednosti  $\pm$  SN.

determined by Atomic Absorption Spectroscopy (AAS), as the amount of metal present in the nodule and soil samples after the treatment with the isolate HGR-13. After analyzing the treated samples in AAS, the inoculated strain HGR-13 has shown maximum biosorption of Al. The results have shown that the Al present in the soil was decreased after deplantation of Horse gram plants when they were inoculated with the Al tolerant rhizobia (HGR-13) i. e. from 5.74 mg/L (control without inoculation) to 3.91 mg/L at 100  $\mu\text{g g}^{-1}$ , 4.8 mg/L to 3.69 mg/L at 200  $\mu\text{g g}^{-1}$  and 3.39 mg/L to 2.95 mg/L at 300  $\mu\text{g g}^{-1}$  of Al concentration. In nodules, the biosorption potential was maximum at 200  $\mu\text{g g}^{-1}$  (1.69 mg/L to 0.99 mg/L), at 100  $\mu\text{g g}^{-1}$  it was 1.92 mg/L to 1.89 mg/L and at 200  $\mu\text{g g}^{-1}$  it was 1.28 mg/L to 1.27 mg/L. The results have shown that the isolate was able to adsorb Al at a concentration of 300  $\mu\text{g g}^{-1}$  in root nodules and also in soil samples inoculated with the strain HGR-13 (Fig. 4, 5).

## Discussion

In our study all the tested Horse gram rhizobia are able to show growth at all the Al concentrations tested. But, they have shown variations in their

growth. Al stress reduced the growth of rhizobial species under laboratory conditions and also in natural environment (Paudyal et al. 2007; Avelar Ferreira et al. 2012). Some cowpea rhizobia tolerate up to 20  $\text{m mol dm}^{-3}$  of  $\text{Al}^{3+}$  (Bruno et al. 2013). *Rhizobium* species like *Sinorhizobium meliloti* and *Bradyrhizobium* growing explanta were extremely sensitive to Al (Arora et al. 2001). The rhizobia from Bambara groundnut were able to grow in a medium pH as low and Al concentration of 50  $\mu\text{M}$  (Laurette et al. 2015). The growth of the *Rhizobium* strain CB756 was also not affected by 50  $\mu\text{M}$  Al at pH 4.5 (Keyser and Munns, 1979). But 50  $\mu\text{M}$  Al at pH 4.6 decreased the growth of *Bradyrhizobium* spp. (Shamsuddin 1987) and the growth of rhizobia was completely inhibited at 50 mM Al (Broos et al. 2004). At high concentrations Al is a potent inhibitor of rhizobial growth. It was observed that root nodule bacteria grow after a lag period in culture medium containing 75  $\mu\text{M}$  Al (Whelan and Alexander 1986). Wild rhizobia from Soybean have shown growth at 200  $\mu\text{mol L}^{-1}$  of  $\text{Al}^{3+}$  (Ping et al. 2014). It has shown negative effect on the survival and growth of *Rhizobium trifolii*. *Bradyrhizobium* strains were able to show growth in the presence of 100  $\mu\text{M}$   $\text{AlCl}_3$  (Lesueur et al. 1993) and some *Bradyrhizobium japonicum* strains were tolerant to Al (Tayler et al. 1991).

*Anoxybacillus* sp. SK 3-4 was found to be the most resistant to Al and significant growth was observed at 300 mg L<sup>-1</sup> to 800 mg L<sup>-1</sup> of Al (Lim et al. 2015). To the best of our knowledge this is the first report that rhizobia nodulating Horse gram were able to grow in culture medium containing 1000 µg g<sup>-1</sup> Al concentration.

Al effects the growth and nodulation of many legumes (Kim et al. 1985). Kushwaha et al. 2017 observed in a study with cowpea, the number of pods formed were more in soil upon treatment with 40 ppm concentration of Al. In the present study the number of pods formed was more at 100 to 400 µg g<sup>-1</sup> of Al. The inoculation of acid tolerant, Al tolerant *Bradyrhizobium japonicum* strains could increase number of pods in soybean (Situmorang et al. 2009). Previous studies reported that at high Al concentrations, nodulation was partially or totally inhibited in several species such as common bean (*Phaseolus vulgaris*), clover (*Trifolium repens*), *Stylosanthes* species and also in other tropical legumes (de Carvalho et al. 1981, Paudyal et al. 2007). de Carvalho et al. 1981 reported that Al has shown its effect on reduction or inhibition of nodulation at 25 iM concentration. Kushwaha et al. 2017 during their study reported that the more number of nodules formed at lower concentrations of Al (0 and 20 ppm) and number of nodules were reduced at higher concentration of Al. Brady et al. 1990 reported that Al at < 5 µM has shown reduction in nodulation of soybean. Inoculation of *Sinorhizobium mexicanum* ITTG 27<sup>T</sup> to *Acaciella angustissima* plants enhanced nitrogen content (Rosales et al. 2011). Al toxicity has shown negative effect on symbiotic nitrogen fixation in common bean plants that were grown hydroponically in acidic nutrient solution containing 70 µM AlCl<sub>3</sub> (Mendoza-Soto et al. 2015). Blamey and Chapman, 1982 reported that in groundnut, poor nodulation and nitrogen fixation was observed under Al toxicity. Horse gram rhizobia i.e. HGR-6 and HGR-13 inoculated plants have shown increased leghaemoglobin content even under 400 µg g<sup>-1</sup> of Al stress. de Carvalho et al. 1982 suggest that the effects of Al on N<sub>2</sub> fixation may be indirect. The rhizospheric microorganisms have intrinsic ability to reduce/detoxify the metal stress by several mechanisms. These mechanisms include the efflux of metal ions outside the cell, biostimulation, bioaugmentation,

metal reduction and biosorption (Outten et al. 2000). Mammari et al. 1997 proved that the metal concentration decreased after rhizobial inoculation, which shows the ability of their rhizobial strain RP5 in the removal of metals through adsorption-desorption mechanism. Bacterial biosorption/bioaccumulation mechanisms together with other plant growth promoting features accounted for improved plant growth in metal contaminated soils (Zaidi et al. 2006). Horse gram rhizobia (HGR-13) have the ability to remove Al even at 300 µg g<sup>-1</sup> Al, even though it was maximum at 100 µg g<sup>-1</sup> in the inoculated plants when compared to control plants. Hence, rhizobia nodulating their hosts may increase metal accumulation in root nodules and may lead to chelation, immobilization and biosorption (Hao et al. 2014).

## Conclusions

Results clearly show that the accumulation of Al in soils reduced upon inoculation of Horse gram plants with Horse gram rhizobia of the current study. The present study demonstrated that the Horse gram plants inoculated with Al tolerant *Rhizobium* strains HGR-4, 6, 13 and 25 besides having nitrogen fixing capacity also have the ability to grow in Al contaminated soils. Hence, these Horse gram plants upon inoculation of the rhizobia associated with them during the study i.e. HGR 4, 6, 13 and 25 could be used in phytoremediation of metal from Al contaminated soils.

## Acknowledgements

The first author E.P. is grateful to University Grants Commission (UGC), New Delhi for financial assistance under Post-Doctoral Fellowship (PDF). Dr. Anthony Johnson is thankful to Universal Grants Commission (UGC) for Dr. D.S. Kothari Post-Doctoral Fellowship (201617-BL/16-17/0364).



## References

- Armonk, N.Y., 2011. IBM Corp. IBM SPSS Statistics for windows, Version 20.0.
- Arora, N.K., Kumar, V., Maheswari, D.K., 2001. Constraints, development and future of the bio inoculants with special reference to rhizobial inoculants. In: Maheswari, D.K., Dubey, R.C. (ed.): Innovative approaches in Microbiology. Dehradun, India. pp. 241-254.
- Avelar Ferreira, P.A., Bomfeti, C.A., Lima Soares, B., de Souza Moreira, F.M. 2012. Efficient nitrogen fixing strains isolated from Amazonian soils are highly tolerant to acidity and aluminium. World Journal of Microbiology and Biotechnology, 28, 1947-1959.
- Beladi, M., Habibi, D., Kashani, A., Paknejad, F., Nooralvandi, T., 2011. Phytoremediation of Lead and Copper by Sainfoin (*Onobrychis vicifolia*): Role of antioxidant enzymes and biochemical biomarkers. American-Eurasian Journal of Agriculture and Environmental Science, 3, 440-449.
- Black, C.A., 1965. Methods of Soil Analysis. Part. 2. Chemical and biological properties. American Society of Agronomy. Madison, Wisconsin, USA.
- Blamey, F.P.C., Chapman, J., 1982. Soil amelioration effects on peanuts growth, yield and quality. Plant and Soil, 65, 319-334.
- Brady, D.J., Hetch-Buchholz, C.H., Asher, C.J., Edwards, D.G., 1990. Effect of low activities of aluminium on Soybean (*Glycine max* L.) early growth and nodulation. In: Van Blusichem, M.L. (ed.): Plant Nutrient Physiology and Application. Kluwer, Dordrecht, pp. 329-334.
- Broos, K., Uyttebroek, M., Mertens, J., Smolders, E., 2004. A survey of symbiotic nitrogen fixation by white clover grown on metal contaminated soils. Soil Biology and Biochemistry, 36, 633-640.
- Bruno, L.S., Paulo, A.A.F., Silvia Maria, de O.L., Leandro, M.M., Marcia, R., Messias Jose, B.de.A., Farima Maria, de.S.M., 2013. Cowpea symbiotic efficiency, pH and aluminium tolerance in nitrogen-fixing bacteria. Science and Agriculture, 71, 171-180.
- Carrasco, J.A., Armario, P., Pajuelo, E., Burgos, A., Caviedes, M.A., Lopez, R., Chamber, M.A., Palomares, A.J., 2005. Isolation and characterization of symbiotically effective *Rhizobium* resistant to arsenic and heavy metals after the toxic spill at the *Aznalcollar* pyrite mine. Soil Biology and Biochemistry, 37, 1131-1140.
- Dary, M., Chamber, P.M.A., Palomares, A.J., Pajuelo, E., 2010. "In situ" phytostabilisation of heavy metal polluted soils using *Lupinus luteus* inoculated with metal resistant plant-growth promoting rhizobacteria. Journal of Hazardous Materials, 177, 323-330.
- de Carvalho, M.M., Edwards, D.G., Asher, C.J., Andrew, C.S., 1981. Aluminium toxicity, nodulation and growth of *Stylosanthes* sp. Agronomy Journal, 73, 261-265.
- de Carvalho, M.M., Edwards, D.G., Asher, C.J., Andrew, C.S., 1982. Effect of aluminium on nodulation of two *Stylosanthes* species grown in nutrient solution. Plant and Soil, 64, 141-152.
- Foy, C.D., Chaney, R.L., White, M.C., 1978. The physiology of metal toxicity in plants. Annual Review of Plant Physiology, 29, 511-566.
- Hao, X., Taghavi, S., Xie, P., Orbach, M.J., Alwathnani, H.A., Rensing, C., Wei, G., 2014. Phytoremediation of heavy and transition metals aided by legume rhizobia symbiosis. International Journal of Phytoremediation, 16, 179-202.
- Jackson, M.L., 1973. Soil Chemical Analysis. Prentice Hall of India Ltd. New Delhi, India.
- Khushwaha, J.K., Pandey, A.K., Dubey, R.K., Singh, V., Mailappa, A.S., Singh, S., 2017. Screening of cowpea [*Vigna unguiculata* (L.) Walp.] for aluminium tolerance in relation to growth, yield and related traits. Legume Research, 40, 434-438.
- Keyser, H.H., Munns, D.N., 1979. Tolerance of rhizobia to acidity aluminium and phosphate. Soil Science Society of America Journal, 43, 519-523.
- Kim, M.M., Asher, C.J., Edward, D.G., Date, R.A., 1985. Aluminium toxicity, effect on growth and nodulation of subterranean clover. In: Kuna, H., Kitahara, T., Okuba, T., Shiyonu, M., Sugawata, K., Tajimi, A., Yamaguchi, H. (ed.): Proceedings of the 15<sup>th</sup> International Grassland Congress (Tokyo). Science Society of Japan & Japanese Society of Grassland Science, Japan, pp. 501-503.



- Kochian, L.V., 1995. Cellular mechanism of aluminium toxicity and resistance on plants. Annual Review of Plant Physiology and Plant Molecular Biology, 46, 237-260.
- Kochian, L.V., Hoekenga, O.A., Pineros, M.A., 2004. How do crop plants tolerate acid soils? Mechanisms of aluminium tolerance and phosphorus efficiency. Annual Review of Plant Biology, 55, 459-493.
- Krujatz, F., Harstrick, A., Neortemann, B., Greis, T., 2011. Assessing the toxic effects of nickel, cadmium and EDTA on growth of the plant growth-promoting rhizobacterium *Pseudomonas brassicacearum*. Water, Air and Soil Pollution, Doi: 10.1007/s11270-011-0944-0.
- Laurette, N.N., Maxemilienne, N.B., Henri, F., Souleymanou, A., Kamdem, K., Albert, G., Dieudonne, N., Francois-Xavier, E. 2015. Isolation and screening of indigenous Bambara Groundnut (*Vigna subterranea*) nodulating bacteria for their tolerance to some environmental stresses. American Journal of Microbiology Research, 3, 65-75.
- Lesueur, D., Diem, H.G., Meyer, J.M. 1993. Iron requirement and siderophore production in *Bradyrhizobium* strains isolated from *Acacia mangium*. Journal of Applied Microbiology, 6, 675-682.
- Lim, J.C., Goh, K.M., Shamsir, M.S., Ibrahim, Z., Chong, C.S., 2015. Characterization of aluminium resistant *Anoxybacillus* sp. SK 3-4 isolated from a hot spring. Journal of Basic Microbiology, 55, 514-519.
- Mamaril, J.C., Paner, E.T., Alpante, B.M., 1997. Biosorption and desorption studies of Cr (III) by free and immobilized *Rhizobium* (BJVr 12) cell biomass. Biodegradation, 8, 275-285.
- Matsumoto, H., 2000. Cell biology of aluminum toxicity and tolerance in higher plants. International Review of Cytology, 200, 1-46.
- Mendoza-Soto, A.B., Naya, L., Leija, A., Hernández, G., 2015. Responses of symbiotic nitrogen-fixing common bean to aluminum toxicity and delineation of nodule responsive micro RNAs. Frontiers in Plant Science, 6, 587.
- Olsen, S.R., Cole, C.V., Watanabe, F.S., Dean, L.A., 1954. Estimation of available phosphorus in soil by extraction with sodium carbonate. 19. USDA Circular No. 939.
- Outten, F.W., Outten, C.E., Halloran, T., 2000. Metallo regulatory systems at the interface between bacterial metal homeostasis and resistance. In: Storz, G., Hengge, A.R. (ed.): Bacterial stress responses. Washington, DC, pp. 29-42.
- Pajuelo, E., Roddriguez, L.I.D., Mary, M., Palomares, A.J., 2008. Toxic effects of arsenic on *Sinorhizobium-Medicago sativa* symbiotic interaction. Environmental Pollution, 154, 203-211.
- Paudyal, S.P., Rishi, R.A., Chauhan, S.V.S., Maheshwari, D.K., 2007. Effect of heavy metals on growth of *Rhizobium* strains and symbiotic efficiency of two species of tropical legumes. Scientific World, 5, 27-32.
- Ping, L.X., Peng, W.R., Hai, N., Ying, H.M., 2014. Screening for Al-tolerant *Rhizobium* and a study on its characters. Journal of South China Agricultural University, 35, 50-55.
- Rosales, R.R., Victor, M., Valdiviezo, R., Joaquin, A., Molina, M., Federico, A., Miceli, G., Dendooven, L., 2011. Aluminium tolerance in the tropical leguminous N<sub>2</sub> fixing shrub *Acaciella angustissima* (Mill.) Britton & Rose inoculated with *Sinorhizobium mexicanum*. Gayana Botany, 68, 188-195.
- Shamsuddin, Z.H., 1987. Growth, infectivity and nodulating abilities of some winged bean rhizobia in acid conditions. Perth [thesis]. Murdoch University, Western Australia.
- Situmorang, A.R.F., Mubarak, N.R., 2009. The use of acid-Aluminium tolerant *Bradyrhizobium japonicum* inoculant for Soybean grown on acid soils. Hayati Journal of Biosciences, 16, 157-160.
- Sledge, M.K., Pechter, P., Payton, M.E., 2005. Aluminum tolerance in *Medicago truncatula* germplasm. Crop Science, 45, 200-204.
- Taylor, G.J., 1991. Current views of the aluminium stress response; the physiological basis of tolerance. Current Topics in Plant Biochemistry and Physiology, 10, 57-93.
- Tu, J.C., Ford, R.E., Garu, C.R., 1970. Some factors affecting the nodulation and nodule efficiency in Soy beans infected by soybean mosaic virus. Phytopathology, 60, 1653-1656.

- Vincent, J.M., 1970. A manual for the practical study of the root nodule bacteria. IBP Hand Book No. 15, Blackwell Scientific publications, Oxford.
- Whelan, A.M., Alexander, M., 1986. Effects of low pH and high Al, Mn and Fe levels on the survival of *Rhizobium trifolii* and the nodulation of subterranean clover. *Plant and Soil*, 78, 381-391.
- Woldeyohannes, W.H., Dasilva, M.C., Gueye, M., 2007. Nodulation and nitrogen fixation of *Stylosanthes hamata* in response to induced drought stress. *Arid Land Research and Management*, 21, 157-163.
- Zaidi, S., Usmani, S., Singh, B.R., Musarrat, J., 2006. Significance of *Bacillus subtilis* strain SJ-101 as a bioinoculant for concurrent plant growth promotion and nickel accumulation in *Brassica juncea*. *Chemosphere*, 64, 991-997.