

ANNALES

Anali za istrske in mediteranske študije
Annali di Studi istriani e mediterraneei
Annals for Istrian and Mediterranean Studies
Series Historia Naturalis, 35, 2025, 1





ANNALES

Anali za istrske in mediteranske študije
Annali di Studi istriani e mediterranei
Annals for Istrian and Mediterranean Studies

Series Historia Naturalis, 35, 2025, 1

ISSN 1408-533X
e-ISSN 2591-1783

UDK 5

Letnik 35, leto 2025 številka 1

**UREDNIŠKI ODBOR/
COMITATO DI REDAZIONE/
BOARD OF EDITORS:**

Alessandro Acquavita (IT), Nicola Bettoso (IT), Christian Capape (FR), Darko Darovec, Dušan Devetak, Jakov Dulčić (HR), Edy Fantinato (IT), Serena Fonda Umani (IT), Andrej Gogala, Daniel Golani (IL), Danijel Ivajnskič, Hakan Kabasakal (TR), Mitja Kaligarič, Marcelo Kovačič (HR), Petar Kružić (HR), Lovrenc Lipej, Vesna Mačić (ME), Alenka Malej, Borut Mavrič, Patricija Mozetič, Martina Orlando-Bonaca, Michael Stachowitsch (AT), Francesco Tiralongo (IT), Tom Turk, Al Vrezec

**Glavni urednik/Redattore capo/
Editor in chief:**

Darko Darovec

**Odgovorni urednik naravoslovja/
Redattore responsabile per le scienze
naturali/Natural Science Editor:**

Lovrenc Lipej

Urednica/Redattrice/Editor:

Martina Orlando-Bonaca

Prevajalci/Traduttori/Translators:

Martina Orlando-Bonaca (sl./it.)

**Oblikovalec/Progetto grafico/
Graphic design:**

Dušan Podgornik, Lovrenc Lipej

Tisk/Stampa/Print:

Založništvo PADRE d.o.o.

Izdajatelj/Editori/Published by:Zgodovinsko društvo za južno Primorsko - Koper / Società storica del Litorale - Capodistria®
Inštitut IRRIS za raziskave, razvoj in strategije družbe, kulture in okolja / Institute IRRIS for Research, Development and Strategies of Society, Culture and Environment / Istituto IRRIS di ricerca, sviluppo e strategie della società, cultura e ambiente®**Sedež uredništva/Sede della redazione/
Address of Editorial Board:**Nacionalni inštitut za biologijo, Morska biološka postaja Piran / Istituto nazionale di biologia, Stazione di biologia marina di Pirano / National Institute of Biology, Marine Biology Station Piran
SI-6330 Piran /Pirano, Fornače/Fornace 41, tel.: +386 5 671 2900, fax +386 5 671 2901;
e-mail: annales@mbss.org, internet: www.zdjp.si

Redakcija te številke je bila zaključena 20. 06. 2025.

**Sofinancirajo/Supporto finanziario/
Financially supported by:**

Javna agencija za znanstveno-raziskovalno in inovacijsko dejavnost Republike Slovenije (ARIS)

Annales - Series Historia Naturalis izhaja dvakrat letno.**Naklada/Tiratura/Circulation:**

300 izvodov/copie/copies

Revija Annales, Series Historia Naturalis je vključena v naslednje podatkovne baze / La rivista Annales, series Historia Naturalis è inserita nei seguenti data base / Articles appearing in this journal are abstracted and indexed in: BIOSIS-Zoological Record (UK); Aquatic Sciences and Fisheries Abstracts (ASFA); Elsevier B.V.: SCOPUS (NL); Directory of Open Access Journals (DOAJ).

To delo je objavljeno pod licenco / Quest'opera è distribuita con Licenza / This work is licensed under a Creative Commons BY 4.0.

Navodila avtorjem in vse znanstvene revije in članki so brezplačno dostopni na spletni strani <https://zdjp.si/en/p/annalesshn/>
The submission guidelines and all scientific journals and articles are available free of charge on the website <https://zdjp.si/en/p/annalesshn/>
Le norme redazionali e tutti le riviste scientifiche e gli articoli sono disponibili gratuitamente sul sito <https://zdjp.si/en/p/annalesshn/>

VSEBINA / *INDICE GENERALE* / *CONTENTS*BIOTSKA GLOBALIZACIJA
GLOBALIZZAZIONE BIOTICA
*BIOTIC GLOBALIZATION***Okan AKYOL, Oğuzhan TAKICAK & Hasan TARUN**

A Fugitive Lessepsian Fish in a Sea-Cage Farm in the Aegean Sea: *Stephanolepis diaspros* (Monacanthidae) 1
Ubežni lesepski migrant iz ribogojnice v Egejskem morju: Stephanolepis diaspros (Monacanthidae)

Nicola BETTOSO, Lisa FARESI, Valentina TORBOLI & Jose A. CUESTA

Additional Record of the Pea Crab *Pinnotheres bicristatus* (Brachyura: Pinnotheridae) in the Adriatic Sea 7
Dodatni zapis o pojavljanju stražne rakovice vrste Pinnotheres bicristatus (Brachyura: Pinnotheridae) v Jadranskem morju

Alan DEIDUN, Sarah BAUMANN, Bruno ZAVA & Maria CORSINI-FOKA

Confirming the Occurrence of the Non-Indigenous *Pteragogus trispilus* (Actinopterygii: Labridae) within Maltese Waters 13
Potrditev pojavljanja tujerodne ustnače vrste Pteragogus trispilus (Actinopterygii: Labridae) znotraj malteških voda

Deniz ERGÜDEN, Yusuf Kenan BAYHAN, Sibel ALAGÖZ ERGÜDEN & Deniz AYAS

A New Ichthyological Record and Distributional Update for *Epigonus denticulatus* Dieuzeide, 1950 in Turkish Mediterranean Waters 21
Nov ihtiološki zapis in podatki o razširjenosti rjavega veleokca, Epigonus denticulatus Dieuzeide, 1950 v turških sredozemskih vodah

Sara LADOUL, Farid HEMIDA, Christian REYNAUD & Christian CAPAPÉ

On the Occurrence of Cornish Blackfish *Schedophilus medusophagus* (Osteichthyes: Centrolophidae) from the Maghreb Shore (Southwestern Mediterranean Sea) 27
Potrjena prisotnost meduzojeda Schedophilus medusophagus (Osteichthyes: Centrolophidae) z magrebske obale (jugozahodno Sredozemsko morje)

Christina MICHAIL & Francesco TIRALONGO

First Occurrence of Ariidae in Cypriot Waters – a Major Contribution to Biodiversity 35
Prvo pojavljanje predstavnikov iz družine Ariidae v ciprskih vodah – velik prispevek k biodiverziteti

SREDOZEMSKA HRUSTANČNICE
SQUALI E RAZZE MEDITERRANEE
*MEDITERRANEAN SHARKS AND RAYS***Lovrenc LIPEJ, Riccardo BATTISTELLA, Borut MAVRIČ & Danijel IVAJNŠIČ**

An Insight into the Diet of the Bull Ray, *Aetomylaeus bovinus* (Geoffroy Saint-Hilaire, 1817) in the Northern Adriatic Sea 43
Vpogled v prehranjevalne navade kljunatega morskega goloba, Aetomylaeus bovinus (Geoffroy Saint-Hilaire, 1817) v severnem Jadranu

Cem ÇEVİK, Deniz ERGÜDEN & Deniz AYAS

A New Capture Record of *Alopias superciliosus* Lowe, 1841 from the Turkish Coast (Northeastern Mediterranean) 55
Nov ulov velikooke morske lisice Alopias superciliosus Lowe, 1841 iz turške obale (severovzhodno Sredozemlje)

Cem DALYAN, N. Bikem KESİCİ, Elif YÜCEDAĞ BAKIR, Yunus GÖNÜL & Hakan KABASAKAL

No Longer as Common as its Name: a Review of the Occurrence of *Torpedo torpedo* (Linnaeus, 1758) (Chondrichthyes: Elasmobranchii) in Turkish Waters, with Photographic Evidence 65
Ni več tako pogost kot njegovo ime: pregled pojavljanja okatega električnega skata Torpedo torpedo (Linnaeus, 1758) (Chondrichthyes: Elasmobranchii) v turških vodah s fotografskimi dokazi

Deniz ERGÜDEN, Cemal TURAN, Servet Ahmet DOĞDU & Deniz AYAS

Disc Deformity in a Juvenile Female Brown Ray, *Raja miraletus* (Family: Rajidae), from Northeastern Mediterranean (Türkiye) 73
Deformacija diska pri juvenilni samici modropege raže, Raja miraletus (družina: Rajidae), iz severovzhodnega Sredozemskega morja (Turčija)

Farid HEMIDA, Christian REYNAUD & Christian CAPAPÉ

On an Old Record of the Smalltooth Sand Tiger Shark *Odontaspis ferox* (Chondrichthyes: Odontaspidae) from the Algerian Coast (Southwestern Mediterranean Sea) 83
O starem zapisu o drobn zobem morskem biku Odontaspis ferox (Chondrichthyes: Odontaspidae) z alžirske obale (jugozahodno Sredozemsko morje)

Hakan KABASAKAL, Uğur UZER & F. Saadet KARAKULAK

Plastic Debris-Induced Fin Damage in the Smoothhound, *Mustelus mustelus* 91
Poškodbe plavuti pri navadnem morskem psu, Mustelus mustelus, zaradi plastičnih odpadkov

Nicolas ZIANI, Florane TONDU, Rémi BRU, Chloé MOSNIER, Sarah FOXONET, Ruben Bao GALLIEN, Mathias POULY, Modan Lou TONIETTO, Lucille VERDON, Eloïse DEYSSON, Alessandro DE MADDALENA & Hakan KABASAKAL

Bite Marks Observed on a Large Female White Shark *Carcharodon carcharias* Off Camargue, France Provide Potential Insights into the Reproduction of the Mediterranean Population 97
Sledovi ugrizov na veliki samici belega morskega volka Carcharodon carcharias pri Camargu (Francija) kažejo na možno razmnoževanje sredozemske populacije

**MORSKA FAVNA
FAUNA MARINA
MARINE FAUNA****Sihem RAFRAFI-NOUIRA, RIMEL BENMESSAOUD, Mourad CHÉRIFF, Christian REYNAUD & Christian CAPAPÉ**

Morphological Deformities in a Common Two-Banded Sea Bream, *Diplodus vulgaris* (Osteichthyes: Sparidae), from Northern Tunisian Waters (Central Mediterranean Sea) 109
Morfološke deformacije pri fratru, Diplodus vulgaris (Osteichthyes: Sparidae), iz severnih tunizijskih vod (osrednje Sredozemsko morje)

Abdelkarim DERBALI, Aymen HADJ TAIEB & Wassim KAMMOUN

The Current Status of *Polititapes aureus* (Mollusca: Bivalvia) in the Coastal Zone of Sfax, Tunisia (Central Mediterranean) 117
Trenutno stanje vrste Polititapes aureus (Mollusca: Bivalvia) na obalnem območju Sfaksa v Tuniziji (osrednje Sredozemlje)

Neža LEBAN & Valentina PITACCO

Current Knowledge on the Distribution of the Poorly Known Echiurid Species *Maxmuelleria gigas* (M. Müller, 1852) in the Slovenian Sea 125
Trenutno poznavanje prostorske razporeditve manj poznane vrste zvezdaša Maxmuelleria gigas (M. Müller, 1852) v slovenskem morju

Jan MALEJ, Tjaša KOGOVŠEK, Martin VODOPIVEC, Janja FRANCÉ, Patricija MOZETIČ, Matevž MALEJ & Alenka MALEJ

Long-Term Study of Zooplankton Biomass in the Gulf of Trieste (Adriatic Sea) 133
Dolgoročna študija zooplanktonske biomase v Tržaškem zalivu (Jadransko morje)

Sihem RAFRAFI-NOUIRA, Rimel BENMESSAOUD, Mourad CHÉRIFF, Christian REYNAUD & Christian CAPAPÉ

Occurrence of the Longjaw Snake Eel, *Ophisurus serpens* (Ophichthidae), in Tunisian Waters (Central Mediterranean Sea) 145
Pojavljanje zobate jegulje, Ophisurus serpens (Ophichthidae), iz tunizijskih voda (osrednje Sredozemsko morje)

FLORA
FLORA
FLORA

**Martina ORLANDO-BONACA, Artur BONACA,
Diego BONACA & Ana ROTTER**

Seagrasses: a Promising Source of
Bioactive Compounds for Human
Health Applications 153
*Morske cvetnice: obetaven vir
bioaktivnih spojin za uporabo
v zdravstvu*

OCENE IN POROČILA
RECENSIONI E RELAZIONI
REVIEWS AND REPORTS

Shin-ichi Uye

Book review: Mirrors of the Sea: When
Science and Art Meet. 30 Years of the
Unesco Intergovernmental Oceanographic
Commission in Slovenia 167
Kazalo k slikam na ovitku 169
Index to images on the cover 169

received: 2025-04-15

DOI 10.19233/ASHN.2025.19

SEAGRASSES: A PROMISING SOURCE OF BIOACTIVE COMPOUNDS FOR HUMAN HEALTH APPLICATIONS

Martina ORLANDO-BONACA

Marine Biology Station Piran, National Institute of Biology, SI-6330 Piran, Fornače 41, Slovenia
E-mail: martina.orlando@nib.si

Artur BONACA & Diego BONACA

Ulica Vena Piloni 5, SI-6000 Koper, Slovenia

Ana ROTTER

Marine Biology Station Piran, National Institute of Biology, SI-6330 Piran, Fornače 41, Slovenia

ABSTRACT

Seagrasses are unique marine flowering plants that provide critical ecological services and can serve as valuable reservoirs of bioactive compounds with potential health benefits. This review explores the bioactive metabolites found in four seagrass species native to European marine waters: Cymodocea nodosa, Posidonia oceanica, Zostera marina, and Nanozostera noltei. These species exhibit diverse chemical properties, including antioxidants, antimicrobials, and anti-inflammatory agents, making them promising candidates for pharmaceutical, nutraceutical, and cosmetic applications. Despite their promising applications in biotechnology, their full potential remains underexplored due to research and technological limitations. Future studies should focus on optimizing extraction methods, exploring synergistic interactions, and ensuring sustainable utilization of these valuable marine resources.

Key words: seagrasses, European marine waters, bioactive compounds, human health applications

FANEROGAME MARINE: UNA PROMETTENTE FONTE DI COMPOSTI BIOATTIVI PER APPLICAZIONI NEL CAMPO DELLA SALUTE UMANA

SINTESI

Le fanerogame sono piante marine che forniscono servizi ecologici critici e possono servire come preziosi serbatoi di composti bioattivi con potenziali benefici per la salute. Questa rassegna esplora i metaboliti bioattivi trovati in quattro specie di fanerogame native delle acque marine europee: Cymodocea nodosa, Posidonia oceanica, Zostera marina e Nanozostera noltei. Queste specie presentano diverse proprietà chimiche, tra cui antiossidanti, antimicrobici e agenti antinfiammatori, che le rendono promettenti candidati per applicazioni farmaceutiche, nutraceutiche e cosmetiche. Nonostante le loro promettenti applicazioni in biotecnologia, il loro pieno potenziale rimane poco esplorato a causa delle limitazioni tecnologiche. Gli studi futuri dovrebbero concentrarsi sull'ottimizzazione dei metodi di estrazione, sull'esplorazione delle interazioni sinergiche e sull'utilizzo sostenibile di queste preziose risorse marine.

Parole chiave: fanerogame, acque marine europee, composti bioattivi, applicazioni per la salute umana

INTRODUCTION

Seagrasses are the only group of marine flowering plants, uniquely adapted to thrive in underwater environments. Their morphological, physiological, ecological, and genetic adaptations facilitate survival in these habitats, including underwater pollination, internal gas transport, and the presence of epidermal chloroplasts (Hemminga & Duarte, 2000). Originating from terrestrial species, seagrasses have transitioned to marine habitats, forming vast meadows in shallow waters (Papenbrock, 2012). These meadows provide essential ecosystem services, serving as key feeding and breeding grounds for marine organisms (Ruiz-Frau *et al.*, 2017). Referred to as the “lungs of the sea,” seagrasses contribute significantly to oxygen production and carbon sequestration, mitigating climate change through the blue carbon storage (Papenbrock, 2012; Bedulli *et al.*, 2020; James *et al.*, 2023). Indeed, seagrass ecosystems are estimated to sequester approximately 140 megagrams of organic carbon (C_{org}) per hectare within the top meter of soil (Fourqurean *et al.*, 2012). This accumulation occurs over centennial to millennial time scales, making seagrass up to 40 times more effective at capturing organic carbon compared to terrestrial forest soils. (Serrano *et al.*, 2021). Additionally, they stabilize sediments, filter pollutants, and enhance water quality (Reynolds *et al.*, 2016; Bonanno & Orlando-Bonaca, 2017). As a result, seagrass meadows are recognized as one of the priority habitats in Europe within the EU Habitat Directive (HD, 92/43/EEC) and the evaluation of their status is crucial for the implementation of the EU Water Framework Directive (WFD), and the Marine Strategy Framework Directive (MSFD) (Orlando-Bonaca *et al.*, 2015). Moreover, they are among the least conserved marine ecosystems (Hu *et al.*, 2021).

Beyond their ecological significance, seagrasses are gaining attention for their rich diversity of bioactive compound with antioxidant, anti-inflammatory, cytotoxic and antimicrobial activities (Ameen *et al.*, 2024). Despite increasing interest in marine-derived bioactives, research on seagrasses remains limited, presenting an opportunity for further exploration (Ribas-Taberner *et al.*, 2025). In recent years, there has been growing interest in seagrass-derived bioactive compounds for biotechnological applications (Rotter *et al.*, 2021, 2023). However, challenges such as preservation of their ecological stability and technological limitations hinder their full integration into biotechnology (Rotter *et al.*, 2021; Gono *et al.*, 2022).

This review provides an in-depth examination of the bioactive compounds of biotechnological relevance in the biomedical field, isolated and characterized from four native seagrass species in

European marine waters: *Cymodocea nodosa* (Ucria) Ascherson, *Posidonia oceanica* (Linnaeus) Delile, *Zostera marina* Linnaeus, and *Nanozostera noltei* (Hornemann) Tomlinson & Posluszny. The paper also highlights some areas for future research on these seagrass species that could improve their use for human health while maintaining sustainable practices.

MATERIAL AND METHODS

A systematic literature review was conducted to gain insights into the breadth of studies published on the bioactive compounds of the four European seagrass species and their potential benefits. The literature search, covering 2004–2025, was performed in Scopus Advanced search, VOSviewer and Google Scholar, focusing on peer-reviewed articles and excluding grey literature. The keywords employed in the search included seagrass, *Cymodocea nodosa*, *Posidonia oceanica*, *Zostera marina* and *Nanozostera noltei*, biotechnology, bioactive compounds, bioactive molecules, bioactive potential, health, pharma and biomedicine.

RESULTS AND DISCUSSION

Bioactive compounds in *Cymodocea nodosa*

Cymodocea nodosa (commonly known as Lesser Neptune grass) is the most common species in shallow, sheltered to semi-exposed Mediterranean sites, forming meadows that can be either mono-specific or mixed with *N. noltei* (Buia *et al.*, 1985). This species has its origins in tropical regions and is currently confined to the Mediterranean Sea and some areas in the North Atlantic, extending from southern Portugal and Spain to Senegal, including the Canary Islands and Madeira (OSPAR, 2010). It can be found in coastal waters, coastal lagoons, inlets and estuaries and other sheltered and semi-exposed habitats (Ivajnsiĉ *et al.*, 2022). It is recognized as a pioneer species, adept at rapidly colonizing bare patches of the sea floor, since its rhizome can grow several meters annually (Boudouresque *et al.*, 1994).

C. nodosa contains several bioactive metabolites with potential pharmaceutical applications (Tab. 1; Fig. 1). Among them, diarylheptanoids and terpenoids have demonstrated antibacterial activity, particularly against mycobacterial strains and methicillin-resistant strains of *Staphylococcus aureus* (MRSA) (Kontiza *et al.*, 2008). The relatively simple structures of these metabolites also suggest they could be targets for further chemical modification to enhance their antibacterial properties (Kontiza *et al.*, 2008). In addition, diarylheptanoids, especially cymodienol, have shown strong cytotoxicity (Kontiza *et al.*, 2005).

Tab. 1: Bioactive compounds extracted from the four European marine seagrasses and their reported bioactivities. CN = *Cymodocea nodosa*; ZM = *Zostera marina*; PO = *Posidonia oceanica*; NN = *Nanozostera noltei*.

Tab. 1: Bioaktivne spojine, pridobljene iz štirih evropskih morskih cvetnic, in njihove potrjene bioaktivnosti. CN = *Cymodocea nodosa*; ZM = *Zostera marina*; PO = *Posidonia oceanica*; NN = *Nanozostera noltei*.

Compound	Bioactivity	Seagrass species	References
Terpenoids	Antibacterial	CN	Kontiza <i>et al.</i> (2008)
Diarylheptanoids	Antibacterial, anticancer/ cytotoxic	CN, ZM	Kontiza <i>et al.</i> (2005); Kontiza <i>et al.</i> (2008); Li <i>et al.</i> (2021); Cacciola <i>et al.</i> (2024)
Polysaccharides	Antioxidant, antimicrobial, cytotoxic, antihypertensive, anti- obesity, antidiabetic	CN, PO, ZM	Kolenchenko <i>et al.</i> (2005); Berfad & Alnour (2014); Kolsi <i>et al.</i> (2015, 2016, 2017a, 2017b, 2018); Benito-Gonzalez <i>et al.</i> (2019); Perumal <i>et al.</i> (2021); Ribas-Taberner <i>et al.</i> (2025)
Phenolics	Antidiabetic, anti-inflammatory, antioxidant, analgesic, antihypertensive, antirheumatic, anti-arthritic, antimicrobial, anticancer/ cytotoxic	CN, PO, ZM, NN	Rees <i>et al.</i> (2008); Choi <i>et al.</i> , (2009); Subhashini <i>et al.</i> (2013); Berfad & Alnour (2014); Barletta <i>et al.</i> (2015); Custódio <i>et al.</i> (2015); Grignon-Dubois <i>et al.</i> (2015); Kolsi <i>et al.</i> (2017b); Manck <i>et al.</i> (2017); Styshova <i>et al.</i> (2017); Farid <i>et al.</i> (2018); Leri <i>et al.</i> (2018); Benito-Gonzalez <i>et al.</i> (2019); Sevimli-Gur & Yesil-Celiktas (2019); Vasarri <i>et al.</i> (2020); Ammar <i>et al.</i> (2021); Mechchate <i>et al.</i> (2021); Vasarri <i>et al.</i> (2021); Razgonova <i>et al.</i> (2022); Abruscato <i>et al.</i> (2023); Grignon- Dubois & Rezzonico (2023); Vasarri <i>et al.</i> (2023); Chabbani <i>et al.</i> (2024); Kevrekidou <i>et al.</i> (2024); Micheli <i>et al.</i> (2024); De Leo <i>et al.</i> (2025)
Proteins and peptides	Antioxidant, antimicrobial	PO	Berfad & Alnour (2014); Benito-Gonzalez <i>et al.</i> (2019); Punginelli <i>et al.</i> (2023)

Furthermore, sulfated polysaccharides derived from *C. nodosa* exhibit a diverse range of bioactive properties, including significant antioxidant, antimicrobial, cytotoxic, antihypertensive activity, anti-obesity and anti-diabetic properties (Kolsi *et al.*, 2015, 2016, 2017a; Perumal *et al.*, 2021; Ribas-Taberner *et al.*, 2025; see Tab. 1). Sulfated polysaccharides derived from *C. nodosa* warrant special attention also for treating male infertility and enhancing sperm quality due to their antioxidant capabilities (Kolsi *et al.*, 2018; Ribas-Taberner *et al.*, 2025).

Among phenolic compounds, the chicoric acid, that was found to be abundant in *C. nodosa*, has high potential for management of metabolic alterations, thus representing a target for treatment of diabetes

and its complications (De Leo *et al.*, 2025). The hydroalcoholic extract of *C. nodosa* containing phenolic compounds exhibits significant antioxidant activity, which may protect against oxidative stress and be applied for managing metabolic disorders, particularly diabetes, by enhancing insulin secretion and reducing hyperglycemia and lipid levels in the blood (Kolsi *et al.*, 2017b). More recently, Chabbani *et al.* (2024) identified additional phenolic compounds in *C. nodosa*, including sinapic acid, myricetin, and quercetin-3-O-rutinoside, which demonstrate potential therapeutic properties, aimed at combating oxidative stress and inflammation. The extract's safety profile, with no cytotoxic effects observed, further supports its application in skincare products (Chabbani *et al.*, 2024).

Bioactive compounds in *Posidonia oceanica*

Posidonia oceanica (the Neptune grass) is the most important Mediterranean endemic seagrass species, that forms meadows extending from the surface to depths of 40–45 m (Telesca *et al.*, 2015). These meadows provide multiple ecosystem services, including protection of coastlines from erosion, provision of breeding and nesting habitats for economically important species, and serving as a vital source of oxygen while facilitating high carbon sequestration within the matte (consisting of intertwined remnants of roots, rhizomes, and leaves, which effectively trap sediment and organic carbon, creating stable substrates) (Pergent-Martini *et al.*, 2021). The widespread distribution of *P. oceanica* in the Mediterranean Sea suggests that these meadows are the product of ecological and evolutionary processes that have unfolded over centuries (Giakoumi *et al.*, 2013; Telesca *et al.*, 2015).

Extracts of *P. oceanica* have shown anti-diabetic, antioxidant, vasoprotective and antimicrobial activities (Gokce & Haznedaroglu, 2008; Ozbil *et al.*, 2024). Phenolic compounds, such as chicoric, gentisic, ferulic and caftaric acid have potential analgesic, anti-inflammatory, antirheumatic, anti-arthritis and anticarcinogenic activities (Grignon-Dubois & Rezzonico, 2015; Mechchate *et al.*, 2021; Micheli *et al.*, 2024), see Tab. 1. These were also observed in *C. nodosa* and such phenolic similarities between *P. oceanica* and *C. nodosa* imply a possible evolutionary connection that may suggest that these families possess adaptive traits developed in response to comparable environmental pressures (Grignon-Dubois & Rezzonico, 2015). Polyphenols including gallic acid, catechin, epicatechin and chlorogenic acid that were isolated in *P. oceanica* extracts were able to inhibit the expression and activity of gelatinases MMP-2 and MMP-9 (enzymes that break down the extracellular matrix), which are associated with cancer cell invasion and metastasis (Barletta *et al.*, 2015). Additionally, Leri *et al.* (2018) highlighted the significant antioxidant properties of these bioactive metabolites, with potential use in therapeutic applications against malignancies and other chronic conditions due to their ability to impair malignant cell migration through autophagy modulation. Furthermore, the (hydro)alcoholic extracts of *P. oceanica* leaves containing phenols has potential anticancer applications, given their antioxidant, anti-inflammatory and anti-glycation properties (Farid *et al.*, 2018; Sevimli-Gur & Yesil-Celiktas, 2019; Abrucato *et al.*, 2023; Vassari *et al.*, 2023; Kevrekidou *et al.*, 2024).

Cornara *et al.* (2018) reported that the extract of *P. oceanica*, rich in CA, exhibits significant radical scavenging activity and lipolytic activity. Regarding skin protection, the polyphenols of *P. oceanica* have potential therapeutic applications in treating conditions like psoriatic skin inflammation, indicating its potential use in complementary medicine for inflammatory skin diseases (Micheli *et al.*, 2024). Messina *et al.* (2021) concluded that the phenolic compounds found in *P. oceanica* are also promising candidates for applications particularly in cosmeceuticals and as protective agents against oxidative stress and UV-induced damage. They highlighted that the dried leaves of *P. oceanica* demonstrated significantly higher antioxidant activity than fresh leaves.

P. oceanica butanol extract containing phenolic acids and flavonoids, encapsulated in gelatin nanoparticles, as a promising and effective antidiabetic therapy (Ammar *et al.*, 2021). Additionally, the extract from *P. oceanica* egagropiles (round-shaped conglomerations created by the progressive disintegration of fibrous material sourced from its foliage and transported to nearby coastal areas by waves) showed a moderate inhibition against H5N1 virus (Farid *et al.*, 2018).

P. oceanica extracts containing phenols and flavonoids exhibit anti-inflammatory activities (Vasari *et al.*, 2020), while those containing polyphenols, polysaccharides and proteins exhibit antioxidant and antimicrobial activities (Berfad & Alnour, 2014; Benito-Gonzalez *et al.*, 2019). The main polysaccharide known from angiosperm land plants in the cell walls is cellulose; however, it is currently lacking biotechnologically relevant biological activity properties for biomedical purposes (Pfeifer & Classen, 2020). The polysaccharides identified in *P. oceanica* that have bioactive qualities include galacturonic acid and xylose, which are components of pectin and hemicellulose. These polysaccharides contribute to the antioxidant, antifungal, and antiviral properties of the extracts, making them potential candidates for use in food and pharmaceutical applications (Benito-Gonzalez *et al.*, 2019).

Bioactive compounds in *Zostera marina*

Zostera marina (commonly known as Eelgrass) is one of the world's most widespread marine phanerogams, predominantly found along the temperate and cold coasts of the Northern Hemisphere, but it is also present in the Mediterranean Sea, where its distribution is more limited. The species fulfills important roles as an ecosystem engineer in estuarine and coastal waters, enhancing biodiversity, contributing to sediment stability, and carbon and nutrient




Fig. 1: Potential bioactive properties of the four European marine seagrasses (photos: Tihomir Makovec).
Sl. 1: Potencialne bioaktivne lastnosti štirih evropskih morskih cvetnic (slike: Tihomir Makovec).

storage (Duarte *et al.*, 2013; Blok *et al.*, 2018). Distinct life history strategies and a wide tolerance to a range of salinity and temperature conditions allow this seagrass species to exploit areas from brackish estuaries to open oceans over a large geographical range (Nejrup & Pedersen 2008).

Z. marina leaves decay very slowly and indeed, extracts from this seagrass exhibited antimicrobial and antioxidant activities (Zheng *et al.*, 2014). Polysaccharides from *Z. marina* exhibit significant antioxidant activities (see Tab. 1). An example is zosterin, a low-etherified pectin with potential in reducing lipid peroxidation products, suggesting its use in therapeutic applications for oxidative stress-related conditions (Kolenchenko *et al.*, 2005). Additionally, zosterin has been associated with various pharmacological effects, including hypocholesterolemic, antibacterial, antiviral, and immunomodulating properties, indicating its broad potential in medical applications (Kolenchenko *et al.*, 2005).

Phenolics, such as zosteric acid and others, have shown potential as antimicrobial agents (Rees *et al.*, 2008; Choi *et al.*, 2009). Rosmarinic

acid (RA) from *Z. marina* was also recognized for its antioxidant properties (Custódio *et al.*, 2016). Rosmarinic acid and flavonoids extracted from *Z. marina* have also evidenced therapeutic potential in the treatment of type 2 diabetes and hyperlipidemia, due to their antioxidant, anti-inflammatory, and immunomodulatory activities (Styshova *et al.*, 2017). Several polyphenols from *Z. marina* (chicoric, p-coumaric, rosmarinic, benzoic, ferulic and caffeic acids) exhibited significant cytotoxic effects against various cancer cell lines, indicating their potential use as chemotherapeutic agents in cancer treatment and as preventive supplements in oncology (Sevimli-Gur & Yesil-Celiktas, 2019). *Z. marina* polyphenols also have potential applications in pharmaceutical industry due to their antioxidant and anti-inflammatory properties (Grignon-Dubois & Rezzonico, 2023; Razgonova *et al.*, 2022).

Diarylheptanoids extracted from *Z. marina* exhibited selective cytotoxic effects and significant inhibition of the HCT116 colon cancer cell line and potential for reducing tumor growth *in vivo* (Li *et al.*, 2021; Cacciola *et al.*, 2024).

Bioactive compounds in *Nanozostera noltei*

Nanozostera noltei (the dwarf eelgrass) is widely distributed along the northeastern Atlantic coasts, from Norway to Mauritania, and in the Mediterranean and Black Seas (Short *et al.*, 2007). This species frequently thrives in meadows of *C. nodosa* in the infralittoral belt, but it can be found in a variety of habitat types with variable salinity. In lagoonal systems, it establishes a narrow band within the mediolittoral zone, demonstrating its ability to endure partial desiccation (Lipej *et al.*, 2006).

Only in recent decades several studies have examined the phenolic profile of this species (see Tab. 1), with rosmarinic, caffeic and zosteric acids having antioxidant properties (Grignon-Dubois *et al.*, 2012; Subhashini *et al.*, 2013). Phenolic compounds, flavons and flavonoids from *N. noltei* suggest potential applications in pharmaceuticals (Manck *et al.*, 2017).

Extracts of *N. noltei* are notable for their ability to chelate both copper and iron ions, potentially aiding in the reduction of oxidative stress and associated conditions (Custódio *et al.*, 2016). This seagrass has demonstrated significant toxicity to various cell lines, suggesting promising applications in cancer research, particularly in targeting hepatocarcinoma cells, and may also contribute to alleviating Alzheimer's disease (Custódio *et al.*, 2016).

Sourcing and extraction of biotechnologically relevant bioactive compounds

Despite the ample evidence that indicates seagrasses as a resource for novel health and pharmaceutical formulations, it is important to highlight that seagrasses provide an excellent substratum for epiphytic organisms (such as bacteria, fungi, protozoa and algae), making these organisms an integral component of seagrass ecosystems (Borowitzka *et al.*, 2006). In addition, distinct microbial communities within the seagrass leaves, roots, flowers, fruits, seeds and the rhizosphere are likely involved in ecologically important processes that benefit plants (Tarquinio *et al.*, 2021). These microbial communities may be promoting the production or might be the main source of biotechnologically relevant secondary metabolites (Panno *et al.*, 2013; Rotter *et al.*, 2021). When bioprospecting for novel compounds, it is therefore important to sterilize the seagrass surfaces (targeting epiphytes) and conduct molecular analyses to assess the presence of microbial genetic markers. This can identify the producer organism of the biotechnologically relevant compound.

Regardless of the source of the main biological activities, it is important to again highlight that seagrass ecosystems are now recognized as critical habitats supporting biodiversity and ecosystem services (Duarte *et al.*, 2025). Hence, when confirming the bioactivity and target sector for compounds from seagrasses, sustainable sourcing of the biomass and supply of biomolecules must be considered to maintain the ecological stability of these fragile communities (Rotter *et al.*, 2020, 2021). In the case of seagrasses, research can rely on harvesting or beach wrack, provided that their harvesting is sustainable and does not affect the ecosystem balance in coastal areas (Rudovica *et al.*, 2021). Wherever possible, harvesting can also be substituted/complemented by modern biotechnology approaches including systems biology and metabolic engineering (Rotter *et al.*, 2024). Finally, to contribute to environmental sustainability to an even higher degree, besides sourcing, extraction methods should be improved by means of greener methodologies avoiding the use of toxic organic solvents (Benito-González *et al.*, 2019).

CONCLUSIONS

Seagrass meadows are among the most essential and productive yet threatened benthic habitats, warranting significant scientific attention, particularly in the field of biotechnology. The bioactive metabolites found in marine plants have been shown to address various human health concerns, including several diseases, and hold promise for developing new natural-based therapeutic products.

The bioactive compounds found in the reviewed seagrass species exhibit a diverse array of health-promoting properties, positioning them as significant resources for the development of novel therapeutic drugs. Their bioactive potential highlights their ability to act as inhibitors of various harmful pathogens within the pharmaceutical sector. These compounds also show promise in treating metabolic disorders and cancers. Nevertheless, this review does not cover the literature on human clinical trials involving bioactive molecules derived from seagrasses or the sustainable production of drugs, considering the yield, maintenance of ecological balance or the use of green extraction processes, as this represents a separate and extensive area of research. Many bioactive metabolites from seagrasses remain unidentified, and ongoing research aims to uncover all relevant compounds that contribute to human well-being. Future research should focus on environmentally and economically sustainable extraction methods for metabolites with significant therapeutic potential.

Further studies should investigate the long-term stability and shelf life of seagrass extracts and their use in various formulations. Additionally, future research should explore the potential synergistic effects of various seagrass bioactive compounds, both among themselves and in combination with other ingredients, to enhance product effectiveness. Finally, as the extraction of bioactive compounds

from marine resources continues to expand globally, it is crucial to assess the environmental impact of harvesting seagrass meadows. This will ensure that sustainable practices are upheld while utilizing this valuable resource. Hence, extraction from seagrasses or their epiphytes should be complemented by using metabolic engineering and chemical synthesis.

MORSKE CVETNICE: OBETAVEN VIR BIOAKTIVNIH SPOJIN ZA UPORABO V ZDRAVSTVU

Martina ORLANDO-BONACA

Morska biološka postaja Piran, Nacionalni inštitut za biologijo, SI-6330 Piran, Fornače 41, Slovenija
E-mail: martina.orlando@nib.si

Artur BONACA & Diego BONACA

Ulica Vena Piloni 5, SI-6000 Koper, Slovenija

Ana ROTTER

Morska biološka postaja Piran, Nacionalni inštitut za biologijo, SI-6330 Piran, Fornače 41, Slovenija

POVZETEK

Morske cvetnice so edinstvene morske rastline, ki zagotavljajo ključne ekološke storitve in lahko služijo kot dragoceni viri bioaktivnih spojin s potencialnimi koristmi za zdravje ljudi. Ta pregled obravnava bioaktivne metabolite, prisotne v štirih vrstah morskih trav, ki so domorodne v evropskih morskih vodah: Cymodocea nodosa, Posidonia oceanica, Zostera marina in Nanozostera noltei. Te vrste izkazujejo raznolike kemijske lastnosti, vključno z antioksidativnim, protimikrobnim in protivnetnim delovanjem, zaradi česar so obetavne za uporabo v farmacevtski, nutracevtski in kozmetični industriji. Kljub njihovem potencialu v biotehnologiji pa ta ostaja premalo raziskana zaradi raziskovalnih in tehnoloških omejitev. Prihodnje študije bi se morale osredotočiti na optimizacijo metod ekstrakcije, raziskovanje sinergijskih učinkov ter zagotavljanje trajnostne uporabe teh dragocenih morskih virov.

Ključne besede: morske trave, evropske morske vode, bioaktivne spojine, uporaba za zdravje ljudi

REFERENCES

- Abruscato, G., R. Chiarelli, V. Lazzara, D. Punginelli, S. Sugár, M. Mauro, M. Librizzi, V. Di Stefano, V. Arizza, A. Vizzini, M. Vazzana & C. Luparello (2023): In Vitro Cytotoxic Effect of Aqueous Extracts from Leaves and Rhizomes of the Seagrass *Posidonia oceanica* (L.) Delile on HepG2 Liver Cancer Cells: Focus on Autophagy and Apoptosis. *Biology*, 12, 616. <https://doi.org/10.3390/biology12040616>.
- Ameen, H.M., A. Jayadev, G. Prasad & D.I. Nair (2024): Seagrass Meadows: Prospective Candidates for Bioactive Molecules. *Molecules*, 29, 4596. <https://doi.org/10.3390/molecules29194596>.
- Ammar, N.M., H.A. Hassan, M.A. Mohammed, A. Serag, S.H. Abd El-Alim, H. Elmotasem, M. El Raey, A.N. El Gendy, M. Sobeh & A.Z. Abdel-Hamid (2021): Metabolomic profiling to reveal the therapeutic potency of *Posidonia oceanica* nanoparticles in diabetic rats. *RSC Adv.*, 11, 8398-8410.
- Barletta, E., M. Ramazzotti, F. Fratianni, D. Pessani & D. Degl'Innocenti (2015): Hydrophilic extract from *Posidonia oceanica* inhibits activity and expression of gelatinases and prevents HT1080 human fibrosarcoma cell line invasion. *Cell Adhes. Migr.*, 9, 422-431.
- Bedulli, C., P.S. Lavery, M. Harvey, C.M. Duarte & O. Serrano (2020): Contribution of Seagrass Blue Carbon Toward Carbon Neutral Policies in a Touristic and Environmentally-Friendly Island. *Front. Mar. Sci.*, 7, 1. doi: 10.3389/fmars.2020.00001.
- Benito-Gonzalez, I., A. Lopez-Rubio, A. Martinez-Abad, A.R. Ballester, I. Falco, L. Gonzalez-Candelas, G. Sanchez, J. Lozano-Sanchez, I. Borrás-Linares, A. Segura-Carretero & M. Martínez-Sanz (2019): In-Depth Characterization of Bioactive Extracts from *Posidonia oceanica* Waste Biomass. *Mar. Drugs*, 17, 409.
- Berfad, M. & T. Alnour (2014): Phytochemical analysis and Antibacterial activity of the 5 different extract from the seagrasses *Posidonia oceanica*. *J. Med. Plants Stud.*, 4, 15-18.
- Blok, S.E., B. Olesen & D. Krause-Jensen (2018): Life history events of eelgrass *Zostera marina* L. populations across gradients of latitude and temperature. *Mar. Ecol. Prog. Ser.*, 590, 79-93. <https://doi.org/10.3354/meps12479>.
- Bonanno, G. & M. Orlando-Bonaca (2017): Trace elements in Mediterranean seagrasses: accumulation, tolerance and biomonitoring. A review. *Marine Pollution Bulletin*, 125, 8-18.
- Borowitzka, M.A., P.S. Lavery & M. van Keulen (2006): Epiphytes of seagrasses. In: *Seagrasses: Biology, Ecology and Conservation*. Larkum, A.W.D., R.J. Orth & C.M. Duarte (Eds.); pp.441-462, Springer. DOI: https://doi.org/10.1007/978-1-4020-2983-7_19.
- Boudouresque, C.F., A. Meinesz, M. Ledoyer & P. Vitiello (1994). Les herbiers à phanérogames marines. In: Bellan-Santini, D., Lacaze, J., Poizat, C. (Eds.), *Les biocénoses marines et littorales de Méditerranée, synthèse, menaces et perspectives*. Muséum National d'Histoire Naturelle publ, pp. 98-118.
- Buía, M., G. Russo & L. Mazzella (1985): Interrelazioni Tra *Cymodocea nodosa* (Ucria) Aschers. e *Zostera noltii* Hornem. in Un Prato Misto Superficiale Dell'isola d'Ischia. *Nova Thalass.*, 7, 406-408.
- Cacciola, N.A., P. De Cicco, R. Amico, F. Sepe, Y. Li, L. Grauso, M.F. Nan., S. Scarpato, C. Zidorn, A. Mangoni & F. Borrelli (2024): Zosterabisphephenone B, a new diarylheptanoid heterodimer from the seagrass *Zostera marina*, induces apoptosis cell death in colon cancer cells and reduces tumour growth in mice. *Phytother. Res.*, 38, 4168-4176.
- Choi, H.-G., J.-H. Lee, H.-H. Park & F. Sayegh (2009). Antioxidant and Antimicrobial Activity of *Zostera marina* L. Extract. *Algae*, 24, 179-184.
- Custódio, L., S. Laukaityte, A.H. Engelen, M.J. Rodrigues, H. Pereira, C. Vizetto-Duarte, L. Barreira, H. Rodriguez, F. Albericio & J. Varela (2016): A comparative evaluation of biological activities and bioactive compounds of the seagrasses *Zostera marina* and *Zostera noltei* from southern Portugal. *Nat. Prod. Res.*, 30, 724-728.
- De Leo, M., L. Ciccone, V. Menicagli, E. Balestri, A. Braca, P. Nieri, L. Testai (2025): *Cymodocea nodosa*, a Promising Seagrass of Nutraceutical Interest: Overview of Phytochemical Constituents and Potential Therapeutic Uses. *Nutrients*, 17, 1236. <https://doi.org/10.3390/nu17071236>.
- Duarte, C.M., I.J. Losada, I.E. Hendriks, I. Mazarrasa & N. Marbà (2013): The role of coastal plant communities for climate change mitigation and adaptation. *Nat. Clim. Chang.*, 3, 961-968.
- Duarte, C.M., E.T. Apostolaki, O. Serrano, A. Steckbauer & R.K.F. Unsworth (2025): Conserving seagrass ecosystems to meet global biodiversity and climate goals. *Nat. Rev. Biodivers.*, 1, 150-165. <https://doi.org/10.1038/s44358-025-00028-x>.
- Farid, M., M. Marzouk, S. Hussein, A. Elkhateeb & E.-S. Saleh (2018): Comparative study of *Posidonia oceanica* L.: LC/ESI/MS analysis, cytotoxic activity and chemosystematic significance. *J. Mater. Environ. Sci.*, 9, 1676-1682.
- Fourqurean, J.W., C.M. Duarte, H. Kennedy, N. Marbà, M. Holmer, M.A. Mateo, E.T. Apostolaki, G.A. Kendrick, D. Krause-Jensen, K.J. McGlathery & O. Serrano (2012): Seagrass ecosystems as a globally significant carbon stock. *Nat. Geosci.*, 5, 505-509. <https://doi.org/10.1038/ngeo1477>.

- Giakoumi, S., M. Sini, V. Gerovasileiou, T. Mazor, J. Beher, H.P. Possingham, A. Abdulla, M.E. Cinar, P. Dendrinis, A.C. Gucu, A.A. Karamanlidis, P. Rodic, P. Panayotidis, E. Taskin, A. Jaklin, E. Voultsiadou, C. Webster, A. Zenetos & S. Katsanevakis (2013): Ecoregion-based conservation planning in the mediterranean: dealing with large-scale heterogeneity. *PloS One*, 8(10), e76449. 10.1371/journal.pone.0076449.
- Gokce, G. & M.Z. Haznedaroglu (2008): Evaluation of antidiabetic, antioxidant and vasoprotective effects of *Posidonia oceanica* extract. *Journal of Ethnopharmacology*, 115(1), 122-130. <https://doi.org/10.1016/j.jep.2007.09.016>.
- Gono, C.M.P., P. Ahmadi, T. Hertiani, E. Septiana, M.Y. Putra & G.A. Chianese (2022): Comprehensive Update on the Bioactive Compounds from Seagrasses. *Mar. Drugs*, 20, 406. <https://doi.org/10.3390/md20070406>.
- Grignon-Dubois, M., B. Rezzonico & T. Alcoverro (2012): Regional scale patterns in seagrass defences: Phenolic acid content in *Zostera noltii*. *Estuarine, Coastal and Shelf Science*, 114, 18-22. <https://doi.org/10.1016/j.ecss.2011.09.010>.
- Grignon-Dubois, M. & B. Rezzonico (2015): Phenolic fingerprint of the seagrass *Posidonia oceanica* from four locations in the Mediterranean Sea: First evidence for the large predominance of chicoric acid. *Bot. Mar.*, 58, 379-391.
- Grignon-Dubois, M. & B. Rezzonico (2023): Phenolic chemistry of the seagrass *Zostera marina* Linnaeus: First assessment of geographic variability among populations on a broad spatial scale. *Phytochemistry*, 213, 113788.
- Hemminga, M.A. & C.M. Duarte (2000): *Seagrass ecology*. Cambridge University Press. <https://doi.org/10.1017/CBO9780511525551>.
- Hu, W., D. Zhang, B. Chen, X. Liu, X. Ye, Q. Jiang, X. Zheng, J. Du & S. Chen (2021): Mapping the seagrass conservation and restoration priorities: Coupling habitat suitability and anthropogenic pressures. *Ecological Indicators*, 129, 107960. <https://doi.org/10.1016/j.ecolind.2021.107960>.
- Ivajnski, D., M. Orlando-Bonaca, D. Donša, V.J. Grujić, D. Trkov, B. Mavrič & L. Lipej (2022): Evaluating Seagrass Meadow Dynamics by Integrating Field-Based and Remote Sensing Techniques. *Plants*, 11, 1196. <https://doi.org/10.3390/plants11091196>.
- James, R.K., L.M. Keyzer, S.J. van de Velde, P.M.J. Herman, M.M. van Katwijk, & T.J. Bouma (2023): Climate Change Mitigation by Coral Reefs and Seagrass Beds at Risk: How Global Change Compromises Coastal Ecosystem Services. *Sci. Total Environ.*, 857(3), 159576. <https://doi.org/10.1016/j.scitotenv.2022.159576>.
- Kevrekidou, A., A.N. Assimopoulou, V. Trachana, D. Stagos & P. Malea (2024): Antioxidant Activity, Inhibition of Intestinal Cancer Cell Growth and Polyphenolic Compounds of the Seagrass *Posidonia oceanica*'s Extracts from Living Plants and Beach Casts. *Mar. Drugs*, 22, 130.
- Kolenchenko, E.A., L.N. Sonina & Y.S. Khotimchenko (2005): Comparative in vitro Assessment of Antioxidant Activities of Low-Etherified Pectin from the Eelgrass *Zostera marina* and Antioxidative Medicines. *Russ. J. Mar. Biol.*, 31, 331-334.
- Kolsi, R.B.A., A. Ben Gara, R. Chaaben, A. El Feki, F.P. Patti, L. El Feki & K. Belghith (2015): Anti-obesity and lipid lowering effects of *Cymodocea nodosa* sulphated polysaccharide on high cholesterol-fed-rats. *Arch. Physiol. Biochem.*, 121, 210-217.
- Kolsi, R.B.A., J. Fakhfakh, F. Krichen, I. Jribi, A. Chiarore, F.P. Patti, C. Blecker, N. Allouche, H. Belghith & K. Belghith (2016): Structural characterization and functional properties of antihypertensive *Cymodocea nodosa* sulfated polysaccharide. *Carbohydr. Polym.*, 151, 511-522.
- Kolsi, R.B.A., B. Gargouri, S. Sassi, D. Frikha, S. Lassoued & K. Belghith (2017a): In vitro biological properties and health benefits of a novel sulfated polysaccharide isolated from *Cymodocea nodosa*. *Lipids Health Dis.*, 16, 252.
- Kolsi, R.B.A., H. Ben Salah, N. Jardak, R. Chaaben, A. El Feki, T. Rebai, K. Jamoussi, N. Allouche, H. Belghith & K. Belghith (2017b): Effects of *Cymodocea nodosa* extract on metabolic disorders and oxidative stress in alloxan-diabetic rats. *Biomed. Pharmacother.*, 89, 257-267.
- Kolsi, R., F.B. Abdallah, M. Elleuch, B. Kolsi, A.E. Feki & K.S. Belghith (2018): Antioxidant and fertility potentials of a marine sulfated polysaccharide. *JIM Sfax*, 28, 36-44.
- Kontiza, I., C. Vagias, J. Jakupovic, D. Moreau, C. Roussakis & V. Roussis (2005): Cymodienol and cymodiene: new cytotoxic diarylheptanoids from the sea grass *Cymodocea nodosa*. *Tetrahedron Letters*, 46(16), 2845-2847. doi: 10.1016/j.tetlet.2005.02.123.
- Kontiza, I., M. Stavri, M. Zloh, C. Vagias, S. Gibbons & V. Roussis (2008): New metabolites with antibacterial activity from the marine angiosperm *Cymodocea nodosa*. *Tetrahedron*, 64, 1696-1702.
- Leri, M., M. Ramazzotti, M. Vasarri, S. Peri, E. Barletta, C. Pretti & D. Degl'Innocenti (2018): Bioactive Compounds from *Posidonia oceanica* (L.) Delile Impair Malignant Cell Migration through Autophagy Modulation. *Mar. Drugs*, 16, 137.
- Li, Y., L. Grauso, S. Scarpato, C.N.A. Cacciola, F. Borrelli, C. Zidorn & A. Mangoni (2021): Stable Catechol Keto Tautomers in Cytotoxic Heterodimeric Cyclic Diarylheptanoids from the Seagrass *Zostera marina*. *Org. Lett.*, 23, 7134-7138.
- Lipej, L., R. Turk & T. Makovec (2006): Ogrožene vrste in habitatni tipi v slovenskem morju = Endangered Species and Habitat Types in the Slovenian Sea. *Zavod RS za varstvo narave, Ljubljana*, 264 pp.

- Manck, L., E. Quintana, R. Suárez, F.G. Brunb, I. Hernández, M.J. Ortega & E. Zubia (2017):** Profiling of Phenolic Natural Products in the Seagrass *Zostera noltei* by UPLC-MS. *Nat Prod Commun.*, 12(5), 687-690.
- Mechchate, H., I. Es-Safi, O. Mohamed Al Kamaly & D. Boustia (2021):** Insight into Gentisic Acid Antidiabetic Potential Using In Vitro and In Silico Approaches. *Molecules*, 26(7), 1932. doi: 10.3390/molecules26071932.
- Messina, C.M., R. Arena, S. Manuguerra, Y. Pericot, E. Curcuraci, F. Kerninon, G. Renda, C. Hellio & A. Santulli (2021):** Antioxidant Bioactivity of Extracts from Beach Cast Leaves of *Posidonia oceanica* (L.) Delile. *Mar. Drugs*, 19, 560.
- Micheli, L., M. Vassarri, D. Degl'Innocenti, L. Di Cesare Mannelli, C. Ghelardini, A. Emiliano, A. Verdelli, M. Caproni & E. Barletta (2024):** *Posidonia oceanica* (L.) Delile Is a Promising Marine Source Able to Alleviate Imiquimod-Induced Psoriatic Skin Inflammation. *Mar. Drugs*, 22, 300.
- Nejrup, L.B. & M.F. Pedersen (2008):** Effects of salinity and water temperature on the ecological performance of *Zostera marina*. *Aquat Bot*, 88, 239-246.
- Orlando-Bonaca, M., J. Francé, B. Mavrič, M. Grego, L. Lipej, V. Flander-Putrlje, M. Šiško & A. Falace (2015):** A new index (MediSkew) for the assessment of the *Cymodocea nodosa* (Ucria) Ascherson meadow's status. *Mar. Environ. Res.*, 110, 132-141.
- OSPAR Commission (2010):** Background Document for *Cymodocea* Meadows. OSPAR Commission, London, 30 pp., 487/2010.
- Ozbil, E., M. Ilktac, S. Ogmen, O. Isbilen, J.M. Duran Ramirez, J. Gomez, J.N. Walker, E. Volkan (2024):** In vitro antibacterial, antibiofilm activities, and phytochemical properties of *Posidonia oceanica* (L.) Delile: An endemic Mediterranean seagrass. *Heliyon*, 10(15), e35592.
- Panno, L., M. Bruno, S. Voyron, A. Anastasi, G. Gnani, L. Miserere & G.C. Varese (2013):** Diversity, ecological role and potential biotechnological applications of marine fungi associated to the seagrass *Posidonia oceanica*. *New Biotechnology*, 30(6), 685-694. <https://doi.org/10.1016/j.nbt.2013.01.010>.
- Papenbrock, J. (2012):** Highlights in Seagrasses' Phylogeny, Physiology, and Metabolism: What Makes Them Special? International Scholarly Research Network, ISRN Botany, Article ID 103892, doi:10.5402/2012/103892.
- Pergent-Martini, C., G. Pergent, B. Monnier, C.-F. Boudouresque, C. Mori & A. Valette-Sansevin (2021):** Contribution of *Posidonia oceanica* meadows in the context of climate change mitigation in the Mediterranean Sea. *Mar. Environ. Res.*, 165, 105236. <https://doi.org/10.1016/j.marenvres.2020.105236>.
- Perumal, P., U. Arthanari & S.A. Elumalai (2021):** A Review on Pharmacologically Potent Bioactive Metabolites of Seagrass Cymodoceaceae Family. *Curr. Trends Biotechnol. Pharm.*, 15, 436-446.
- Pfeifer, L. & B. Classen B (2020):** The Cell Wall of Seagrasses: Fascinating, Peculiar and a Blank Canvas for Future Research. *Front. Plant Sci.*, 11, 588754. doi: 10.3389/fpls.2020.588754.
- Razgonova, M.P., L.A. Tekutyeva, A.B. Podvolotskaya, V.D. Stepanchikina, A.M. Zakharenko & K. Golokhvast (2022):** *Zostera marina* L.: Supercritical CO₂-Extraction and Mass Spectrometric Characterization of Chemical Constituents Recovered from Seagrass. *Separations*, 9, 182.
- Rees, C.R., J.M. Costin, R.C. Fink, M. McMichael, K.A. Fontaine, S. Isern & S.F. Michael (2008):** In vitro inhibition of dengue virus entry by p-sulfoxy-cinnamic acid and structurally related combinatorial chemistries. *Antivir. Res.*, 80, 135-142.
- Reynolds, L.K., M. Waycott, K.J. McGlathery & R.J. Orth (2016):** Ecosystem Services Returned through Seagrass Restoration. *Restoration Ecology*, 24, 583-588. <https://doi.org/10.1111/rec.12360>.
- Ribas-Taberner, M.d.M., P.M. Mir-Rossello, L. Gil, A. Sureda & X. Capó (2025):** Potential Use of Marine Plants as a Source of Bioactive Compounds. *Molecules*, 30, 485. <https://doi.org/10.3390/molecules30030485>.
- Rotter, A., A. Bacu, M. Barbier, F. Bertoni, A.M. Bones, M.L. Cancela, J. Carlsson, M. F. Carvalho, M. Ceglowska, M.C. Dalay, T. Dailianis, I. Deniz, D. Drakulovic, A. Dubnika, H. Einarsson, A. Erdogan, O.T. Eroldogan, D. Ezra, S. Fazi, R.J. FitzGerald, L.M. Gargan, S.P. Gaudêncio, N. Ivošević DeNardis, D. Joksimović, M. Katarzyte, J. Kotta, M. Mandalakis, I. Matijošyte, H. Mazur-Marzec, A. Massa-Gallucci, M. Mehiri, S.L. Nielsen, L. Novoveská, D. Overlinge, M.E. Portman, K. Pyrc, C. Rebours, T. Reinsch, F. Reyes, B. Rinkevich, J. Robbins, V. Rudovica, J. Sabotič, I. Safarik, S. Talve, D. Tasdemir, X.T. Schneider, O.P. Thomas, A. Torunska-Sitarz, G.C. Varese & M.I. Vasquez (2020):** A New Network for the Advancement of Marine Biotechnology in Europe and Beyond. *Front. Mar. Sci.*, 7, 278. doi: 10.3389/fmars.2020.00278.
- Rotter, A., M. Barbier, F. Bertoni, A.M. Bones, M.L. Cancela, J. Carlsson, M.F. Carvalho, M. Ceglowska, J. Chirivella-Martorell, M. Conk Dalay, M. Cueto, T. Dailianis, I. Deniz, A.R. Díaz-Marrero, D. Drakulovic, A. Dubnika, C. Edwards, H. Einarsson, A. Erdogan, O.T. Eroldogan, D. Ezra, S. Fazi, R.J. FitzGerald, L.M. Gargan, S.P. Gaudêncio, M. Gligora Udovič, N. Ivošević DeNardis, R. Jónsdóttir, M. Katarzyte, K. Klun, J. Kotta, L. Ktari, Z. Ljubešić, L. Lukić Bilela, M. Mandalakis, A. Massa-Gallucci, I. Matijošyte, H. Mazur-Marzec, M. Mehiri, S.L. Nielsen, L. Novoveská, D. Overlinge, G. Perale, P. Ramasamy, C. Rebours, T. Reinsch, F. Reyes, B. Rinkevich, J. Robbins, E. Röttinger, V. Rudovica, J. Sabotič, I. Safarik, S. Talve, D. Tasdemir, X. Theodotou Schneider, O.P. Thomas, A. Torunska-Sitarz, G.C. Varese & M.I. Vasquez (2021):** The Essentials of Marine Biotechnology. *Front. Mar. Sci.* 8:629629. doi: 10.3389/fmars.2021.629629.

- Rotter, A., A. Giannakourou, J.E. Argente García, G.M. Quero, C. Auregan, G. Triantaphyllidis, A. Venetsanopoulou, R. De Carolis, C. Efstratiou, M. Aboal, M.Á.E. Abad, E. Grigalionyte-Bembič, Y. Kotzamanis, M. Kovač, M. Ljubić Čmelar, G.M. Luna, C. Aguilera, F.G. Acien Fernández, J.L. Gómez Pinchetti, S. Manzo, I. Milašinčić, A. Nadarmija, L. Parrella, M. Pinat, E. Roussos, C. Ruel, E. Salvatori, F.J. Sánchez Vázquez, M. Semitiel García, A.F. Skarmeta Gómez, J. Ulčar & C. Chiavetta (2023): Identification of Marine Biotechnology Value Chains with High Potential in the Northern Mediterranean Region. *Marine Drugs*, 21(7), 416. <https://doi.org/10.3390/md21070416>.
- Rotter, A., D. Varamogianni-Mamatsi, A. Zvonar Pöbirk, M. Gosenca Matjaž, M. Cueto, A.R. Díaz-Marrero, R. Jonsdóttir, K. Sveinsdóttir, T.S. Catalá, G. Romano, B.A. Guler, E. Atak, M. Berden Zrimec, D. Bosch, I. Deniz, S. P. Gaudencio, E. Grigalionyte-Bembič, K. Klun, L. Zidar, A. Coll Rius, Š. Baebler, L. Lukić Bilela, B. Rinkevich & M. Mandalakis (2024): Marine cosmetics and the blue bioeconomy: From sourcing to success stories. *iScience*, 27(12), 111339.
- Rudovica, V., A. Rotter, S.P. Gaudêncio, L. Novoveská, F. Akgül, L.K. Akslen-Hoel, D.A.M. Alexandrino, O. Anne, L. Arbidans, M. Atanassova, M. Beldowska, J. Beldowski, A. Bhatnagar, O. Bikovens, V. Bisters, M.F. Carvalho, T.S. Catalá, A. Dubnika, A. Erdogan, L. Ferrans, B.Z. Haznedaroglu, R. Hendroko Setyobudi, B. Graca, I. Grinfelde, W. Hogland, E. Ioannou, Y. Jani, M. Kataržyte, S. Kikionis, K. Klun, J. Kotta, M. Kriipsalu, J. Labidi, L. Lukić Bilela, M. Martínez-Sanz, J. Oliveira, R. Ozola-Davidane, J. Pilecka-Ulcugaceva, K. Pospiskova, C. Rebours, V. Roussis, A. López-Rubio, I. Safarik, F. Schmieder, K. Stankevica, T. Tamm, D. Tasdemir, C. Torres, G.C. Varese, Z. Vincevica-Gaile, I. Zekker & J. Burlakovs (2021): Valorization of Marine Waste: Use of Industrial By-Products and Beach Wrack Towards the Production of High Added-Value Products. *Front. Mar. Sci.*, 8, 723333. doi: 10.3389/fmars.2021.723333.
- Ruiz-Frau, A., S. Gelcich, I.E. Hendriks, C.M. Duarte & N. Marbà (2017): Current State of Seagrass Ecosystem Services: Research and Policy Integration. *Ocean Coast. Manag.*, 149, 107-115. <https://doi.org/10.1016/j.ocecoaman.2017.10.004>.
- Serrano, O., D.I. Gómez-López, L. Sánchez-Valencia, A. Acosta-Chaparro, R. Navas-Camacho, J. González-Corredor, C. Salinas, P. Masque, C.A. Bernal & N. Marbà (2021): Seagrass blue carbon stocks and sequestration rates in the Colombian Caribbean. *Sci Rep* 11, 11067. <https://doi.org/10.1038/s41598-021-90544-5>.
- Sevimli-Gur, C. & O. Yesil-Celiktas (2019): Cytotoxicity screening of supercritical fluid extracted seaweeds and phenylpropanoids. *Mol. Biol. Rep.*, 46, 3691-3699.
- Short, F., T. Carruthers, W. Dennison & M. Waycott (2007): Global seagrass distribution and diversity: A bioregional model. *J. Exp. Mar. Biol. Ecol.*, 350, 3-20.
- Styshova, O.N., A.M. Popov, A.A. Artyukov & A.A. Klimovich (2017): Main constituents of polyphenol complex from seagrasses of the genus *Zostera*, their antidiabetic properties and mechanisms of action. *Exp. Ther. Med.*, 13, 1651-1659.
- Subhashini, P., E. Dilipan, T. Thangaradjou & J. Papenbrock (2013): Bioactive natural products from marine angiosperms: abundance and functions. *Nat. Prod. Bioprospect.*, 3, 129-136. <https://doi.org/10.1007/s13659-013-0043-6>.
- Tarquinio, F., O. Attlan, M.A. Vanderklift, O. Berry & A. Bissett (2021): Distinct Endophytic Bacterial Communities Inhabiting Seagrass Seeds. *Front. Microbiol.*, 12, 703014. doi: 10.3389/fmicb.2021.703014.
- Telesca, L., A. Belluscio, A. Criscoli, G. Ardizzone, E.T. Apostolaki, S. Frascetti, M. Gristina, L. Knittweis, C.S. Martin, G. Pergent, A. Alagna, F. Badalamenti, G. Garofalo, V. Gerakaris, M.L. Pace, C. Pergent-Martini & M. Salomidi (2015). Seagrass meadows (*Posidonia oceanica*) distribution and trajectories of change. *Sci Rep* 5, 12505. <https://doi.org/10.1038/srep12505>.
- Vasarri, M., E. Barletta, M. Stio, M.C. Bergonzi, A. Galli & D. Degl'Innocenti (2023): Ameliorative Effect of *Posidonia oceanica* on High Glucose-Related Stress in Human Hepatoma HepG2 Cells. *Int. J. Mol. Sci.*, 24, 5203. <https://doi.org/10.3390/ijms24065203>.
- Vasarri, M., M. Leri, E. Barletta, M. Ramazzotti, R. Marzocchini, D. Degl'Innocenti (2020): Anti-inflammatory properties of the marine plant *Posidonia oceanica* (L.) Delile. *Journal of Ethnopharmacology*, 247, 112252. <https://doi.org/10.1016/j.jep.2019.112252>.
- Zheng, H., X. Sun, N. Guo & R. Li (2014): New insight on antioxidants and anti-obesity properties of two seagrasses *Thalassia hemprichii* and *Zostera marina*: an integrated molecular docking simulation with in vitro study. *AJMR*, 8(23), 2315-2321. Doi: 10.5897/AJMR2014.6790.