

**IMFM**

INSTITUTE OF MATHEMATICS, PHYSICS AND MECHANICS  
JADRANSKA 19, 1000 LJUBLJANA, SLOVENIA

**Preprint series**

**Vol. 50 (2012), 1178**

ISSN 2232-2094

**COUNTING GRAPHS WITH  
DIFFERENT NUMBERS OF  
SPANNING TREES  
THROUGH THE COUNTING  
OF PRIME PARTITIONS**

Jernej Azarija

Ljubljana, June 28, 2012

# Counting graphs with different numbers of spanning trees through the counting of prime partitions

Jernej Azarija \*

## Abstract

Let  $A_n$  ( $n \geq 1$ ) be the set of all integers  $x$  such that there exists a connected graph on  $n$  vertices with precisely  $x$  spanning trees. In this paper, we show that  $|A_n|$  grows faster than  $\sqrt{n}e^{\frac{2\pi}{\sqrt{3}}\sqrt{n/\log n}}$ . This settles a question of Sedláček posed in [6].

## 1 Introduction

J. Sedláček is regarded as one of the pioneers of Czech graph theory. He devoted much of his work to the study of subjects related to the number of spanning trees  $\tau(G)$  of a graph  $G$ . In [7] he studied the function  $\alpha(n)$  defined as the least number  $k$  for which there exists a graph on  $k$  vertices having precisely  $n$  spanning trees. He showed that for every  $n > 6$ , it holds:

$$\alpha(n) \leq \begin{cases} \frac{n+6}{3} & \text{if } n \equiv 0 \pmod{3}, \\ \frac{n+4}{3} & \text{if } n \equiv 2 \pmod{3}. \end{cases}$$

Azarija and Škrekovski [1] later found out that if  $n > 25$  then:

$$\alpha(n) \leq \begin{cases} \frac{n+4}{3} & \text{if } n \equiv 2 \pmod{3}, \\ \frac{n+9}{4} & \text{otherwise.} \end{cases}$$

Sedláček continued to study quantities related to the function  $\tau$ . In [6] and [8] he considered the set  $B_n^t$  defined as the set of integers so that  $x \in B_n^t$  whenever there is a  $t$ -regular graph on  $n$  vertices with precisely  $x$  spanning trees. He showed that for odd integers  $t \geq 3$ ,  $\lim_{a \rightarrow \infty} |B_{2a}^t| = \infty$  and whenever  $t \geq 4$  is an even integer  $\lim_{a \rightarrow \infty} |B_a^t| = \infty$ . In [6] he also studied a more general set -  $A_n$  defined as a set of numbers such that  $x \in A_n$  whenever there exists a connected graph on  $n$  vertices having precisely  $x$  spanning trees. One could think about  $|A_n|$  as the maximal number of connected graphs on  $n$  vertices with mutually different numbers of spanning trees. Using a simple construction, he has shown that:

$$\lim_{n \rightarrow \infty} \frac{|A_n|}{n} = \infty$$

---

\*Department of Mathematics, University of Ljubljana, Jadranska 21, 1000 Ljubljana, Slovenia, e-mail: jernej.azarija@gmail.com

and remarked: *it is not clear how the fraction  $\frac{|A_n|}{n^2}$  behaves when  $n$  tends to infinity.* In modern terminology, we could write his result as  $|A_n| = \omega(n)$  since  $f(n) = \omega(g(n))$  whenever  $|f(n)| \geq c|g(n)|$  for every  $c > 0$  and  $n > n_0$  for an appropriately chosen  $n_0$ .

In this paper, we extend his work and show that  $|A_n| = \omega(\sqrt{n}e^{\frac{2\pi}{\sqrt{3}}\sqrt{n/\log n}})$ . In order to prove the result we define the graph  $C_{x_1, \dots, x_k}$  as follows. Let  $3 \leq x_1 \leq \dots \leq x_k$  be integers. By  $C_{x_1, \dots, x_k}$  we denote the graph that is obtained after identifying a vertex from the disjoint cycles  $C_{x_1}, \dots, C_{x_k}$ . Since  $C_{x_1}, \dots, C_{x_k}$  are the blocks of  $C_{x_1, \dots, x_k}$ , it follows that

$$\tau(C_{x_1, \dots, x_k}) = \prod_{i=1}^k x_i \quad \text{and} \quad |V(C_{x_1, \dots, x_k})| = \sum_{i=1}^k x_i - k + 1.$$

We also introduce some number theoretical concepts. We say that  $\langle x_1, \dots, x_k \rangle$  is a *partition* of  $n$  with integer *parts*  $1 \leq x_1 \leq \dots \leq x_k$  if  $\sum_{i=1}^k x_i = n$ . The study of partitions covers an extensive part of the research done in combinatorics and number theory. If we denote by  $p(n)$  the number of partitions of  $n$  then, the celebrated theorem of Hardy and Ramanujan [3] states that

$$p(n) \sim \frac{1}{4n\sqrt{3}} e^{\pi\sqrt{\frac{2n}{3}}}$$

or, equivalently

$$\lim_{n \rightarrow \infty} \frac{p(n)}{\frac{1}{4n\sqrt{3}} e^{\pi\sqrt{\frac{2n}{3}}}} = 1.$$

Since then, asymptotics for many types of partition functions have been studied [2]. For the interest of our paper is the function  $p_p(n)$  which we define as the number of partitions of  $n$  into prime parts. In [5] Roth and Szekeres presented a theorem which can be used to derive the following asymptotic relation for  $p_p$

$$p_p(n) \sim e^{\frac{2\pi}{\sqrt{3}}\sqrt{n/\log n}}.$$

Somehow surprising is the fact that if we disallow a constant number of primes as parts, the same asymptotic relation still holds. More specifically, if  $p_{op}(n)$  is the number of partitions into odd primes then

$$p_{op}(n) \sim p_p(n).$$

## 2 Main Result

In this section we prove that  $|A_n| = \omega(\sqrt{n}e^{\frac{2\pi}{\sqrt{3}}\sqrt{n/\log n}})$  by showing that

$$\lim_{n \rightarrow \infty} \frac{|A_n|}{\sqrt{n}e^{\frac{2\pi}{\sqrt{3}}\sqrt{n/\log n}}} = \infty.$$

We do so by establishing a lower bound for the number of partitions whose parts are only odd primes and whose sum is less than or equal to a given number  $n$ .

**Lemma 1.** Let  $P_n$  be the set of all partitions  $\langle x_1, \dots, x_k \rangle$  with  $\sum_{i=1}^k x_i \leq n$  and all the numbers  $x_1, \dots, x_k$  being odd primes. Then, there exists a  $n_0$  such that for all  $n \geq n_0$

$$|P_n| \geq \frac{1}{4} \sqrt{n \log n} e^{\frac{2\pi}{\sqrt{3}} \sqrt{n/\log n}}.$$

*Proof.* Let  $n_1$  be such a positive integer so that for all  $n \geq n_1$

$$p_{op}(n) \geq \frac{1}{2} e^{\frac{2\pi}{\sqrt{3}} \sqrt{n/\log n}}.$$

For  $n \geq n_1$  we then have

$$|P_n| = \sum_{i=3}^n p_{op}(i) \geq \frac{1}{2} \sum_{i=n_1}^n e^{\frac{2\pi}{\sqrt{3}} \sqrt{i/\log i}} \geq \frac{1}{2} \int_{n_1-1}^n e^{\frac{2\pi}{\sqrt{3}} \sqrt{x/\log x}} dx.$$

Showing that  $\int_{n_1-1}^n e^{\frac{2\pi}{\sqrt{3}} \sqrt{x/\log x}} dx \sim \frac{\sqrt{3}}{\pi} \sqrt{n \log n} e^{\frac{2\pi}{\sqrt{3}} \sqrt{n/\log n}}$  would imply the existence of a positive integer  $n_2$  such that

$$\int_{n_1-1}^n e^{\frac{2\pi}{\sqrt{3}} \sqrt{x/\log x}} dx \geq \frac{1}{2} \sqrt{n \log n} e^{\frac{2\pi}{\sqrt{3}} \sqrt{n/\log n}}$$

for all  $n \geq n_2$ . The statement of the theorem would then immediately follow for all  $n \geq n_0$  where  $n_0 = \max\{n_1, n_2\}$ . To prove the last asymptotic identity we observe that it follows from L'Hospital's rule that

$$\lim_{n \rightarrow \infty} \frac{\int_{n_1-1}^n e^{\frac{2\pi}{\sqrt{3}} \sqrt{x/\log x}} dx}{\frac{\sqrt{3}}{\pi} \sqrt{n \log n} e^{\frac{2\pi}{\sqrt{3}} \sqrt{n/\log n}}} = \lim_{n \rightarrow \infty} \frac{e^{\frac{2\pi}{\sqrt{3}} \sqrt{n/\log n}}}{\frac{d}{dn} \left( \frac{\sqrt{3}}{\pi} \sqrt{n \log n} e^{\frac{2\pi}{\sqrt{3}} \sqrt{n/\log n}} \right)} = 1.$$

□

Lemma 1 readily gives an asymptotic lower bound for  $|A_n|$ .

**Theorem 2.**  $|A_n| = \omega\left(\sqrt{n} e^{\frac{2\pi}{\sqrt{3}} \sqrt{n/\log n}}\right)$ .

*Proof.* Let  $P_n$  be defined in the same way as in the statement of Lemma 1. To every partition  $\langle x_1, \dots, x_k \rangle \in P_n$  with sum  $s = \sum_{i=1}^k x_i$  we associate the graph obtained after identifying a vertex from  $C_{x_1, \dots, x_k}$  with a vertex from the disjoint path  $P_{n-s+k}$ . Observe that the resulting graph has precisely  $n$  vertices and  $\prod_{i=1}^k x_i$  spanning trees. Since all the parts in the partitions are primes it follows that any pair of graphs that were obtained from two different partitions in  $P_n$  have a different number of spanning trees. Thus

$$|A_n| \geq |P_n|$$

and therefore from Lemma 1 we know that for large enough  $n$

$$|A_n| \geq \frac{1}{4} \sqrt{n \log n} e^{\frac{2\pi}{\sqrt{3}} \sqrt{n/\log n}}.$$

Since

$$\lim_{n \rightarrow \infty} \frac{\frac{1}{4} \sqrt{n \log n} e^{\frac{2\pi}{\sqrt{3}} \sqrt{n/\log n}}}{\sqrt{n} e^{\frac{2\pi}{\sqrt{3}} \sqrt{n/\log n}}} = \infty$$

it follows from the squeeze theorem that

$$\lim_{n \rightarrow \infty} \frac{|A_n|}{\sqrt{n} e^{\frac{2\pi}{\sqrt{3}}} \sqrt{n/\log n}} = \infty$$

from where the stated claim follows.  $\square$

Observe, that all the graphs constructed in the proof of Theorem 2 contain a cut vertex. Since almost all graphs are 2-connected [4] it is reasonable to expect that there exists a construction of a class  $C_n$  of 2-connected graphs of order  $n$  with mutually different number of spanning trees such that  $|C_n| = \omega(|P_n|)$ . We thus leave it as a further investigation to try to improve the bound on the growth rate of  $|A_n|$ .

### 3 Acknowledgements

The author is thankful to Riste Škrekovski and Martin Raič for fruitful discussions.

### References

- [1] J. Azarija, R. Škrekovski, Euler's idoneal numbers and an inequality concerning minimal graphs with a prescribed number of spanning trees, manuscript, 2011.
- [2] P. Flajolet, R. Sedgewick, *Analytic Combinatorics*, Cambridge University Press 2009, p. 574.
- [3] G. H. Hardy, S. Ramanujan, Asymptotic formulæ in combinatory analysis, *Proc. London Math. Soc.* (2) **17** (1918) 75–115.
- [4] F. Harary, E.M Palmer, *Graphical Enumeration* Academic Press, New York, 1973.
- [5] K. F. Roth, G. Szekeres, Some asymptotic formulae in the theory of partitions, *Quarterly Journal of Mathematics*, Oxford Series **5** (1954) 241–259.
- [6] J. Sedláček, On the number of spanning trees of finite graphs, *Čas. Pro. Pěst Mat.*, Vol. **94** (1969) 217–221.
- [7] J. Sedláček, On the minimal graph with a given number of spanning trees, *Canad. Math. Bull.* **13** (1970) 515–517.
- [8] J. Sedláček, Regular graphs and their spanning trees, *Čas. Pro. Pěst Mat.*, Vol. **95** (1970) 420–426.