

REGULAR GRAPHS ARE 'DIFFICULT' FOR COLOURING

Janez Žerovnik

University of Maribor, Faculty of Technical Sciences,
Smetanova 17, Maribor,
Slovenia

Keywords: graph colouring, decision problem, NP-completeness, regular graphs

Edited by: Rudi Murn

Received: December 19, 1992

Revised: February 16, 1993

Accepted: March 1, 1993

Abstract: Let k be 3 or 4. In this two cases we prove that the decision problem of k -colourability when restricted to Δ -regular graphs is NP-complete for any $\Delta \geq k + 1$.

1 Introduction

In this note we consider the time complexity of the decision problem of (vertex) k -colourability restricted to regular graphs.

It is known that 'almost all k -colourable graphs are easy to colour', namely the proportion of 'difficult' graphs for the usual backtrack algorithm vanishes with growing problem size [9]. Knowing this it is not surprising that there are algorithms with average polynomial time complexity [1], when average is taken over all graphs and even when the average is taken over all 3-colourable graphs with a given number of vertices[5].

If $P \neq NP$, then for every algorithm there has to be a class of 'counterexamples', i.e. graphs on which the algorithm either has superpolynomial time complexity or it fails to produce a correct answer.

For example, Petford and Welsh noticed that one of the situations in which the 3-colourable graphs were not efficiently coloured by their randomised algorithm is when graphs are approximately regular of a low vertex degree [8]. Similarly, approximately regular graphs of a relative low vertex degree are 'difficult' also for the k -colouring generalisation of their algorithm [10]. Petford and Welsh conjectured that 'dense' graphs are easy. Indeed, Edwards showed that, when restricted to class of graphs with lowest vertex degree $\delta > \alpha n$ for arbitrary $\alpha > 0$, the decision problem of 3-colouring is polynomial [4].

This may be understood that the 'difficult' graphs are likely to be found among 'sparse' graphs. It is known that the problem of 3-

colouring is NP-complete (even) when restricted to graphs of maximal vertex degree 4 [6]. Here we show that the problem can be further 'simplified', proving that the decision problem of 3-colouring is NP-complete when restricted to Δ -regular graphs (for $\Delta \geq 4$). We also show that the decision problem of 4-colouring is NP-complete when restricted to Δ -regular graphs (for $\Delta \geq 5$).

We assume that the reader is familiar with some standard definitions of graph theory and of computational complexity theory (given, for example, in [2] and [7]).

2 3-colourability of 4-regular graphs is NP-complete

Let us define the problem $\Pi(k, \Delta)$ as follows:

Input: Δ -regular graph G

Question: Is G k -colourable?

Lemma 1 For any graph G there is a graph G' with no vertex of degree 1 or 2 such that:

G is 3-colourable iff G' is 3-colourable

Remark: G' in the Lemma is either a graph with minimal vertex degree ≥ 3 or the empty graph, which is the case when G is, for example, a cycle. If G' is empty, it is trivially 3-colourable, and from the proof of the Lemma 1 it follows that also G is 3-colourable.

Proof (of Lemma 1): It is easy to see that the following assertions are true:

- (a) If there is a vertex $v \in V(G)$ with degree 1, then G is 3-colourable if and only if the





Figure 1: Vertices of degree 1 or 2 may be omitted

Figure 2: G is 3-colourable iff G' is 3-colourable

induced graph on $V \setminus \{v\}$ is 3-colourable (see Fig. 1(a)).

- (b) If there is a vertex $v \in V(G)$ with degree 2, then G is 3-colourable if and only if the induced graph on $V \setminus \{v\}$ is 3-colourable (see Fig. 1(b)).

In this way we can reduce any graph G to a graph G' with minimal vertex degree at least 3 which is 3-colourable exactly when G is. **Q.E.D.**

Remark: Extracting G' successively using (a) and (b) can clearly be done efficiently.

Remark: It is obvious that maximal vertex degree of the graph G' is not greater than maximal vertex degree of the original graph G .

Lemma 2 G is 3-colourable iff G' is 3-colourable

where G' is a graph, obtained from G by the construction given in Fig. 2.

The construction, given in Fig. 2, can be done as follows. Take two sets, say M and N , of three vertices each. Connect every pair x, y ; $x \in M$ and $y \in N$. Add two vertices, say u and v and connect u to all the vertices of N and v to all the vertices of M . Now choose arbitrary pair of distinct vertices of G , say w and z , and connect u with w and v with z to get the graph G' .

Proof: Since the graph H is bipartite, it is easy to see that 3-colouring of arbitrary graph G (on Fig. 2) can be extended to 3-colouring of graph G' . On the other hand, since G is subgraph of G' , G is 3-colourable if G' is. **Q.E.D.**

Now we shall prove

Lemma 3 The problem $\Pi(3, 4)$ is NP-complete.

Proof: We will reduce the problem of 3-colourability of graphs with vertex degree at most 4 (which is known to be NP-complete [6]) to the problem $\Pi(3, 4)$.

Let G be arbitrary graph with maximal degree $\Delta \leq 4$. By Lemma 1 there is a graph G_1 (which has at most as many vertices as G) and G_1 is 3-colourable exactly when G is 3-colourable. If G_1 is empty, then we know that G is 3-colourable.

Now consider the case when G_1 is nonempty. By construction, G_1 is a graph with vertex degrees 3 and 4. Since the sum of all the vertex degrees is twice the number of edges ($\sum_{v \in V} \delta_v = 2|E|$), the number of vertices with degree 3 must be even.

Now couple vertices of degree 3 in G_1 arbitrarily. Connect a copy of the graph H to each couple of vertices of degree 3, as defined in Fig. 2. By Lemma 2, this construction gives a graph G_2 which is 3-colourable exactly when G_1 is 3-colourable. **Q.E.D.**

Remark: Graph H has 8 vertices. Since we added at most $\frac{|V|}{2} 8$ new vertices, the resulting graph G_2 has at most a constant factor more vertices than G_1 .

Remark: The construction can clearly be done efficiently.

Thus 3-colourability of 4-regular graphs is NP-complete. Now we reduce the problem of 3-colourability of Δ -regular graphs to the same problem on $\Delta + 1$ -regular graphs.




Figure 3: Joining two copies of a Δ -regular graph we get a $\Delta + 1$ -regular graph

3 3-colourability of Δ -regular graphs is NP-complete

Lemma 4 $\Pi(3, \Delta) \propto \Pi(3, \Delta + 1)$

Proof: Let G be arbitrary Δ -regular graph. Now we give a construction of a graph G' .

Take two copies of G , $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$. Denote with $f : G \rightarrow G_1$ and $g : G \rightarrow G_2$ the corresponding isomorphisms.

Graph $G' = (V', E')$ is defined with: $V' = V_1 \cup V_2$ and $E' = E_1 \cup E_2 \cup \{\{f(v), g(v)\} \mid v \in V\}$ (see Fig. 3).

If G' is 3-colourable, then also G is 3-colourable, since it is isomorphic to a subgraph in G' .

On the other hand, if we have a 3-colouring b of G , it is easy to construct a 3-colouring b' of G' , for example with a 'shift' of colours: $b'(f(v)) = b(v)$ and $b'(g(v)) = (b(v) \bmod 3) + 1$.

Since for any Δ size blow up is a constant factor, the assertion of the Lemma follows. **Q.E.D.**

By induction, from Lemma 3 and Lemma 4 we have:

Proposition 1 *The decision problem of 3-colouring restricted to Δ -regular graphs $\Pi(3, \Delta)$ is NP-complete for $\Delta \geq 4$.*

It is known that for graphs of maximal vertex degree 3 the problem is polynomial [6]. Hence we know for all problems $\Pi(3, \Delta)$ whether they are polynomial or NP-complete.

4 4-colourability of Regular Graphs

Here we discuss an attempt to generalise the proposition 1 on the problem of k -colouring. With analogous proof as for the case of 3-colourings we prove a proposition for 4-colouring, while for $k > 4$ the time complexity of the decision problem of k -colouring of Δ -regular graphs remains open for some Δ .

Two of the previous lemmas are easily generalised:

Lemma 5 *Let G' be any subgraph of G obtained by the following process: if there is a vertex of degree less than k , delete it. Graph G is k -colourable if and only if graph G' is k -colourable.*

Proof: Assume we coloured the graph G' with k -colours. It is easy to see that there is algorithm, which properly extends the proper colouring of G' to a proper colouring of G . (Take, for example, vertices of G in opposite order as they were deleted from G . When a vertex was deleted, it had less than k neighbours, therefore there is at least one free colour for it.) **Q.E.D.**

Lemma 6 *For any graph G with vertex degrees k and $k + 1$ there is a $k + 1$ -regular graph G' , such that:*

G is k -colourable iff G' is k -colourable




Figure 4: G is 4-colourable iff G' is 4-colourable

Proof: If there are at least two vertices of degree k in G , then we add a copy of graph H . For given k the graph H is defined as follows. Take a complete bipartite graph $K_{k,k}$. Add two vertices

and connect one vertex with all the vertices of one independent set of the $K_{k,k}$ and the other vertex with the second independent set of the $K_{k,k}$ (for the case $k = 4$ see Fig. 4). In this way we reduce the number of vertices of degree k by two.

If there is only one vertex of degree k in G , then we construct a new graph as follows: Take two copies of G , connect the two vertices of degree k with an edge. The resulting graph is obviously $k + 1$ -regular and it is easy to see that it is k -colourable exactly when G is k -colourable. **Q.E.D.**

For a proof of a generalization of the proposition 1 we need a lemma of the following type: decision problem of k -colouring on arbitrary graph can be reduced to the same problem on a graph of maximal vertex degree $k + 1$.

In the proof of the proposition for 3-colouring we used the result of Garey, Johnson and Stockmayer. Here we give the idea of a proof for $k = 4$. We were not able to generalise the idea for $k > 4$.

Lemma 7 *The decision problem of 4-colouring of graphs of vertex degree ≤ 5 is NP-complete.*




Figure 5: Graph for substituting vertices of degree 6

Proof (outline): The key of the proof is the idea of how to substitute vertices of large degree with a graph of small enough maximal vertex degree and with property that any 4-colouring of the resulting graph G' defines a 4-colouring of the original graph G . Such graphs are given in Figures 5, 6 and 7. The graphs in Fig. 5 and Fig. 6 are used for substituting vertices of degrees 6 and 7, respectively. For vertices of larger degrees, a longer chain is used, as indicated on Fig. 7. The graphs given have the property, that in any proper 4-colouring all the vertices with 'free edges' have to be coloured with the same colour. (This colour




Figure 6: Graph for substituting vertices of degree 7

can be then assigned to the substituted vertex in the original graph. The other vertices of G can then be assigned the same colours as they had in the 4-colouring of G' .) We omit the details. **Q.E.D.**

With a straightforward generalization of the proof of Lemma 4 we have also:

Lemma 8 $\Pi(k, \Delta) \propto \Pi(k, \Delta + 1)$

Therefore:

Proposition 2 *The decision problem of 4-colouring of Δ -regular graphs $\Pi(4, \Delta)$ is NP-complete for any $\Delta \geq 5$.*


Again, because of the theorem of Brooks [3], the problem $\Pi(4, \Delta)$ has polynomial time complexity for $\Delta \leq 4$. Thus for all the problems $\Pi(4, \Delta)$ we know whether they are polynomial or NP-complete. Let us conclude with a couple of conjectures. Since we were unable to generalise the Lemma 7 we state

Conjecture 1 *The decision problem of k -colouring of graphs with vertex degree $\leq k + 1$ is NP-complete.*

If the first conjecture was true, then we would have a nice classification of time complexity for all the problems $\Pi(k, \Delta)$.

Conjecture 2 *For any $k > 2$, $\Delta > 2$ the decision problem of k -colourability of Δ -regular graphs $\Pi(k, \Delta)$ is NP-complete if $\Delta > k$ and is polynomial otherwise.*

Let us conclude with a simple consequence of the proposition. Assume we have an algorithm

Figure 7: Graph for substituting vertices of degree > 5

$\mathcal{A}$ for 3-colouring and we want to characterise graphs, for which the algorithm does not provide the correct solution in polynomial time. If $P \neq NP$ then for any algorithm $\mathcal{A}$ for each $\Delta \geq 4$ there exists an infinite family $F(\mathcal{A}, \Delta)$ of Δ -regular graphs such that the algorithm $\mathcal{A}$ has superpolynomial complexity on each family $F(\mathcal{A}, \Delta)$. If this were not the case for some Δ then $\mathcal{A}$ would be a polynomial algorithm for 3-colouring of Δ -regular graphs, which would imply $P=NP$!

Acknowledgement. The contents of the paper is based on a part of author's dissertation submitted to University of Ljubljana under supervision of Tomaž Pisanski. Thanks are also due to the anonymous referees, who helped to improve the quality of presentation considerably.

References

- [1] E.A. Bender, H.S. Wilf, A Theoretical Analysis of Backtracking in the Graph Coloring Problem, *Journal of Algorithms* **6** (1985) 175-282.
- [2] J.A. Bondy, U.S.R. Murty, *Graph Theory with Applications*, Amer. Elsevier, New York, 1976.
- [3] R.L. Brooks, On Coloring the Nodes of a Network, *Proc. Camb. Phil. Soc* **37** (1941) 194-197.
- [4] K. Edwards, The Complexity of Colouring Problems on Dense Graphs, *Theoretical Computer Science* **43**(1986) 123-147.
- [5] J.A. Ellis, P.M. Lepolesa, A Las Vegas Graph Colouring Algorithm, *The Computer Journal* **32** (1989) 474-476.
- [6] M.R. Garey, D.S. Johnson, L. Stockmeyer, Some Simplified NP-complete Graph Problems, *Theoretical Computer Science* **1** (1976) 237-267.
- [7] M.R. Garey, D.S. Johnson, *Computers and Intractability*, W.H. Freeman and Co., San Francisco, 1979.
- [8] A.D. Petford, D.J.A. Welsh, A Randomised 3-Colouring Algorithm, *Discrete Math.*, **74** (1989) 253-261.
- [9] J.S. Turner, Almost All k -Colorable Graphs Are Easy to Color, *Journal of Algorithms* **9** (1988) 63-82.
- [10] J. Žerovnik, A Randomized Algorithm for k -colorability, *Discrete Mathematics* (to appear).